wwPDB EM Validation Summary Report (i)

Nov 9, 2026 – 02:26 PM EST

PDB ID	:	10BG / pdb_000010bg
EMDB ID	:	EMD-75041
Title	:	SARS-CoV-2 Spike in complex with PVI.V6-4 Fab
Authors	:	Jaiswal, D.; Altomare, C.; Adelsberg, D.; Bajic, G.
Deposited on	:	2026-01-09
Resolution	:	4.00 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:


EMDB validation analysis	:	0.0.1.dev133
Mogul	:	2022.3.0, CSD as543be (2022)
MolProbity	:	4-5-2 with Phenix2.2
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics	:	202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 4.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	7587 (3.50 - 4.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	125	18%, 90%, 10%
2	L	114	18%, 89%, 10%
3	S	1195	6%, 16%, 84%

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 3356 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PVI.V6-4 Fab heavy chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	H	124	Total	C	N	O	S	0	0
			940	595	158	182	5		

- Molecule 2 is a protein called PVI.V6-4 Fab light chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	L	113	Total	C	N	O	S	0	0
			877	551	145	178	3		


- Molecule 3 is a protein called Spike glycoprotein.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	S	193	Total	C	N	O	S	0	0
			1525	975	255	287	8		

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
S	682	GLY	ARG	engineered mutation	UNP P0DTC2
S	683	SER	ARG	engineered mutation	UNP P0DTC2
S	685	SER	ARG	engineered mutation	UNP P0DTC2
S	817	PRO	PHE	engineered mutation	UNP P0DTC2
S	892	PRO	ALA	engineered mutation	UNP P0DTC2
S	899	PRO	ALA	engineered mutation	UNP P0DTC2
S	942	PRO	ALA	engineered mutation	UNP P0DTC2
S	986	PRO	LYS	engineered mutation	UNP P0DTC2
S	987	PRO	VAL	engineered mutation	UNP P0DTC2

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				AltConf
			Total	C	N	O	
4	S	1	14	8	1	5	0


3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: PVI.V6-4 Fab heavy chain




• Molecule 2: PVI.V6-4 Fab light chain



• Molecule 3: Spike glycoprotein




Q498 P499 T500 N501 G504 S514 L518 H519 A520 P521 A522 T523 V524 C525 G526
PRO LYS LYS SER THR ASN LEU VAL LYS ASN LYS CYS VAL ASN PHE ASN PHE ASN GLY LEU THR GLY THR GLY VAL LEU THR GLU SER ASN LYS LYS PHE LEU PRO PHE GLN GLN PHE GLY ARG ASP
ILE ALA ASP THR THR ASP ALA VAL ARG ASP PRO GLN THR LEU GLU ILE LEU ASP ILE THR PRO CYS SER PHE GLY GLY VAL SER VAL ILE THR PRO GLY THR ASN THR SER ASN GLN VAL ALA VAL LEU TYR GLN ASP VAL ASN CYS THR GLU VAL PRO VAL ALA ILE HIS ALA ASP GLN
LEU THR PRO THR TRP ARG VAL TYR SER THR GLY SER ASN VAL PHE GLN THR ARG ALA GLY CYS LEU ILE GLY ALA GLU HIS VAL ASN ASN SER TYR GLU CYS ASP ILE PRO ILE GLY ALA GLY ILE CYS ALA SER TYR GLN THR GLN THR ASN SER PRO GLY SER ALA SER SER VAL ALA
SER GLN SER ILE ILE ALA TYR THR MET SER LEU GLY ALA GLU ASN SER VAL ALA TYR SER ASN ASN SER ILE ALA ILE PRO THR ASN PHE THR ILE SER VAL THR THR GLU ILE LEU PRO VAL SER MET THR LYS THR SER VAL ASP CYS THR MET TYR ILE CYS GLY ASP SER THR GLU
CYS SER ASN LEU LEU LEU GLN TYR GLY SER PHE CYS THR GLN LEU ASN ARG ALA LEU THR GLY ILE ALA VAL GLU GLN ASP LYS ASN THR GLN GLU VAL PHE ALA GLN VAL LYS GLN ILE TYR LYS THR PRO PRO ILE LYS ASP PHE GLY GLY PHE ASN PHE SER GLN ILE LEU PRO ASP
PRO SER LYS PRO SER LYS ARG SER PRO ILE GLU ASP LEU LEU PHE ASN LYS VAL THR LEU ALA ASP ALA GLY PHE ILE LYS GLN TYR GLY ASP CYS LEU GLY ASP ILE ALA ALA ARG ASP LEU ILE CYS ALA GLN LYS PHE ASN GLY LEU LEU THR VAL LEU PRO PRO LEU LEU THR ASP GLU
MET ILE ALA GLN TYR THR SER ALA LEU LEU LEU ALA GLY THR ILE THR SER GLY TRP THR PHE GLY ALA GLY PRO ALA LEU GLN LEU ILE PRO PHE PRO MET GLN MET ALA TYR ARG PHE ASN GLY ILE GLY VAL THR GLN ASN VAL LEU TYR GLU ASN GLN LYS LEU ILE ALA ASN GLN PHE ASN
SER ALA ILE GLY LYS ILE GLN ASP SER LEU SER SER SER THR PRO SER ALA LEU GLY LYS LEU GLN ASP VAL VAL ASN GLN ASN ALA ALA GLN ALA LEU ASN THR LEU VAL LYS GLN LEU SER SER ASN PHE GLY ALA ILE SER SER VAL LEU LEU ASN ASP ILE LEU PRO GLU PRO GLU
ALA GLU VAL GLN ILE ASP ARG LEU ILE THR GLY ARG LEU GLN SER LEU GLN THR TYR VAL THR GLN GLN LEU ILE ARG ALA ALA GLU ILE ARG ALA SER ALA ASN LEU ALA ALA THR LYS MET SER GLU CYS VAL LEU GLY GLN SER LYS ARG VAL ALA ASP PHE CYS GLY LYS GLY TYR HIS
LEU MET SER PHE PRO GLN SER ALA PRO HIS GLY VAL VAL PHE LEU HIS VAL THR TYR VAL PRO ALA GLU GLU LYS ASN PHE THR THR ALA PRO ALA ILE CYS HIS ASP GLY LYS ALA HIS PHE PRO ARG GLU CYS VAL GLY VAL PHE VAL SER ASN GLY THR HIS TRP PHE VAL THR GLN ARG ASN
PHE TYR GLU PRO GLN ILE ILE THR THR ASP ASN GLY ASN CYS ASP VAL VAL ILE GLY ILE VAL ASN ASN THR VAL TYR ASP PRO LEU GLN PRO GLU LEU ASP SER PHE LYS GLU GLU LEU ASP LYS TYR PHE LYS ASN HIS THR SER PRO ASP VAL ASP LEU GLY ASP
ILE SER GLY ILE ASN ALA SER VAL VAL ASN ILE GLN LYS GLU ILE ASP ARG LEU ASN GLU VAL ALA LYS ASN LEU ASN GLU SER LEU ILE ASP LEU GLN GLU LEU GLY LYS TYR GLU GLN

4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	1489246	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING ONLY	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/Å^2$)	52.75	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	1500	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	2.539	Depositor
Minimum map value	-0.002	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.008	Depositor
Recommended contour level	0.2	Depositor
Map size (Å)	447.616, 447.616, 447.616	wwPDB
Map dimensions	416, 416, 416	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.076, 1.076, 1.076	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|$ >5	RMSZ	#$\|Z\|$ >5
1	H	0.22	0/959	0.57	0/1297
2	L	0.27	0/898	0.72	2/1224 (0.2%)
3	S	0.19	0/1566	0.47	0/2130
All	All	0.22	0/3423	0.57	2/4651 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	L	6	GLN	CA-C-N	7.02	130.87	120.83
2	L	6	GLN	C-N-CA	7.02	130.87	120.83

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	940	0	924	7	0
2	L	877	0	840	5	0
3	S	1525	0	1442	4	0
4	S	14	0	13	0	0
All	All	3356	0	3219	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 15 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:39:LEU:HB3	2:L:57:ALA:HB2	1.82	0.60
2:L:53:LEU:HB3	2:L:54:ILE:HD12	1.90	0.53
3:S:396:TYR:HB2	3:S:514:SER:HB3	1.91	0.53
1:H:103:TYR:HB3	3:S:417:LYS:HG2	1.92	0.51
3:S:358:ILE:HB	3:S:395:VAL:HB	1.93	0.51

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	122/125 (98%)	111 (91%)	11 (9%)	0	100	100
2	L	111/114 (97%)	97 (87%)	14 (13%)	0	100	100
3	S	189/1195 (16%)	180 (95%)	9 (5%)	0	100	100
All	All	422/1434 (29%)	388 (92%)	34 (8%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	99/100 (99%)	99 (100%)	0	100	100
2	L	99/100 (99%)	99 (100%)	0	100	100
3	S	166/1043 (16%)	166 (100%)	0	100	100
All	All	364/1243 (29%)	364 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	H	39	GLN
3	S	493	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	NAG	S	1301	3	14,14,15	0.77	0	17,19,21	1.32	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	S	1301	3	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
4	S	1301	NAG	C1-O5-C5	3.30	116.61	112.19

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand NAG S 1301




Bond lengths



Bond angles



Torsions



Rings

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.


6 Map visualisation

This section contains visualisations of the EMDB entry EMD-75041. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections


6.1.1 Primary map



X



Y




Z


6.1.2 Raw map



X



Y



Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)


6.2.1 Primary map



X Index: 208



Y Index: 208




Z Index: 208


6.2.2 Raw map



X Index: 208



Y Index: 208




Z Index: 208


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)


6.3.1 Primary map



X Index: 197




Y Index: 206




Z Index: 182


6.3.2 Raw map



X Index: 197



Y Index: 210




Z Index: 415

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) (i)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views (i)

6.5.1 Primary map



X Y Z

The images above show the 3D surface view of the map at the recommended contour level 0.2. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



X Y Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate (i)



The volume at the recommended contour level is 17 nm^3; this corresponds to an approximate mass of 15 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum (i)




*Reported resolution corresponds to spatial frequency of 0.250 $Å^{-1}$

8 Fourier-Shell correlation

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC



*Reported resolution corresponds to spatial frequency of 0.250 Å^{-1}

8.2 Resolution estimates (i)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	4.00	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	5.09	9.17	6.35

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 5.09 differs from the reported value 4.0 by more than 10 %


9 Map-model fit

This section contains information regarding the fit between EMDB map EMD-75041 and PDB model 10BG. Per-residue inclusion information can be found in section 3 on page 5.


9.1 Map-model overlay



X




Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.2 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.


9.2 Q-score mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.2).

9.4 Atom inclusion (i)



At the recommended contour level, 67% of all backbone atoms, 55% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.2) and Q-score for the entire model and for each chain.



Chain	Atom inclusion	Q-score
All	0.5490	0.2790
H	0.5950	0.3390
L	0.5940	0.3110
S	0.4950	0.2240