Full wwPDB EM Validation Report ⓘ

Nov 12, 2026 – 12:17 PM EST

PDB ID	:	40BG / pdb_000040bg
EMDB ID	:	EMD-79944
Title	:	MC4R co-expressed with a heterotrimeric Gs protein and the stabilizing antibody fragment scFv16, in complex with setmelanotide.
Authors	:	Catalano, C.; Scapin, G.; Muccini, A.J.; Senko, S.; Poweleit, N.; Luksic, W.R.; Fisher, B.; Byrne, S.; Haineault, J.; Fencil, K.; Mishra, A.; Gottler, L.
Deposited on	:	2026-11-02
Resolution	:	2.60 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references ⓘ) were used in the production of this report:


EMDB validation analysis	:	0.0.1.dev133
Mogul	:	2022.3.0, CSD as543be (2022)
MolProbity	:	4-5-2 with Phenix2.2
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics	:	202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	8728 (2.10 - 3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	360	82% 12% 6%
2	G	71	11% 75% 24%
3	R	350	63% 11% 25%
4	C	10	30% 60% 10% 30%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	A	373	
6	S	307	

2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 8694 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	B	337	Total	C	N	O	S	0	0
			2579	1594	465	499	21		

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-19	MET	-	initiating methionine	UNP P54311
B	-18	GLY	-	expression tag	UNP P54311
B	-17	SER	-	expression tag	UNP P54311
B	-16	HIS	-	expression tag	UNP P54311
B	-15	HIS	-	expression tag	UNP P54311
B	-14	HIS	-	expression tag	UNP P54311
B	-13	HIS	-	expression tag	UNP P54311
B	-12	HIS	-	expression tag	UNP P54311
B	-11	HIS	-	expression tag	UNP P54311
B	-10	LEU	-	expression tag	UNP P54311
B	-9	GLU	-	expression tag	UNP P54311
B	-8	VAL	-	expression tag	UNP P54311
B	-7	LEU	-	expression tag	UNP P54311
B	-6	PHE	-	expression tag	UNP P54311
B	-5	GLN	-	expression tag	UNP P54311
B	-4	GLY	-	expression tag	UNP P54311
B	-3	PRO	-	expression tag	UNP P54311
B	-2	GLY	-	expression tag	UNP P54311
B	-1	SER	-	expression tag	UNP P54311
B	0	SER	-	expression tag	UNP P54311
B	1	GLY	-	expression tag	UNP P54311

- Molecule 2 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	G	54	Total	C	N	O	S	0	0
			414	260	73	78	3		

- Molecule 3 is a protein called Melanocortin receptor 4.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	R	261	Total	C	N	O	S	0	0
			2020	1346	313	338	23		

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
R	333	LEU	-	expression tag	UNP P32245
R	334	GLU	-	expression tag	UNP P32245
R	335	VAL	-	expression tag	UNP P32245
R	336	LEU	-	expression tag	UNP P32245
R	337	PHE	-	expression tag	UNP P32245
R	338	GLN	-	expression tag	UNP P32245
R	339	GLY	-	expression tag	UNP P32245
R	340	PRO	-	expression tag	UNP P32245
R	341	SER	-	expression tag	UNP P32245
R	342	GLY	-	expression tag	UNP P32245
R	343	SER	-	expression tag	UNP P32245
R	344	GLY	-	expression tag	UNP P32245
R	345	HIS	-	expression tag	UNP P32245
R	346	HIS	-	expression tag	UNP P32245
R	347	HIS	-	expression tag	UNP P32245
R	348	HIS	-	expression tag	UNP P32245
R	349	HIS	-	expression tag	UNP P32245
R	350	HIS	-	expression tag	UNP P32245

- Molecule 4 is a protein called Setmelanotide.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	C	10	Total	C	N	O	S	0	1
			78	49	18	9	2		

- Molecule 5 is a protein called Guanine nucleotide-binding protein G(s) subunit alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	A	226	Total	C	N	O	S	0	0
			1837	1163	330	336	8		

- Molecule 6 is a protein called Stabilizing antibody fragment scFv16.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	S	230	Total	C	N	O	S	0	0
			1759	1118	292	339	10		

- Molecule 7 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		AltConf
7	R	1	Total	Ca	0
			1	1	

- Molecule 8 is water.


Mol	Chain	Residues	Atoms		AltConf
8	R	3	Total	O	0
			3	3	
8	C	3	Total	O	0
			3	3	


3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1

Chain B:






• Molecule 2: Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2

Chain G:






• Molecule 3: Melanocortin receptor 4

Chain R:






• Molecule 4: Setmelanotide

Chain C:






• Molecule 5: Guanine nucleotide-binding protein G(s) subunit alpha

Chain A:






• Molecule 6: Stabilizing antibody fragment scFv16

Chain S:





4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	226360	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/Å^2$)	50.85	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	1.276	Depositor
Minimum map value	-0.801	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.017	Depositor
Recommended contour level	0.112	Depositor
Map size (Å)	318.72, 318.72, 318.72	wwPDB
Map dimensions	384, 384, 384	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.83, 0.83, 0.83	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ACE, DAL, NH2, CA, DPN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| > 5$	RMSZ	$\#\|Z\| > 5$
1	B	0.12	0/2626	0.32	0/3562
2	G	0.09	0/420	0.27	0/568
3	R	0.14	0/2061	0.34	0/2805
4	C	0.70	0/59	0.93	0/75
5	A	0.14	0/1871	0.28	0/2518
6	S	0.10	0/1803	0.28	0/2447
All	All	0.13	0/8840	0.32	0/11975

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
4	C	0	2
5	A	0	1
All	All	0	3

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
5	A	245	ARG	Sidechain
4	C	2	ARG	Sidechain
4	C	7	ARG	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2579	0	2484	25	0
2	G	414	0	420	1	0
3	R	2020	0	2115	29	0
4	C	78	0	66	11	0
5	A	1837	0	1812	13	0
6	S	1759	0	1688	9	0
7	R	1	0	0	0	0
8	C	3	0	0	1	0
8	R	3	0	0	0	0
All	All	8694	0	8585	80	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:C:2:ARG:HG2	4:C:3:CYS:H	1.13	1.08
3:R:188:SER:OG	4:C:7:ARG:HD3	1.73	0.88
4:C:2:ARG:HG2	4:C:3:CYS:N	1.93	0.82
3:R:185:ILE:HD12	4:C:7:ARG:HE	1.46	0.79
4:C:2:ARG:CG	4:C:3:CYS:H	1.94	0.77
3:R:78:PRO:HB3	3:R:166:VAL:HG21	1.70	0.72
5:A:35:ARG:HB3	5:A:202:PHE:HE2	1.56	0.71
5:A:53:MET:HG3	5:A:201:MET:HE1	1.75	0.69
3:R:69:ILE:HA	3:R:75:LEU:HD11	1.74	0.68
5:A:53:MET:HA	5:A:352:ILE:HD11	1.76	0.66
1:B:61:MET:HE2	1:B:70:LEU:HD11	1.78	0.65
3:R:141:LEU:HD11	3:R:207:LEU:HD23	1.81	0.62
3:R:185:ILE:CD1	4:C:7:ARG:HE	2.13	0.61
6:S:202:ARG:HH12	6:S:220:GLU:HG3	1.66	0.61
1:B:158:VAL:HG12	1:B:190:LEU:HD21	1.83	0.61
3:R:185:ILE:HD12	4:C:7:ARG:NE	2.15	0.61
3:R:200:MET:HE2	3:R:204:MET:HE3	1.83	0.59
3:R:188:SER:HG	4:C:7:ARG:HD3	1.67	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:271:CYS:HB2	1:B:290:ASP:HB2	1.87	0.57
5:A:232:SER:CB	5:A:245:ARG:HD2	2.35	0.57
3:R:300:LEU:HD23	3:R:304:LEU:HD12	1.88	0.55
3:R:257:CYS:HB3	3:R:294:ASN:HB2	1.90	0.54
1:B:51:LEU:HB2	1:B:336:LEU:HB2	1.91	0.53
1:B:61:MET:HE3	1:B:328:ALA:HB3	1.90	0.52
6:S:22:CYS:HB3	6:S:79:LEU:HB3	1.92	0.52
6:S:91:THR:HG23	6:S:118:THR:HA	1.91	0.52
5:A:232:SER:OG	5:A:245:ARG:HD2	2.10	0.52
1:B:112:VAL:HG13	1:B:126:LEU:HD11	1.91	0.51
4:C:2:ARG:NH1	8:C:101:HOH:O	2.43	0.50
6:S:200:PRO:HB2	6:S:202:ARG:HG2	1.91	0.50
1:B:187:VAL:HA	1:B:203:ALA:HA	1.93	0.50
1:B:45:MET:HB3	1:B:339:TRP:HB3	1.92	0.49
5:A:187:ILE:HD11	5:A:202:PHE:HB3	1.95	0.49
3:R:73:LYS:HA	3:R:76:HIS:CD2	2.47	0.49
5:A:232:SER:HB2	5:A:245:ARG:HD2	1.95	0.49
1:B:79:LEU:HB2	1:B:93:ILE:HB	1.94	0.48
1:B:49:ARG:HB2	1:B:338:ILE:HB	1.95	0.48
5:A:49:ILE:O	5:A:53:MET:HG2	2.13	0.47
1:B:121:CYS:HB3	1:B:139:LEU:HD12	1.96	0.47
3:R:50:VAL:O	3:R:54:LEU:HD23	2.14	0.47
3:R:164:LYS:O	3:R:168:ILE:HG13	2.15	0.47
6:S:174:LEU:HD22	6:S:212:PHE:CG	2.49	0.47
3:R:220:ARG:HH12	3:R:240:ASN:N	2.13	0.47
3:R:146:ASP:O	3:R:150:THR:HG23	2.16	0.46
5:A:230:SER:HB3	5:A:271:LEU:HD12	1.97	0.46
3:R:285:ASN:O	3:R:289:ILE:HG12	2.16	0.45
5:A:258:ASN:HA	5:A:263:ARG:HH22	1.79	0.45
3:R:257:CYS:CB	3:R:294:ASN:HB2	2.47	0.45
5:A:5:LEU:HD12	5:A:9:ASP:HB3	1.98	0.45
1:B:310:GLY:O	1:B:337:LYS:HE3	2.17	0.45
3:R:216:PHE:O	3:R:220:ARG:HG2	2.16	0.45
5:A:281:VAL:HB	5:A:314:VAL:HG12	1.99	0.45
1:B:33:ILE:HG22	2:G:38:MET:HE1	1.98	0.44
1:B:192:LEU:HD23	1:B:199:PHE:HB3	1.99	0.44
1:B:6:GLN:O	1:B:9:GLN:HG3	2.17	0.44
3:R:138:CYS:SG	3:R:204:MET:HE1	2.57	0.44
1:B:250:CYS:HB2	1:B:264:TYR:HB2	2.00	0.44
3:R:202:PHE:HE2	3:R:266:ILE:HD11	1.83	0.44
3:R:75:LEU:HB2	3:R:80:TYR:CD2	2.53	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:190:LEU:HD12	1:B:201:SER:HB3	2.00	0.44
3:R:59:LEU:O	3:R:63:ILE:HG12	2.18	0.43
1:B:211:TRP:CZ3	1:B:218:CYS:HB2	2.54	0.43
3:R:185:ILE:CD1	4:C:7:ARG:NE	2.79	0.43
1:B:231:ALA:HB1	1:B:276:VAL:HG22	2.00	0.43
1:B:105:TYR:CZ	1:B:109:GLY:HA2	2.54	0.43
6:S:174:LEU:HB3	6:S:192:MET:HB2	2.01	0.42
3:R:296:ILE:O	3:R:299:PRO:HD2	2.18	0.42
1:B:13:GLN:NE2	1:B:14:LEU:HG	2.33	0.42
1:B:187:VAL:HG22	1:B:203:ALA:HB2	2.00	0.42
6:S:151:PRO:HD3	6:S:247:LEU:HD11	2.00	0.42
1:B:283:ARG:HA	1:B:283:ARG:HD2	1.93	0.42
4:C:2:ARG:O	4:C:3:CYS:C	2.62	0.42
3:R:75:LEU:HD12	3:R:75:LEU:O	2.20	0.42
6:S:48:VAL:HG13	6:S:64:VAL:HG21	2.02	0.41
3:R:183:LEU:HA	3:R:186:ILE:HG12	2.01	0.41
3:R:260:PRO:HB2	3:R:287:TYR:CD1	2.56	0.41
1:B:64:GLY:HA2	1:B:105:TYR:CD2	2.56	0.41
1:B:294:CYS:HB3	1:B:308:LEU:HB2	2.02	0.41
3:R:305:ARG:HB3	5:A:372:GLU:OE2	2.21	0.41
6:S:98:ARG:HG2	6:S:110:PHE:HB3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	335/360 (93%)	319 (95%)	16 (5%)	0	100	100
2	G	52/71 (73%)	51 (98%)	1 (2%)	0	100	100
3	R	255/350 (73%)	246 (96%)	9 (4%)	0	100	100
4	C	6/10 (60%)	2 (33%)	4 (67%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
5	A	218/373 (58%)	213 (98%)	5 (2%)	0	100	100
6	S	226/307 (74%)	223 (99%)	3 (1%)	0	100	100
All	All	1092/1471 (74%)	1054 (96%)	38 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	276/299 (92%)	276 (100%)	0	100	100
2	G	43/58 (74%)	43 (100%)	0	100	100
3	R	231/310 (74%)	231 (100%)	0	100	100
4	C	6/6 (100%)	4 (67%)	2 (33%)	0	0
5	A	196/330 (59%)	195 (100%)	1 (0%)	81	92
6	S	192/247 (78%)	192 (100%)	0	100	100
All	All	944/1250 (76%)	941 (100%)	3 (0%)	84	94

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	C	3	CYS
4	C	9	CYS
5	A	246	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	156	GLN
3	R	283	HIS
5	A	193	GLN
6	S	39	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
6	S	82	GLN
6	S	179	GLN
6	S	194	ASN

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains

2 non-standard protein/DNA/RNA residues are modelled in this entry.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.


6 Map visualisation

This section contains visualisations of the EMDB entry EMD-79944. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)


6.2.1 Primary map



X Index: 192




Y Index: 192




Z Index: 192


6.2.2 Raw map



X Index: 192



Y Index: 192




Z Index: 192


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)


6.3.1 Primary map



X Index: 190




Y Index: 211




Z Index: 225


6.3.2 Raw map



X Index: 190



Y Index: 211




Z Index: 225


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) (i)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views (i)

6.5.1 Primary map



X Y Z

The images above show the 3D surface view of the map at the recommended contour level 0.112. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



X Y Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate



The volume at the recommended contour level is 52 nm^3; this corresponds to an approximate mass of 47 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum (i)




*Reported resolution corresponds to spatial frequency of 0.385 Å^{-1}

8 Fourier-Shell correlation

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC



*Reported resolution corresponds to spatial frequency of 0.385 Å^{-1}

8.2 Resolution estimates (i)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.60	-	-
Author-provided FSC curve	2.59	2.89	2.61
Unmasked-calculated*	3.00	3.49	3.04

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.00 differs from the reported value 2.6 by more than 10 %


9 Map-model fit

This section contains information regarding the fit between EMDB map EMD-79944 and PDB model 40BG. Per-residue inclusion information can be found in section 3 on page 7.


9.1 Map-model overlay



X




Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.112 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.


9.2 Q-score mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.112).

9.4 Atom inclusion (i)



At the recommended contour level, 88% of all backbone atoms, 87% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.112) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8740	0.6150
A	0.8580	0.6020
B	0.9170	0.6340
C	0.7500	0.5860
G	0.7050	0.5570
R	0.8620	0.6040
S	0.8970	0.6260

