



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 02:48 PM UTC

PDB ID : 41BI / pdb_000041bi
Title : INTERLEUKIN-1 BETA (IL-1 BETA) (MUTANT WITH CYS 8 REPLACED BY ALA (C8A)
Authors : Veerapandian, B.; Poulos, T.L.; Gilliland, G.L.; Masui, Y.; Hirai, Y.
Deposited on : 1992-02-03
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	153	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1147 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called INTERLEUKIN-1 BETA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	151	Total	C	N	O	S	0	0	1
			1147	743	187	210	7			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	8	ALA	CYS	conflict	UNP P01584


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: INTERLEUKIN-1 BETA

Chain A: 



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, α , β , γ	55.14Å 55.14Å 76.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 2.90	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-2.90)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	RESTRAIN	Depositor
R, R_{free}	0.190 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	1147	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.31	12/1168 (1.0%)	1.61	13/1573 (0.8%)

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	30	HIS	CE1-NE2	7.60	1.40	1.32
1	A	30	HIS	ND1-CE1	6.09	1.38	1.32
1	A	120	TRP	NE1-CE2	-5.35	1.31	1.37
1	A	15	GLN	CD-OE1	5.29	1.33	1.23
1	A	38	GLN	CD-OE1	5.28	1.33	1.23
1	A	89	ASN	CG-OD1	5.22	1.33	1.23
1	A	81	GLN	CD-OE1	5.20	1.33	1.23
1	A	32	GLN	CD-OE1	5.19	1.33	1.23
1	A	31	LEU	N-CA	5.18	1.52	1.46
1	A	141	GLN	CD-OE1	5.14	1.33	1.23
1	A	48	GLN	CD-OE1	5.09	1.33	1.23
1	A	14	GLN	CD-OE1	5.05	1.33	1.23

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	146	PHE	CA-CB-CG	-7.67	106.13	113.80
1	A	142	ASP	CA-CB-CG	-7.21	105.39	112.60
1	A	46	PHE	CA-CB-CG	-7.15	106.65	113.80
1	A	56	ILE	CB-CA-C	6.92	117.15	110.16
1	A	76	ASP	CA-CB-CG	-6.89	105.71	112.60
1	A	58	VAL	CA-CB-CG2	6.15	120.85	110.40
1	A	152	SER	N-CA-CB	5.73	120.25	110.50
1	A	145	ASP	N-CA-C	5.51	118.38	109.40
1	A	51	GLU	N-CA-C	5.50	118.69	109.95
1	A	53	ASN	CA-C-N	5.19	131.38	122.09
1	A	53	ASN	C-N-CA	5.19	131.38	122.09
1	A	3	VAL	CA-CB-CG2	5.10	119.07	110.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	56	ILE	CA-C-O	5.03	122.56	119.94

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1147	0	1120	35	0
All	All	1147	0	1120	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:44:MET:HE3	1:A:148:MET:HE1	1.42	1.00
1:A:71:CYS:HG	1:A:117:PHE:HE2	0.92	0.91
1:A:71:CYS:SG	1:A:72:VAL:N	2.52	0.82
1:A:10:LEU:HD21	1:A:148:MET:HE3	1.62	0.81
1:A:44:MET:CE	1:A:148:MET:HE1	2.11	0.81
1:A:34:GLN:CG	1:A:34:GLN:NE2	2.47	0.77
1:A:72:VAL:HG23	1:A:74:LYS:HG3	1.68	0.76
1:A:71:CYS:SG	1:A:117:PHE:HE2	2.08	0.75
1:A:35:ASP:O	1:A:38:GLN:HG2	1.95	0.67
1:A:10:LEU:CD2	1:A:148:MET:HE3	2.25	0.66
1:A:69:LEU:HD12	1:A:82:LEU:CD2	2.27	0.64
1:A:90:TYR:HD1	1:A:95:MET:HE1	1.64	0.62
1:A:69:LEU:HD12	1:A:82:LEU:HD23	1.83	0.60
1:A:140:GLY:O	1:A:141:GLN:HB2	2.01	0.60
1:A:23:PRO:HB2	1:A:24:TYR:CD2	2.36	0.60
1:A:36:MET:O	1:A:39:GLN:HB3	2.04	0.56
1:A:19:VAL:HA	1:A:40:VAL:HG23	1.90	0.53
1:A:90:TYR:CD1	1:A:95:MET:HE1	2.46	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:26:LEU:HD13	1:A:82:LEU:HD11	1.95	0.49
1:A:69:LEU:HD12	1:A:69:LEU:N	2.27	0.48
1:A:21:SER:HB2	1:A:27:LYS:HG3	1.93	0.48
1:A:69:LEU:N	1:A:69:LEU:CD1	2.76	0.48
1:A:90:TYR:CD1	1:A:95:MET:CE	2.99	0.46
1:A:72:VAL:CG2	1:A:74:LYS:HG3	2.44	0.44
1:A:12:ASP:C	1:A:14:GLN:H	2.25	0.44
1:A:77:LYS:HE2	1:A:77:LYS:HB3	1.64	0.44
1:A:90:TYR:HD1	1:A:95:MET:CE	2.30	0.44
1:A:71:CYS:SG	1:A:72:VAL:CA	3.06	0.43
1:A:69:LEU:HD12	1:A:82:LEU:HD21	2.01	0.42
1:A:141:GLN:HB3	1:A:142:ASP:H	1.49	0.42
1:A:8:ALA:HB3	1:A:148:MET:HE2	2.02	0.41
1:A:71:CYS:SG	1:A:117:PHE:CE2	2.96	0.41
1:A:71:CYS:HA	1:A:79:THR:O	2.20	0.41
1:A:106:ILE:HG22	1:A:106:ILE:O	2.21	0.40
1:A:140:GLY:O	1:A:141:GLN:CB	2.64	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	149/153 (97%)	138 (93%)	9 (6%)	2 (1%)	9	32

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	141	GLN
1	A	140	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	117/139 (84%)	109 (93%)	8 (7%)	14	42

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	GLN
1	A	20	MET
1	A	26	LEU
1	A	29	LEU
1	A	35	ASP
1	A	39	GLN
1	A	71	CYS
1	A	92	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	14	GLN
1	A	102	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.