wwPDB X-ray Structure Validation Summary Report (i)

Nov 12, 2026 – 12:14 PM JST

PDB ID	:	23AO / pdb_000023ao
Title	:	Structure of the Rieske Non-heme Iron Oxygenase VanA with vanillate bound
Authors	:	Cai, C.G.
Deposited on	:	2026-01-30
Resolution	:	2.68 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.2
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.68 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5070 (2.70-2.66)
Clashscore	190562	5409 (2.70-2.66)
Ramachandran outliers	187476	5324 (2.70-2.66)
Sidechain outliers	187428	5324 (2.70-2.66)
RSRZ outliers	180081	5070 (2.70-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	353	93% 7%
1	B	353	2% 94% 6%
1	C	353	% 93% 6%
1	D	353	% 90% 8%
1	E	353	7% 92% 8%
1	F	353	6% 90% 10%

2 Entry composition (i)


There are 6 unique types of molecules in this entry. The entry contains 16974 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Vanillate O-demethylase oxygenase subunit.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	353	Total	C	N	O	S	0	0	0
			2774	1761	485	511	17			
1	B	353	Total	C	N	O	S	0	0	0
			2774	1761	485	511	17			
1	C	349	Total	C	N	O	S	0	0	0
			2747	1746	478	506	17			
1	D	348	Total	C	N	O	S	0	0	0
			2739	1740	477	505	17			
1	E	353	Total	C	N	O	S	0	0	0
			2774	1761	485	511	17			
1	F	353	Total	C	N	O	S	0	0	0
			2774	1761	485	511	17			

- Molecule 2 is FE2/S2 (INORGANIC) CLUSTER (CCD ID: FES) (formula: Fe_2S_2) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Fe S 4 2 2	0	0
2	B	1	Total Fe S 4 2 2	0	0
2	C	1	Total Fe S 4 2 2	0	0
2	D	1	Total Fe S 4 2 2	0	0
2	E	1	Total Fe S 4 2 2	0	0
2	F	1	Total Fe S 4 2 2	0	0

- Molecule 3 is 4-HYDROXY-3-METHOXYBENZOATE (CCD ID: VNL) (formula: $C_8H_7O_4$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 12 8 4	0	0
3	B	1	Total C O 12 8 4	0	0
3	C	1	Total C O 12 8 4	0	0
3	D	1	Total C O 12 8 4	0	0
3	E	1	Total C O 12 8 4	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	F	1	Total C O 12 8 4	0	0

- Molecule 4 is FE (III) ION (CCD ID: FE) (formula: Fe).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Fe 1 1	0	0
4	B	1	Total Fe 1 1	0	0
4	C	1	Total Fe 1 1	0	0
4	D	1	Total Fe 1 1	0	0
4	E	1	Total Fe 1 1	0	0
4	F	1	Total Fe 1 1	0	0

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	D	1	Total C O 4 2 2	0	0

- Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	52	Total O 52 52	0	0
6	B	72	Total O 72 72	0	0
6	C	64	Total O 64 64	0	0
6	D	49	Total O 49 49	0	0
6	E	31	Total O 31 31	0	0
6	F	10	Total O 10 10	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Vanillate O-demethylase oxygenase subunit

Chain A:




• Molecule 1: Vanillate O-demethylase oxygenase subunit

Chain B:




• Molecule 1: Vanillate O-demethylase oxygenase subunit

Chain C:




• Molecule 1: Vanillate O-demethylase oxygenase subunit


Chain D:




• Molecule 1: Vanillate O-demethylase oxygenase subunit


Chain E:





• Molecule 1: Vanillate O-demethylase oxygenase subunit





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	97.72Å 103.79Å 130.31Å 90.00° 100.11° 90.00°	Depositor
Resolution (Å)	49.51 – 2.68 49.51 – 2.68	Depositor EDS
% Data completeness (in resolution range)	99.6 (49.51-2.68) 99.6 (49.51-2.68)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.95 (at 2.69Å)	Xtriage
Refinement program	PHENIX 1.15.1_3469	Depositor
R, R_{free}	0.193 , 0.234 0.193 , 0.233	Depositor DCC
R_{free} test set	3590 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å^2)	43.1	Xtriage
Anisotropy	0.232	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.33 , 47.6	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	16974	wwPDB-VP
Average B, all atoms (Å^2)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.56% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FE, VNL, FES, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| > 5$	RMSZ	$\#\|Z\| > 5$
1	A	0.09	0/2847	0.27	0/3869
1	B	0.09	0/2847	0.27	0/3869
1	C	0.09	0/2820	0.27	0/3833
1	D	0.09	0/2812	0.27	0/3822
1	E	0.09	0/2847	0.26	0/3869
1	F	0.09	0/2847	0.27	0/3869
All	All	0.09	0/17020	0.27	0/23131

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2774	0	2685	15	0
1	B	2774	0	2685	12	0
1	C	2747	0	2657	10	0
1	D	2739	0	2647	17	0
1	E	2774	0	2685	16	0
1	F	2774	0	2685	16	0
2	A	4	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	4	0	0	0	0
2	C	4	0	0	1	0
2	D	4	0	0	0	0
2	E	4	0	0	0	0
2	F	4	0	0	0	0
3	A	12	0	6	0	0
3	B	12	0	6	0	0
3	C	12	0	6	0	0
3	D	12	0	6	0	0
3	E	12	0	6	0	0
3	F	12	0	6	1	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
4	E	1	0	0	0	0
4	F	1	0	0	0	0
5	B	8	0	12	0	0
5	D	4	0	6	0	0
6	A	52	0	0	2	0
6	B	72	0	0	0	0
6	C	64	0	0	0	0
6	D	49	0	0	1	0
6	E	31	0	0	0	0
6	F	10	0	0	0	0
All	All	16974	0	16098	82	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 82 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:86:ARG:HH21	1:A:88:GLN:HG2	1.54	0.73
1:B:274:ARG:NH1	1:B:277:ARG:O	2.24	0.70
1:D:274:ARG:NH2	1:D:284:THR:OG1	2.29	0.66
1:F:165:ILE:HG23	1:F:198:PHE:HB3	1.80	0.63
1:E:245:PRO:HD2	1:E:248:TYR:HD2	1.65	0.62

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	351/353 (99%)	341 (97%)	9 (3%)	1 (0%)	36	57
1	B	351/353 (99%)	342 (97%)	8 (2%)	1 (0%)	36	57
1	C	347/353 (98%)	333 (96%)	14 (4%)	0	100	100
1	D	346/353 (98%)	336 (97%)	9 (3%)	1 (0%)	36	57
1	E	351/353 (99%)	343 (98%)	8 (2%)	0	100	100
1	F	351/353 (99%)	337 (96%)	13 (4%)	1 (0%)	36	57
All	All	2097/2118 (99%)	2032 (97%)	61 (3%)	4 (0%)	43	65

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	126	ALA
1	A	126	ALA
1	B	126	ALA
1	F	49	HIS

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	284/284 (100%)	282 (99%)	2 (1%)	76	89
1	B	284/284 (100%)	282 (99%)	2 (1%)	76	89
1	C	282/284 (99%)	277 (98%)	5 (2%)	51	76

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	D	281/284 (99%)	277 (99%)	4 (1%)	59 80
1	E	284/284 (100%)	279 (98%)	5 (2%)	51 76
1	F	284/284 (100%)	273 (96%)	11 (4%)	28 54
All	All	1699/1704 (100%)	1670 (98%)	29 (2%)	53 77

5 of 29 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	127	ASP
1	F	218	GLN
1	E	243	ASP
1	F	175	VAL
1	E	239	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	27	ASN
1	E	69	HIS
1	F	328	GLN
1	C	5	ASN
1	B	279	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 21 ligands modelled in this entry, 6 are monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	EDO	B	405	-	3,3,3	0.45	0	2,2,2	0.36	0
5	EDO	D	401	-	3,3,3	0.46	0	2,2,2	0.33	0
2	FES	B	402	1	0,4,4	-	-	-		
3	VNL	D	403	-	12,12,12	1.45	1 (8%)	16,16,16	1.33	1 (6%)
3	VNL	F	402	-	12,12,12	1.49	1 (8%)	16,16,16	1.41	1 (6%)
2	FES	F	401	1	0,4,4	-	-	-		
3	VNL	A	402	-	12,12,12	1.49	1 (8%)	16,16,16	1.40	1 (6%)
2	FES	D	402	1	0,4,4	-	-	-		
5	EDO	B	401	-	3,3,3	0.46	0	2,2,2	0.32	0
2	FES	E	401	1	0,4,4	-	-	-		
3	VNL	E	402	-	12,12,12	1.48	1 (8%)	16,16,16	1.39	1 (6%)
3	VNL	C	402	-	12,12,12	1.46	1 (8%)	16,16,16	1.34	2 (12%)
2	FES	A	401	1	0,4,4	-	-	-		
2	FES	C	401	1	0,4,4	-	-	-		
3	VNL	B	403	-	12,12,12	1.45	1 (8%)	16,16,16	1.32	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	D	401	-	-	0/1/1/1	-
3	VNL	D	403	-	-	0/6/6/6	0/1/1/1
3	VNL	F	402	-	-	4/6/6/6	0/1/1/1
2	FES	B	402	1	-	-	0/1/1/1
2	FES	F	401	1	-	-	0/1/1/1
3	VNL	A	402	-	-	0/6/6/6	0/1/1/1
2	FES	D	402	1	-	-	0/1/1/1
5	EDO	B	401	-	-	0/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FES	C	401	1	-	-	0/1/1/1
2	FES	A	401	1	-	-	0/1/1/1
2	FES	E	401	1	-	-	0/1/1/1
3	VNL	C	402	-	-	0/6/6/6	0/1/1/1
5	EDO	B	405	-	-	0/1/1/1	-
3	VNL	E	402	-	-	0/6/6/6	0/1/1/1
3	VNL	B	403	-	-	0/6/6/6	0/1/1/1

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	402	VNL	CZ-CM2	4.74	1.48	1.40
3	F	402	VNL	CZ-CM2	4.74	1.48	1.40
3	E	402	VNL	CZ-CM2	4.70	1.48	1.40
3	C	402	VNL	CZ-CM2	4.63	1.48	1.40
3	D	403	VNL	CZ-CM2	4.59	1.48	1.40

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	F	402	VNL	OM-CM2-CZ	4.05	120.44	114.57
3	E	402	VNL	OM-CM2-CZ	3.93	120.27	114.57
3	A	402	VNL	OM-CM2-CZ	3.92	120.24	114.57
3	D	403	VNL	OM-CM2-CZ	3.68	119.90	114.57
3	B	403	VNL	OM-CM2-CZ	3.65	119.85	114.57

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	F	402	VNL	CO1-C1-CC-O1
3	F	402	VNL	CO1-C1-CC-O2
3	F	402	VNL	CO2-C1-CC-O2
3	F	402	VNL	CO2-C1-CC-O1

There are no ring outliers.

3 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	402	VNL	1	0
2	A	401	FES	1	0

Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	401	FES	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand VNL D 403




Bond lengths



Bond angles




Torsions




Rings


Ligand VNL F 402




Bond lengths



Bond angles




Torsions




Rings


Ligand FES F 401




Bond lengths



Bond angles




Torsions




Rings


Ligand VNL A 402




Bond lengths



Bond angles




Torsions




Rings


Ligand FES D 402




Bond lengths



Bond angles




Torsions




Rings


Ligand FES E 401




Bond lengths



Bond angles




Torsions




Rings


Ligand VNL E 402




Bond lengths



Bond angles




Torsions




Rings


Ligand VNL C 402




Bond lengths



Bond angles




Torsions




Rings


Ligand FES A 401




Bond lengths



Bond angles




Torsions




Rings


Ligand FES C 401




Bond lengths



Bond angles




Torsions




Rings


Ligand VNL B 403




Bond lengths



Bond angles



Torsions



Rings

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	353/353 (100%)	-0.22	1 (0%)	90	89	24, 42, 74, 100	0
1	B	353/353 (100%)	-0.28	6 (1%)	69	66	22, 37, 73, 154	0
1	C	349/353 (98%)	-0.25	4 (1%)	78	76	24, 41, 72, 148	0
1	D	348/353 (98%)	0.02	5 (1%)	73	71	25, 52, 96, 148	0
1	E	353/353 (100%)	0.35	24 (6%)	23	20	29, 61, 112, 152	0
1	F	353/353 (100%)	0.69	20 (5%)	29	25	49, 81, 128, 219	0
All	All	2109/2118 (99%)	0.05	60 (2%)	55	51	22, 51, 108, 219	0

The worst 5 of 60 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	348	LEU	3.9
1	E	246	ALA	3.8
1	E	238	GLY	3.8
1	E	248	TYR	3.7
1	B	347	ALA	3.7

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no oligosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	VNL	F	402	12/12	0.81	0.24	67,85,93,94	0
5	EDO	B	405	4/4	0.85	0.16	44,46,50,50	0
5	EDO	B	401	4/4	0.87	0.16	60,62,67,67	0
5	EDO	D	401	4/4	0.91	0.15	35,39,45,48	0
3	VNL	E	402	12/12	0.92	0.10	47,49,51,51	0
3	VNL	C	402	12/12	0.94	0.11	38,42,46,48	0
3	VNL	A	402	12/12	0.95	0.07	22,33,40,40	0
2	FES	F	401	4/4	0.95	0.08	51,54,58,59	0
4	FE	F	403	1/1	0.95	0.05	60,60,60,60	0
3	VNL	D	403	12/12	0.96	0.09	20,27,31,31	0
4	FE	B	404	1/1	0.96	0.16	51,51,51,51	0
3	VNL	B	403	12/12	0.96	0.09	29,32,34,37	0
2	FES	D	402	4/4	0.97	0.05	58,61,61,63	0
4	FE	C	403	1/1	0.98	0.06	36,36,36,36	0
4	FE	E	403	1/1	0.98	0.04	37,37,37,37	0
2	FES	B	402	4/4	0.98	0.04	32,33,35,36	0
2	FES	C	401	4/4	0.98	0.04	33,37,37,37	0
2	FES	A	401	4/4	0.98	0.05	32,32,35,39	0
2	FES	E	401	4/4	0.98	0.04	36,37,38,45	0
4	FE	D	404	1/1	0.99	0.04	33,33,33,33	0
4	FE	A	403	1/1	0.99	0.04	32,32,32,32	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around VNL F 402:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)









Electron density around VNL E 402:


$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around VNL C 402:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)










Electron density around VNL A 402:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)









Electron density around FES F 401:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around VNL D 403:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around VNL B 403:


$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around FES D 402:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)










Electron density around FES B 402:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)










Electron density around FES C 401:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)










Electron density around FES A 401:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)








Electron density around FES E 401:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers

There are no such residues in this entry.