



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 02:22 PM UTC

PDB ID : 244D / pdb_0000244d
Title : THE HIGH-RESOLUTION CRYSTAL STRUCTURE OF A PARALLEL-STRANDED GUANINE TETRAPLEX
Authors : Laughlan, G.; Murchie, A.I.H.; Norman, D.G.; Moore, M.H.; Moody, P.C.E.; Lilley, D.M.J.; Luisi, B.
Deposited on : 1995-10-19
Resolution : 1.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.20 Å.









Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1265 (1.20-1.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	6	 83% 17%
1	B	6	 67% 33%
1	C	6	 83% 17%
1	D	6	 83% 17%
1	E	6	 83% 17%
1	F	6	 100%
1	G	6	 67% 33%
1	H	6	 50% 50%
1	I	6	 83% 17%
1	J	6	 50% 33% 17%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	K	6	 50% 50%
1	L	6	 67% 33%
1	M	6	 83% 17%
1	N	6	 67% 33%
1	O	6	 67% 33%
1	P	6	 33% 50% 17%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2453 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(*TP*GP*GP*GP*GP*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	A	6	125	60	24	36	5	0	0	0
1	B	6	108	50	22	31	5	0	0	0
1	C	6	109	50	22	32	5	0	0	1
1	D	6	125	60	24	36	5	0	0	0
1	E	6	108	50	22	31	5	0	0	0
1	F	6	125	60	24	36	5	0	0	0
1	G	6	125	60	24	36	5	0	0	0
1	H	6	125	60	24	36	5	0	0	0
1	I	6	125	60	24	36	5	0	0	0
1	J	6	125	60	24	36	5	0	0	0
1	K	6	108	50	22	31	5	0	0	0
1	L	6	125	60	24	36	5	0	0	0
1	M	6	125	60	24	36	5	0	0	0
1	N	6	125	60	24	36	5	0	0	0
1	O	6	108	50	22	31	5	0	0	0
1	P	6	125	60	24	36	5	0	0	0

- Molecule 2 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total 3	Na 3	0	0
2	B	1	Total 1	Na 1	0	0
2	E	3	Total 3	Na 3	0	0
2	I	3	Total 3	Na 3	0	0
2	L	1	Total 1	Na 1	0	0
2	M	3	Total 3	Na 3	0	0

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total 2	Ca 2	0	0
3	B	1	Total 1	Ca 1	0	0
3	C	1	Total 1	Ca 1	0	0
3	D	1	Total 1	Ca 1	0	0
3	L	1	Total 1	Ca 1	0	0
3	M	1	Total 1	Ca 1	0	0
3	N	1	Total 1	Ca 1	0	0
3	P	1	Total 1	Ca 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	37	Total 37	O 37	0	0
4	B	32	Total 32	O 32	0	0
4	C	26	Total 26	O 26	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	D	49	Total O 49 49	0	0
4	E	28	Total O 28 28	0	0
4	F	28	Total O 28 28	0	0
4	G	31	Total O 31 31	0	0
4	H	28	Total O 28 28	0	0
4	I	36	Total O 36 36	0	0
4	J	24	Total O 24 24	0	0
4	K	19	Total O 19 19	0	0
4	L	37	Total O 37 37	0	0
4	M	41	Total O 41 41	0	0
4	N	35	Total O 35 35	0	0
4	O	28	Total O 28 28	0	0
4	P	35	Total O 35 35	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*GP*T)-3')

Chain A:  83% 17%




- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*GP*T)-3')

Chain B:  67% 33%




- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*GP*T)-3')

Chain C:  83% 17%




- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*GP*T)-3')

Chain D:  83% 17%



- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*GP*T)-3')

Chain E:  83% 17%



- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*GP*T)-3')

Chain F:  100%

There are no outlier residues recorded for this chain.

- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*GP*T)-3')

Chain G:  67% 33%




- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*GP*T)-3')

Chain H:  50% 50%



- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*GP*T)-3')

Chain I:  83% 17%



- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*GP*T)-3')

Chain J:  50% 33% 17%



- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*GP*T)-3')

Chain K:  50% 50%




- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*GP*T)-3')

Chain L:  67% 33%



- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*GP*T)-3')

Chain M:  83% 17%




- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*GP*T)-3')



- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*GP*T)-3')



- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*GP*T)-3')



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	28.76Å 35.47Å 56.77Å 74.39° 77.64° 89.73°	Depositor
Resolution (Å)	8.00 – 1.20	Depositor
% Data completeness (in resolution range)	97.7 (8.00-1.20)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	SHELX-93	Depositor
R, R_{free}	0.124 , 0.176	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	2453	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.04	0/140	1.48	1/216 (0.5%)
1	B	0.89	0/121	1.73	2/186 (1.1%)
1	C	1.01	0/122	1.65	1/189 (0.5%)
1	D	0.99	0/140	1.46	1/216 (0.5%)
1	E	0.87	0/121	1.49	1/186 (0.5%)
1	F	1.02	0/140	1.57	0/216
1	G	1.05	0/140	1.77	3/216 (1.4%)
1	H	0.91	0/140	1.92	5/216 (2.3%)
1	I	0.95	0/140	1.50	1/216 (0.5%)
1	J	0.92	0/140	1.79	4/216 (1.9%)
1	K	0.94	0/121	1.99	5/186 (2.7%)
1	L	1.12	0/140	1.73	3/216 (1.4%)
1	M	1.00	0/140	1.46	1/216 (0.5%)
1	N	1.10	0/140	2.07	4/216 (1.9%)
1	O	0.98	0/121	1.67	3/186 (1.6%)
1	P	0.94	0/140	1.80	4/216 (1.9%)
All	All	0.99	0/2146	1.70	39/3309 (1.2%)

There are no bond length outliers.

The worst 5 of 39 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	N	131	DT	P-O3'-C3'	19.11	148.87	120.20
1	B	16	DT	OP1-P-OP2	10.68	152.05	120.00
1	K	106	DT	OP1-P-OP2	-9.66	91.01	120.00
1	K	101	DT	N1-C1'-C2'	9.59	127.89	113.50
1	H	76	DT	P-O5'-C5'	8.74	133.12	120.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	125	0	70	0	0
1	B	108	0	57	0	0
1	C	109	0	57	0	0
1	D	125	0	70	0	0
1	E	108	0	57	0	0
1	F	125	0	70	0	0
1	G	125	0	69	0	0
1	H	125	0	70	0	0
1	I	125	0	70	0	0
1	J	125	0	70	1	0
1	K	108	0	57	0	0
1	L	125	0	70	0	0
1	M	125	0	70	0	0
1	N	125	0	70	0	0
1	O	108	0	57	0	0
1	P	125	0	70	1	0
2	A	3	0	0	0	0
2	B	1	0	0	0	0
2	E	3	0	0	0	0
2	I	3	0	0	0	0
2	L	1	0	0	0	0
2	M	3	0	0	0	0
3	A	2	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	L	1	0	0	0	0
3	M	1	0	0	0	0
3	N	1	0	0	0	0
3	P	1	0	0	0	0
4	A	37	0	0	0	0
4	B	32	0	0	0	0
4	C	26	0	0	0	0
4	D	49	0	0	0	0
4	E	28	0	0	0	0
4	F	28	0	0	0	0
4	G	31	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	H	28	0	0	0	0
4	I	36	0	0	0	0
4	J	24	0	0	0	0
4	K	19	0	0	0	0
4	L	37	0	0	0	0
4	M	41	0	0	0	0
4	N	35	0	0	0	0
4	O	28	0	0	0	0
4	P	35	0	0	0	0
All	All	2453	0	1054	1	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All (1) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:J:91:DT:O4'	1:P:151:DT:H2'	2.19	0.42

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 23 ligands modelled in this entry, 23 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.