



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:46 PM UTC

PDB ID : 15C8 / pdb_000015c8
Title : CATALYTIC ANTIBODY 5C8, FREE FAB
Authors : Gruber, K.; Wilson, I.A.
Deposited on : 1998-03-18
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	213	2% 59% 32% 7%
2	H	217	7% 64% 30% 6%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3350 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IGG 5C8 FAB (LIGHT CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	213	Total	C	N	O	S	0	0	0
			1633	1012	276	337	8			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
L	32	ASN	TYR	conflict	UNP P01837
L	94	SER	PHE	conflict	UNP P01837
L	96	TYR	HIS	conflict	UNP P01837

- Molecule 2 is a protein called IGG 5C8 FAB (HEAVY CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	217	Total	C	N	O	S	46	0	0
			1626	1028	262	330	6			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	3	GLN	LYS	conflict	UNP P01869
H	5	GLN	LEU	conflict	UNP P01869
H	6	GLN	GLU	conflict	UNP P01869
H	14	PRO	SER	conflict	UNP P01869
H	40	LYS	ARG	conflict	UNP P01869
H	49	ALA	GLY	conflict	UNP P01869
H	50	GLN	ARG	conflict	UNP P01869
H	56	ASN	GLU	conflict	UNP P01869
H	57	THR	ILE	conflict	UNP P01869
H	66	LYS	THR	conflict	UNP P01869
H	75	SER	THR	conflict	UNP P01869
H	81	HIS	GLN	conflict	UNP P01869
H	87	SER	THR	conflict	UNP P01869

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	93	ALA	VAL	conflict	UNP P01869
H	94	ALA	ARG	conflict	UNP P01869
H	95	ASP	ARG	conflict	UNP P01869
H	96	PRO	GLY	conflict	UNP P01869
H	97	PRO	TYR	conflict	UNP P01869
H	98	TYR	GLY	conflict	UNP P01869
H	99	TYR	SER	conflict	UNP P01869
H	100	GLY	SER	conflict	UNP P01869
H	100A	HIS	GLN	conflict	UNP P01869
H	100B	GLY	GLU	conflict	UNP P01869
H	101	ASP	PRO	conflict	UNP P01869


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	L	54	Total O 54 54	0	0
3	H	37	Total O 37 37	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: IGG 5C8 FAB (LIGHT CHAIN)



• Molecule 2: IGG 5C8 FAB (HEAVY CHAIN)



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	201.00Å 80.50Å 39.30Å 90.00° 98.70° 90.00°	Depositor
Resolution (Å)	30.00 – 2.50 30.00 – 2.50	Depositor EDS
% Data completeness (in resolution range)	90.0 (30.00-2.50) 89.7 (30.00-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.02 (at 2.45Å)	Xtriage
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.190 , 0.251 0.185 , 0.245	Depositor DCC
R_{free} test set	1448 reflections (7.35%)	wwPDB-VP
Wilson B-factor (Å ²)	27.4	Xtriage
Anisotropy	0.377	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 53.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.043 for -h-2*k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3350	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	L	0.41	0/1672	0.88	4/2271 (0.2%)
2	H	0.44	0/1671	0.91	7/2291 (0.3%)
All	All	0.42	0/3343	0.90	11/4562 (0.2%)

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	27(A)	SER	N-CA-C	6.57	119.29	110.35
2	H	60	ASP	N-CA-C	-6.48	100.23	109.24
2	H	97	PRO	N-CA-C	-6.13	106.00	114.27
2	H	96	PRO	N-CA-C	5.84	116.83	110.58
1	L	66	GLY	N-CA-C	5.57	118.03	111.63
2	H	100(B)	GLY	N-CA-C	5.49	119.54	112.18
2	H	82(C)	LEU	N-CA-C	5.41	117.27	110.24
1	L	122	SER	N-CA-C	-5.35	104.49	111.02
2	H	99	TYR	N-CA-C	5.25	117.40	111.11
2	H	145	LYS	N-CA-C	5.20	118.19	110.14
1	L	91	TYR	N-CA-C	-5.11	103.95	111.81

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1633	0	1554	59	0
2	H	1626	0	1571	40	0
3	H	37	0	0	0	0
3	L	54	0	0	2	0
All	All	3350	0	3125	97	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (97) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:198:HIS:CD2	1:L:200:THR:HB	2.22	0.75
2:H:50:GLN:OE1	2:H:58:LYS:HD3	1.89	0.73
1:L:199:LYS:HE2	1:L:199:LYS:H	1.54	0.71
1:L:161:ASN:HB3	1:L:175:MET:HE3	1.72	0.71
1:L:198:HIS:HD2	1:L:200:THR:HB	1.56	0.71
1:L:14:SER:O	1:L:17:GLU:HB2	1.97	0.64
1:L:199:LYS:HE2	1:L:199:LYS:N	2.13	0.63
1:L:137:ASN:HB3	1:L:138:ASN:HD22	1.63	0.63
2:H:51:ILE:HG13	2:H:57:THR:HG22	1.81	0.63
1:L:112:ALA:CB	1:L:200:THR:HG21	2.30	0.61
2:H:138:VAL:HG22	2:H:139:THR:N	2.17	0.60
2:H:35:HIS:CE1	2:H:50:GLN:HE21	2.21	0.59
1:L:137:ASN:HB3	1:L:138:ASN:ND2	2.18	0.58
2:H:149:PRO:HD2	2:H:214:ALA:CB	2.32	0.58
2:H:180:SER:O	2:H:183:ASP:HB2	2.03	0.58
1:L:189:HIS:HB2	1:L:192:TYR:OH	2.03	0.57
2:H:36:TRP:HD1	2:H:69:ILE:HD12	1.69	0.57
2:H:95:ASP:OD1	2:H:95:ASP:C	2.48	0.57
2:H:138:VAL:HG22	2:H:139:THR:H	1.70	0.57
1:L:78:MET:HE2	1:L:106:ILE:HD13	1.87	0.56
2:H:29:ILE:HG12	2:H:76:ASN:OD1	2.05	0.56
2:H:43:GLN:HE21	2:H:43:GLN:HA	1.68	0.56
2:H:38:LYS:HB2	2:H:48:ILE:HD11	1.88	0.56
1:L:211:ARG:HH11	1:L:211:ARG:HB2	1.71	0.56
2:H:199:TRP:CG	2:H:200:PRO:HA	2.40	0.56
2:H:35:HIS:CE1	2:H:50:GLN:HB2	2.41	0.56
1:L:45:LYS:HE3	1:L:45:LYS:HA	1.88	0.55
2:H:140:LEU:HD23	2:H:223:ILE:HG21	1.89	0.54
1:L:42:SER:HB3	3:L:220:HOH:O	2.07	0.54
1:L:94:SER:HA	1:L:95:PRO:C	2.32	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:112:ALA:HA	1:L:200:THR:HG21	1.90	0.53
1:L:155:ARG:HE	1:L:157:ASN:HB3	1.72	0.53
1:L:195:GLU:HG3	1:L:206:VAL:HG22	1.89	0.53
1:L:199:LYS:H	1:L:199:LYS:CE	2.21	0.53
1:L:1:ASP:OD1	1:L:1:ASP:N	2.42	0.53
1:L:150:ILE:HD11	1:L:179:LEU:HD21	1.91	0.53
2:H:60:ASP:OD1	2:H:62:LYS:HG2	2.10	0.52
2:H:38:LYS:CB	2:H:48:ILE:HD11	2.41	0.51
2:H:123:PRO:HD3	2:H:221:LYS:HG2	1.92	0.51
2:H:51:ILE:HG12	2:H:52:ASP:N	2.25	0.50
2:H:137:MET:HA	2:H:195:SER:H	1.76	0.50
1:L:78:MET:HE2	1:L:106:ILE:CD1	2.41	0.49
2:H:125:ALA:HB2	2:H:223:ILE:HG22	1.93	0.49
1:L:141:PRO:O	1:L:198:HIS:HE1	1.95	0.49
1:L:193:THR:HG23	1:L:208:SER:OG	2.12	0.49
1:L:197:THR:HG23	1:L:204:PRO:HB3	1.94	0.49
2:H:32:THR:OG1	2:H:95:ASP:O	2.29	0.49
2:H:124:LEU:HB2	2:H:141:GLY:C	2.39	0.48
1:L:187:GLU:O	1:L:211:ARG:NH2	2.46	0.48
1:L:150:ILE:O	1:L:151:ASP:HB2	2.13	0.48
2:H:119:PRO:HB3	2:H:147:TYR:HB3	1.96	0.48
2:H:206:THR:HG23	2:H:221:LYS:C	2.39	0.47
1:L:112:ALA:HB2	1:L:200:THR:HG21	1.96	0.47
1:L:159:VAL:HG22	1:L:179:LEU:HD13	1.95	0.47
1:L:37:GLN:HG3	1:L:86:TYR:CE2	2.50	0.47
1:L:150:ILE:HG23	1:L:192:TYR:CE1	2.51	0.46
2:H:153:THR:O	2:H:210:VAL:HA	2.16	0.46
2:H:54:ASN:OD1	2:H:56:ASN:HB2	2.16	0.46
1:L:112:ALA:CA	1:L:200:THR:HG21	2.46	0.46
1:L:61:ARG:HD2	1:L:77:SER:O	2.16	0.45
1:L:43:SER:HB2	2:H:91:TYR:CE1	2.51	0.45
1:L:159:VAL:O	1:L:160:LEU:HD23	2.16	0.45
1:L:187:GLU:HA	1:L:211:ARG:HE	1.80	0.45
1:L:33:LEU:HD22	1:L:71:TYR:CG	2.52	0.45
1:L:115:VAL:HG12	1:L:116:SER:N	2.32	0.45
1:L:170:ASP:OD2	1:L:172:THR:OG1	2.32	0.45
2:H:51:ILE:O	2:H:52(A):PRO:HD3	2.17	0.45
1:L:62:PHE:CE1	1:L:75:ILE:HG12	2.52	0.45
1:L:73:LEU:C	1:L:73:LEU:HD23	2.41	0.44
1:L:198:HIS:O	1:L:199:LYS:C	2.60	0.44
2:H:96:PRO:HA	2:H:97:PRO:HD3	1.71	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:124:GLN:HG2	1:L:129:GLY:O	2.18	0.44
1:L:149:LYS:HA	1:L:153:SER:O	2.18	0.44
2:H:32:THR:OG1	2:H:33:TYR:N	2.50	0.44
1:L:33:LEU:HD22	1:L:71:TYR:CD2	2.52	0.43
2:H:124:LEU:HB2	2:H:141:GLY:O	2.18	0.43
2:H:199:TRP:CD1	2:H:200:PRO:HA	2.53	0.43
2:H:51:ILE:HG13	2:H:57:THR:CG2	2.47	0.43
1:L:1:ASP:HA	3:L:215:HOH:O	2.18	0.43
1:L:45:LYS:HE3	1:L:45:LYS:CA	2.49	0.42
2:H:166:LEU:HD23	2:H:166:LEU:HA	1.91	0.42
1:L:138:ASN:ND2	1:L:138:ASN:N	2.67	0.42
2:H:87:SER:O	2:H:88:ALA:HB2	2.19	0.42
2:H:56:ASN:HD22	2:H:56:ASN:HA	1.61	0.42
2:H:137:MET:HA	2:H:195:SER:N	2.34	0.42
1:L:27(A):SER:HA	1:L:69:THR:HG22	2.01	0.41
1:L:91:TYR:CE2	2:H:95:ASP:OD2	2.73	0.41
1:L:170:ASP:CG	1:L:172:THR:HG1	2.28	0.41
1:L:211:ARG:O	1:L:212:ASN:CB	2.68	0.41
2:H:35:HIS:HE1	2:H:50:GLN:HE21	1.66	0.41
1:L:161:ASN:HB3	1:L:175:MET:CE	2.47	0.41
1:L:192:TYR:HB2	1:L:209:PHE:CE1	2.55	0.41
1:L:151:ASP:C	1:L:153:SER:H	2.29	0.41
1:L:199:LYS:C	1:L:201:SER:H	2.29	0.41
1:L:16:GLY:HA2	1:L:77:SER:OG	2.22	0.40
1:L:58:VAL:HA	1:L:59:PRO:HD3	1.96	0.40
1:L:113:PRO:HB3	1:L:139:PHE:CD2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	211/213 (99%)	200 (95%)	8 (4%)	3 (1%)	9	17
2	H	215/217 (99%)	194 (90%)	18 (8%)	3 (1%)	9	17
All	All	426/430 (99%)	394 (92%)	26 (6%)	6 (1%)	9	17

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	202	SER
2	H	135	ASN
2	H	137	MET
1	L	138	ASN
1	L	184	ASP
1	L	200	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	187/187 (100%)	150 (80%)	37 (20%)	1	2
2	H	184/184 (100%)	150 (82%)	34 (18%)	1	3
All	All	371/371 (100%)	300 (81%)	71 (19%)	1	3

All (71) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	1	ASP
1	L	2	ILE
1	L	10	ILE
1	L	12	SER
1	L	14	SER
1	L	27	SER
1	L	27(A)	SER
1	L	33	LEU
1	L	45	LYS
1	L	65	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	L	78	MET
1	L	94	SER
1	L	105	GLU
1	L	121	SER
1	L	122	SER
1	L	132	VAL
1	L	138	ASN
1	L	142	LYS
1	L	145	ASN
1	L	147	LYS
1	L	149	LYS
1	L	153	SER
1	L	155	ARG
1	L	156	GLN
1	L	168	SER
1	L	169	LYS
1	L	170	ASP
1	L	172	THR
1	L	176	SER
1	L	181	LEU
1	L	183	LYS
1	L	191	SER
1	L	199	LYS
1	L	200	THR
1	L	203	SER
1	L	207	LYS
1	L	211	ARG
2	H	3	GLN
2	H	11	LEU
2	H	25	SER
2	H	40	LYS
2	H	42	GLU
2	H	43	GLN
2	H	50	GLN
2	H	51	ILE
2	H	58	LYS
2	H	62	LYS
2	H	64	GLN
2	H	66	LYS
2	H	74	SER
2	H	85	GLU
2	H	95	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	H	105	GLN
2	H	113	SER
2	H	133	GLN
2	H	142	CYS
2	H	150	GLU
2	H	156	THR
2	H	178	LEU
2	H	179	GLN
2	H	180	SER
2	H	187	LEU
2	H	202	SER
2	H	208	CYS
2	H	209	ASN
2	H	216	SER
2	H	218	LYS
2	H	221	LYS
2	H	222	LYS
2	H	223	ILE
2	H	226	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	38	GLN
1	L	137	ASN
1	L	138	ASN
1	L	157	ASN
1	L	190	ASN
2	H	39	GLN
2	H	43	GLN
2	H	50	GLN
2	H	56	ASN
2	H	64	GLN
2	H	105	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	L	213/213 (100%)	0.03	5 (2%) 61 57	6, 33, 77, 96	0
2	H	210/217 (96%)	0.20	15 (7%) 22 19	8, 33, 81, 95	0
All	All	423/430 (98%)	0.11	20 (4%) 36 32	6, 33, 77, 96	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	226	VAL	4.5
2	H	127	GLY	3.8
2	H	137	MET	3.6
2	H	99	TYR	3.2
2	H	97	PRO	3.0
2	H	98	TYR	2.7
2	H	204	THR	2.7
1	L	199	LYS	2.6
2	H	194	PRO	2.6
2	H	96	PRO	2.5
2	H	163	SER	2.4
2	H	167	SER	2.4
2	H	138	VAL	2.1
2	H	100	GLY	2.1
1	L	160	LEU	2.1
1	L	211	ARG	2.1
1	L	200	THR	2.1
2	H	223	ILE	2.1
1	L	212	ASN	2.0
2	H	205	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.