



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 29, 2026 – 10:28 am BST

PDB ID : 29WJ / pdb_000029wj
Title : Structure of monomerized native leukocyte myeloperoxidase in complex with the Staphylococcal Peroxidase Inhibitor SPIN
Authors : Leitgeb, U.; Pfanzagl, V.
Deposited on : 2026-04-10
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.2
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 2.2
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.015 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.51


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1091 (2.54-2.54)
Clashscore	190562	1120 (2.54-2.54)
Ramachandran outliers	187476	1106 (2.54-2.54)
Sidechain outliers	187428	1106 (2.54-2.54)
RSRZ outliers	180081	1091 (2.54-2.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	114	80% 8% 11%
1	C	114	79% 7% 14%
2	B	467	87% 13%
3	E	73	7% 85% 10% 5%
3	F	73	38% 81% 15%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	D	467	3% 89% 9%
5	G	6	50% 50%
6	H	2	50% 50%
6	I	2	50% 50%
6	L	2	50% 50%
7	J	7	14% 43% 43%
8	K	3	100%

2 Entry composition [i](#)

There are 12 unique types of molecules in this entry. The entry contains 10519 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Myeloperoxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	101	Total	C	N	O	S	0	0	0
			815	516	145	150	4			
1	C	98	Total	C	N	O	S	0	0	0
			797	506	142	145	4			

- Molecule 2 is a protein called Myeloperoxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	465	Total	C	N	O	S	0	0	0
			3728	2348	687	666	27			


- Molecule 3 is a protein called Myeloperoxidase inhibitor SPIN.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	69	Total	C	N	O	0	0	0
			557	352	94	111			
3	F	70	Total	C	N	O	0	0	0
			566	357	95	114			

- Molecule 4 is a protein called Myeloperoxidase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	458	Total	C	N	O	S	0	0	0
			3683	2323	680	655	25			

- Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	G	6	Total	C	N	O	0	0	0
			71	40	2	29			

- Molecule 6 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	H	2	Total	C	N	O	0	0	0
			28	16	2	10			
6	I	2	Total	C	N	O	0	0	0
			28	16	2	10			
6	L	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 7 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
7	J	7	Total	C	N	O	0	0	0
			85	48	3	34			

- Molecule 8 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
8	K	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 9 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	1	Total Cl 1 1	0	0
9	C	1	Total Cl 1 1	0	0

- Molecule 10 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: C₃₄H₃₂FeN₄O₄) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	A	1	Total C Fe N O 43 34 1 4 4	0	0
10	C	1	Total C Fe N O 43 34 1 4 4	0	0

- Molecule 11 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	B	1	Total Ca 1 1	0	0
11	D	1	Total Ca 1 1	0	0

- Molecule 12 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
12	B	3	Total 3	O 3	0	0
12	D	1	Total 1	O 1	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Myeloperoxidase





- Molecule 1: Myeloperoxidase




- Molecule 2: Myeloperoxidase




- Molecule 3: Myeloperoxidase inhibitor SPIN





- Molecule 3: Myeloperoxidase inhibitor SPIN






• Molecule 4: Myeloperoxidase



• Molecule 5: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose




• Molecule 6: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




• Molecule 6: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



• Molecule 6: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




• Molecule 7: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J:  14% 43% 43%



- Molecule 8: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K:  100%



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	111.47Å 111.47Å 241.91Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	78.95 – 2.55 78.95 – 2.55	Depositor EDS
% Data completeness (in resolution range)	60.2 (78.95-2.55) 60.2 (78.95-2.55)	Depositor EDS
R_{merge}	0.21	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.97 (at 2.55Å)	Xtriage
Refinement program	REFMAC 5.8.0431	Depositor
R, R_{free}	0.202 , 0.264 0.203 , 0.264	Depositor DCC
R_{free} test set	1509 reflections (2.99%)	wwPDB-VP
Wilson B-factor (Å ²)	53.1	Xtriage
Anisotropy	0.100	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 47.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	10519	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, FUC, YCM, CL, HEM, MAN, CA, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.63	0/828	1.13	1/1126 (0.1%)
1	C	0.57	0/810	1.09	2/1101 (0.2%)
2	B	0.58	0/3803	1.06	7/5158 (0.1%)
3	E	0.51	0/566	1.11	0/761
3	F	0.49	0/575	1.08	0/773
4	D	0.54	0/3747	1.03	5/5080 (0.1%)
All	All	0.56	0/10329	1.06	15/13999 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	6
3	F	0	1
4	D	0	2
All	All	0	9

There are no bond length outliers.

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	622	ARG	CB-CA-C	-6.37	102.51	111.80
2	B	413	THR	CA-CB-OG1	-6.04	100.54	109.60
4	D	593	ASP	CA-CB-CG	5.75	118.36	112.60
2	B	577	MET	CG-SD-CE	-5.60	88.58	100.90
2	B	544	ASP	CA-CB-CG	5.47	118.07	112.60
4	D	438	ASP	CA-CB-CG	5.39	117.99	112.60
2	B	380	ASP	CA-CB-CG	5.38	117.98	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	438	ASP	CA-CB-CG	5.32	117.92	112.60
1	A	262	ASP	CA-CB-CG	5.30	117.90	112.60
1	C	202	GLU	CB-CA-C	5.24	119.33	110.79
4	D	487	ASP	CA-CB-CG	5.23	117.83	112.60
2	B	487	ASP	CA-CB-CG	5.18	117.78	112.60
2	B	593	ASP	CA-CB-CG	5.14	117.74	112.60
4	D	524	ARG	N-CA-CB	-5.09	102.55	110.55
1	C	262	ASP	CA-CB-CG	5.03	117.62	112.60

There are no chirality outliers.

All (9) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	302	ARG	Sidechain
2	B	351	ARG	Sidechain
2	B	392	ARG	Sidechain
2	B	473	ARG	Sidechain
2	B	524	ARG	Sidechain
2	B	622	ARG	Sidechain
4	D	480	ARG	Sidechain
4	D	524	ARG	Sidechain
3	F	30	LYS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	815	0	778	10	0
1	C	797	0	764	5	0
2	B	3728	0	3720	36	0
3	E	557	0	540	4	0
3	F	566	0	546	10	0
4	D	3683	0	3683	24	0
5	G	71	0	61	0	0
6	H	28	0	25	3	0
6	I	28	0	25	0	0
6	L	28	0	25	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	J	85	0	73	2	0
8	K	39	0	34	0	0
9	A	1	0	0	0	0
9	C	1	0	0	0	0
10	A	43	0	30	1	0
10	C	43	0	30	1	0
11	B	1	0	0	0	0
11	D	1	0	0	0	0
12	B	3	0	0	0	0
12	D	1	0	0	0	0
All	All	10519	0	10334	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:502:HIS:HD1	2:B:587:ASN:HD21	1.25	0.84
4:D:502:HIS:HD1	4:D:587:ASN:HD21	1.25	0.82
2:B:319:YCM:HB3	4:D:319:YCM:SG	2.25	0.77
4:D:357:SER:O	3:F:65:THR:HG23	1.93	0.67
1:C:268:GLU:HG2	3:F:33:SER:O	2.00	0.61
1:A:262:ASP:HB2	2:B:341:MET:CE	2.31	0.60
1:A:233:GLU:HG2	2:B:633:GLN:HE21	1.67	0.59
2:B:365:VAL:H	6:H:1:NAG:H81	1.68	0.59
2:B:577:MET:HE1	2:B:581:LEU:HD11	1.86	0.57
4:D:392:ARG:HG3	4:D:392:ARG:HH11	1.68	0.57
4:D:381:ASN:HD21	3:F:91:ASP:HB3	1.70	0.56
1:A:194:ALA:HB1	2:B:328:ASN:HD22	1.71	0.56
4:D:720:SER:HB3	4:D:726:ASP:HB3	1.88	0.54
2:B:720:SER:HB3	2:B:726:ASP:HB3	1.90	0.54
2:B:364:ALA:HA	6:H:1:NAG:C8	2.39	0.53
2:B:346:GLU:OE2	3:E:32:TYR:OH	2.27	0.53
10:A:302:HEM:HBA2	2:B:499:ARG:NH1	2.24	0.53
2:B:347:GLU:N	2:B:348:PRO:HD2	2.23	0.52
2:B:628:ARG:HH11	2:B:628:ARG:HG2	1.74	0.52
4:D:405:ARG:HG2	3:F:39:LEU:HD13	1.91	0.51
7:J:2:NAG:H3	7:J:5:NAG:H3	1.92	0.51
1:C:182:ASN:O	1:C:186:PRO:HA	2.11	0.50
2:B:364:ALA:HA	6:H:1:NAG:H81	1.94	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:305:ASN:HD22	2:B:308:ASP:HB2	1.77	0.48
1:A:241:GLN:O	2:B:562:GLN:NE2	2.46	0.48
2:B:294:LEU:HB2	2:B:310:ILE:HB	1.95	0.47
4:D:699:LEU:N	4:D:700:PRO:CD	2.78	0.47
1:A:247:GLU:O	2:B:719:MET:HG3	2.15	0.47
4:D:628:ARG:HG2	4:D:628:ARG:HH11	1.80	0.47
2:B:422:GLU:OE1	2:B:706:ASN:ND2	2.36	0.47
2:B:409:MET:HE2	2:B:411:GLU:CD	2.40	0.46
4:D:513:ASP:HA	4:D:519:MET:HE3	1.98	0.46
4:D:409:MET:HE2	4:D:411:GLU:CD	2.41	0.46
2:B:714:LYS:HE2	2:B:726:ASP:HA	1.98	0.46
1:A:182:ASN:O	1:A:186:PRO:HA	2.16	0.45
2:B:425:ARG:HG2	2:B:425:ARG:HH11	1.80	0.45
2:B:382:LEU:HD21	3:E:40:HIS:HA	1.99	0.44
1:C:195:PHE:CE1	4:D:331:ASN:HB2	2.52	0.44
2:B:699:LEU:N	2:B:700:PRO:CD	2.80	0.44
1:A:195:PHE:CE1	2:B:331:ASN:HB2	2.53	0.44
4:D:535:TRP:CH2	6:L:1:NAG:H2	2.52	0.44
4:D:404:THR:O	3:F:39:LEU:HB3	2.18	0.44
4:D:688:MET:HE3	4:D:692:GLN:OE1	2.18	0.43
2:B:281:CYS:HB2	2:B:313:PHE:CE2	2.54	0.43
4:D:381:ASN:CB	3:F:45:PHE:O	2.67	0.43
2:B:281:CYS:HB2	2:B:313:PHE:CZ	2.54	0.42
3:E:66:LYS:O	3:E:70:ARG:HG2	2.19	0.42
4:D:450:VAL:O	4:D:454:VAL:HG23	2.19	0.42
4:D:463:LEU:N	4:D:464:PRO:CD	2.82	0.42
2:B:603:ARG:NH1	2:B:649:GLU:OE2	2.49	0.42
4:D:381:ASN:HB3	3:F:45:PHE:O	2.20	0.42
2:B:513:ASP:HA	2:B:519:MET:HE3	2.02	0.42
2:B:742:ARG:HG2	2:B:742:ARG:HH11	1.84	0.42
10:C:302:HEM:HMC1	10:C:302:HEM:HAC	1.89	0.42
2:B:319:YCM:HB2	4:D:325:THR:HG21	2.02	0.41
2:B:463:LEU:N	2:B:464:PRO:CD	2.83	0.41
3:E:89:LYS:HB2	3:E:95:VAL:HG21	2.02	0.41
1:A:262:ASP:CB	2:B:341:MET:HE3	2.51	0.41
1:A:195:PHE:CZ	2:B:495:THR:HG21	2.56	0.41
4:D:319:YCM:HA	4:D:319:YCM:HD3	1.88	0.41
2:B:474:LYS:NZ	7:J:6:MAN:O6	2.45	0.41
2:B:723:TYR:HA	2:B:724:PRO:HA	1.87	0.41
1:C:188:LEU:HB3	4:D:488:PRO:HD2	2.03	0.41
1:A:236:ARG:HH11	1:A:236:ARG:HD3	1.76	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:F:66:LYS:O	3:F:70:ARG:HG2	2.21	0.41
4:D:294:LEU:HB2	4:D:310:ILE:HB	2.03	0.40
4:D:356:MET:HG3	3:F:72:TYR:CD1	2.55	0.40
1:C:268:GLU:OE2	3:F:35:ASN:N	2.47	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	98/114 (86%)	95 (97%)	3 (3%)	0	100	100
1	C	95/114 (83%)	92 (97%)	3 (3%)	0	100	100
2	B	462/467 (99%)	448 (97%)	13 (3%)	1 (0%)	43	56
3	E	67/73 (92%)	67 (100%)	0	0	100	100
3	F	68/73 (93%)	67 (98%)	1 (2%)	0	100	100
4	D	454/467 (97%)	435 (96%)	16 (4%)	3 (1%)	18	25
All	All	1244/1308 (95%)	1204 (97%)	36 (3%)	4 (0%)	36	45

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	321	GLY
4	D	321	GLY
4	D	523	PRO
4	D	521	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	85/96 (88%)	84 (99%)	1 (1%)	63	78
1	C	84/96 (88%)	83 (99%)	1 (1%)	63	78
2	B	409/411 (100%)	407 (100%)	2 (0%)	81	88
3	E	61/65 (94%)	60 (98%)	1 (2%)	55	73
3	F	62/65 (95%)	61 (98%)	1 (2%)	55	73
4	D	401/410 (98%)	399 (100%)	2 (0%)	81	88
All	All	1102/1143 (96%)	1094 (99%)	8 (1%)	76	85

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	239	THR
2	B	322	SER
2	B	663	CYS
3	E	31	VAL
1	C	239	THR
4	D	341	MET
4	D	663	CYS
3	F	31	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	220	ASN
1	A	241	GLN
2	B	633	GLN
2	B	692	GLN
2	B	715	ASN
3	E	90	GLN
1	C	192	ASN
1	C	241	GLN
1	C	261	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
4	D	305	ASN
4	D	517	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

5 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	YCM	C	180	1	7,9,10	0.47	0	4,10,12	0.59	0
2	YCM	B	319	2	7,9,10	0.54	0	4,10,12	0.86	0
4	YCM	D	309	4	7,9,10	0.46	0	4,10,12	0.44	0
4	YCM	D	319	4	7,9,10	0.69	0	4,10,12	0.99	0
1	YCM	A	180	1	7,9,10	0.47	0	4,10,12	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	YCM	C	180	1	-	1/6/8/10	-
2	YCM	B	319	2	-	4/6/8/10	-
4	YCM	D	309	4	-	1/6/8/10	-
4	YCM	D	319	4	-	2/6/8/10	-
1	YCM	A	180	1	-	3/6/8/10	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	180	YCM	CE-CD-SG-CB
1	A	180	YCM	SG-CD-CE-OZ1
1	A	180	YCM	SG-CD-CE-NZ2
2	B	319	YCM	N-CA-CB-SG
2	B	319	YCM	C-CA-CB-SG
4	D	309	YCM	CE-CD-SG-CB
4	D	319	YCM	CE-CD-SG-CB
2	B	319	YCM	SG-CD-CE-OZ1
2	B	319	YCM	SG-CD-CE-NZ2
1	C	180	YCM	SG-CD-CE-OZ1
4	D	319	YCM	SG-CD-CE-OZ1

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	319	YCM	2	0
4	D	319	YCM	2	0

5.5 Carbohydrates

22 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	G	1	2,5	14,14,15	0.50	0	17,19,21	1.50	3 (17%)
5	NAG	G	2	5	14,14,15	0.32	0	17,19,21	1.09	2 (11%)
5	BMA	G	3	5	11,11,12	0.55	0	15,15,17	0.90	0
5	MAN	G	4	5	11,11,12	0.33	0	15,15,17	1.13	1 (6%)
5	MAN	G	5	5	11,11,12	0.37	0	15,15,17	0.93	0
5	FUC	G	6	5	10,10,11	0.62	0	14,14,16	0.66	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	NAG	H	1	6,2	14,14,15	0.31	0	17,19,21	1.22	1 (5%)
6	NAG	H	2	6	14,14,15	0.34	0	17,19,21	1.01	1 (5%)
6	NAG	I	1	6,2	14,14,15	0.37	0	17,19,21	0.74	0
6	NAG	I	2	6	14,14,15	0.36	0	17,19,21	1.50	2 (11%)
7	NAG	J	1	7,4	14,14,15	0.49	0	17,19,21	1.09	2 (11%)
7	NAG	J	2	7	14,14,15	0.38	0	17,19,21	1.52	2 (11%)
7	BMA	J	3	7	11,11,12	0.85	1 (9%)	15,15,17	0.96	0
7	MAN	J	4	7	11,11,12	0.37	0	15,15,17	1.42	2 (13%)
7	NAG	J	5	7	14,14,15	0.28	0	17,19,21	1.10	1 (5%)
7	MAN	J	6	7	11,11,12	0.61	0	15,15,17	1.12	1 (6%)
7	FUC	J	7	7	10,10,11	0.60	0	14,14,16	0.56	0
8	NAG	K	1	8,4	14,14,15	0.31	0	17,19,21	0.98	1 (5%)
8	NAG	K	2	8	14,14,15	0.37	0	17,19,21	0.94	2 (11%)
8	BMA	K	3	8	11,11,12	0.50	0	15,15,17	0.82	1 (6%)
6	NAG	L	1	4,6	14,14,15	0.34	0	17,19,21	1.40	1 (5%)
6	NAG	L	2	6	14,14,15	0.34	0	17,19,21	0.88	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	G	1	2,5	-	4/6/23/26	0/1/1/1
5	NAG	G	2	5	-	2/6/23/26	0/1/1/1
5	BMA	G	3	5	-	0/2/19/22	0/1/1/1
5	MAN	G	4	5	-	2/2/19/22	0/1/1/1
5	MAN	G	5	5	-	2/2/19/22	0/1/1/1
5	FUC	G	6	5	-	-	0/1/1/1
6	NAG	H	1	6,2	-	2/6/23/26	0/1/1/1
6	NAG	H	2	6	-	1/6/23/26	0/1/1/1
6	NAG	I	1	6,2	-	2/6/23/26	0/1/1/1
6	NAG	I	2	6	-	2/6/23/26	0/1/1/1
7	NAG	J	1	7,4	-	2/6/23/26	0/1/1/1
7	NAG	J	2	7	-	2/6/23/26	0/1/1/1
7	BMA	J	3	7	-	0/2/19/22	0/1/1/1
7	MAN	J	4	7	-	2/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	NAG	J	5	7	-	4/6/23/26	0/1/1/1
7	MAN	J	6	7	-	2/2/19/22	0/1/1/1
7	FUC	J	7	7	-	-	0/1/1/1
8	NAG	K	1	8,4	-	0/6/23/26	0/1/1/1
8	NAG	K	2	8	-	2/6/23/26	0/1/1/1
8	BMA	K	3	8	-	1/2/19/22	0/1/1/1
6	NAG	L	1	4,6	-	3/6/23/26	0/1/1/1
6	NAG	L	2	6	-	3/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	J	3	BMA	C2-C3	-2.05	1.49	1.52

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	L	1	NAG	C1-C2-N2	4.98	119.00	110.49
6	I	2	NAG	C1-C2-N2	4.11	117.50	110.49
7	J	4	MAN	C1-C2-C3	4.08	114.69	109.67
7	J	2	NAG	C2-N2-C7	3.82	128.34	122.90
6	H	1	NAG	C1-C2-N2	3.71	116.82	110.49
5	G	1	NAG	C1-C2-N2	-3.70	104.17	110.49
7	J	2	NAG	C1-O5-C5	3.68	117.18	112.19
7	J	1	NAG	O5-C1-C2	-3.19	106.26	111.29
7	J	5	NAG	C1-O5-C5	3.10	116.40	112.19
7	J	6	MAN	C1-O5-C5	3.05	116.33	112.19
5	G	4	MAN	C1-O5-C5	2.94	116.17	112.19
6	I	2	NAG	C4-C3-C2	-2.93	106.73	111.02
8	K	2	NAG	C1-O5-C5	2.72	115.88	112.19
8	K	3	BMA	C1-O5-C5	2.72	115.87	112.19
7	J	4	MAN	C1-O5-C5	2.70	115.84	112.19
8	K	1	NAG	C1-O5-C5	2.62	115.74	112.19
5	G	1	NAG	C2-N2-C7	2.59	126.59	122.90
6	L	2	NAG	C1-C2-N2	2.50	114.75	110.49
5	G	1	NAG	O5-C5-C6	-2.42	103.41	107.20
5	G	2	NAG	C1-O5-C5	-2.26	109.14	112.19
5	G	2	NAG	C1-C2-N2	2.25	114.34	110.49
8	K	2	NAG	C1-C2-N2	2.13	114.12	110.49
6	H	2	NAG	C4-C3-C2	2.11	114.10	111.02
7	J	1	NAG	C1-C2-N2	2.07	114.03	110.49

There are no chirality outliers.

All (38) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	G	1	NAG	C8-C7-N2-C2
5	G	1	NAG	O7-C7-N2-C2
6	H	1	NAG	C8-C7-N2-C2
6	I	1	NAG	C8-C7-N2-C2
6	I	1	NAG	O7-C7-N2-C2
6	L	1	NAG	C1-C2-N2-C7
6	L	1	NAG	C8-C7-N2-C2
6	L	1	NAG	O7-C7-N2-C2
6	L	2	NAG	C1-C2-N2-C7
6	L	2	NAG	C8-C7-N2-C2
6	L	2	NAG	O7-C7-N2-C2
6	H	1	NAG	O7-C7-N2-C2
5	G	5	MAN	O5-C5-C6-O6
7	J	5	NAG	O5-C5-C6-O6
7	J	6	MAN	O5-C5-C6-O6
7	J	2	NAG	C8-C7-N2-C2
5	G	5	MAN	C4-C5-C6-O6
7	J	6	MAN	C4-C5-C6-O6
6	I	2	NAG	C8-C7-N2-C2
7	J	2	NAG	O7-C7-N2-C2
7	J	5	NAG	C8-C7-N2-C2
8	K	2	NAG	C8-C7-N2-C2
5	G	1	NAG	C1-C2-N2-C7
6	I	2	NAG	O7-C7-N2-C2
7	J	5	NAG	O7-C7-N2-C2
8	K	2	NAG	O7-C7-N2-C2
5	G	4	MAN	O5-C5-C6-O6
7	J	5	NAG	C4-C5-C6-O6
6	H	2	NAG	O5-C5-C6-O6
7	J	4	MAN	C4-C5-C6-O6
7	J	1	NAG	O7-C7-N2-C2
7	J	1	NAG	C8-C7-N2-C2
7	J	4	MAN	O5-C5-C6-O6
5	G	2	NAG	C8-C7-N2-C2
8	K	3	BMA	O5-C5-C6-O6
5	G	2	NAG	O7-C7-N2-C2
5	G	4	MAN	C4-C5-C6-O6
5	G	1	NAG	C3-C2-N2-C7


There are no ring outliers.


5 monomers are involved in 6 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	J	2	NAG	1	0
7	J	5	NAG	1	0
6	H	1	NAG	3	0
7	J	6	MAN	1	0
6	L	1	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.














5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
10	HEM	C	302	4,1	50,50,50	2.48	19 (38%)	66,82,82	1.45	9 (13%)
10	HEM	A	302	2,1	50,50,50	1.78	11 (22%)	66,82,82	1.68	16 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	HEM	C	302	4,1	-	3/14/54/54	-
10	HEM	A	302	2,1	-	6/14/54/54	-

All (30) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
10	C	302	HEM	FE-NB	8.47	2.21	1.94
10	C	302	HEM	FE-NC	7.63	2.21	1.95
10	C	302	HEM	C4D-C3D	-4.55	1.37	1.45
10	C	302	HEM	FE-NA	4.10	2.09	1.95
10	A	302	HEM	FE-NA	4.04	2.08	1.95
10	A	302	HEM	C4D-C3D	-4.04	1.38	1.45
10	A	302	HEM	C1B-C2B	-3.92	1.37	1.44
10	C	302	HEM	C1C-C2C	-3.83	1.37	1.45
10	C	302	HEM	C3C-C4C	-3.59	1.39	1.46
10	A	302	HEM	C3C-C4C	-3.44	1.40	1.46
10	C	302	HEM	C1A-C2A	-3.43	1.37	1.44
10	A	302	HEM	C1D-C2D	-3.32	1.38	1.44
10	C	302	HEM	C1D-C2D	-3.27	1.38	1.44
10	C	302	HEM	C3B-C2B	3.19	1.43	1.37
10	A	302	HEM	FE-ND	3.06	2.04	1.94
10	A	302	HEM	C1A-C2A	-3.06	1.38	1.44
10	C	302	HEM	C4A-C3A	-2.97	1.36	1.43
10	C	302	HEM	C3B-C4B	-2.97	1.38	1.44
10	A	302	HEM	C1C-C2C	-2.85	1.39	1.45
10	C	302	HEM	C1B-C2B	-2.82	1.39	1.44
10	A	302	HEM	C3B-C4B	-2.81	1.39	1.44
10	A	302	HEM	C1B-NB	-2.80	1.35	1.40
10	C	302	HEM	C4D-ND	-2.51	1.36	1.40
10	C	302	HEM	C1B-NB	-2.46	1.36	1.40
10	C	302	HEM	CHD-C4C	2.42	1.43	1.38
10	C	302	HEM	C1A-NA	-2.40	1.35	1.39
10	C	302	HEM	CHD-C1D	2.38	1.44	1.39
10	A	302	HEM	C4D-ND	-2.14	1.36	1.40
10	C	302	HEM	C4A-NA	-2.11	1.35	1.39
10	C	302	HEM	O2D-CGD	-2.05	1.23	1.30

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
10	C	302	HEM	CAC-C3C-C4C	4.21	135.07	124.90
10	A	302	HEM	CAC-C3C-C4C	4.18	134.99	124.90
10	C	302	HEM	CHD-C4C-NC	-4.10	120.00	124.44
10	A	302	HEM	C3B-C2B-C1B	3.88	109.37	106.49
10	C	302	HEM	CHD-C4C-C3C	3.53	131.30	125.26
10	A	302	HEM	CHC-C4B-NB	-3.52	120.59	124.42
10	A	302	HEM	CAB-C3B-C2B	-3.45	117.24	128.60
10	C	302	HEM	CAC-C3C-C2C	-3.09	118.42	128.60
10	A	302	HEM	CHD-C4C-NC	-3.00	121.19	124.44
10	A	302	HEM	CMD-C2D-C1D	2.96	129.54	125.04
10	A	302	HEM	CAC-C3C-C2C	-2.88	119.10	128.60
10	A	302	HEM	CAB-C3B-C4B	2.87	137.85	124.47
10	A	302	HEM	C4B-C3B-C2B	-2.78	104.91	107.11
10	C	302	HEM	O2D-CGD-CBD	2.70	122.69	114.03
10	A	302	HEM	CHD-C1D-C2D	2.52	128.92	124.98
10	A	302	HEM	CAD-CBD-CGD	2.45	118.87	113.60
10	C	302	HEM	O1A-CGA-CBA	-2.41	115.35	123.08
10	C	302	HEM	CAD-CBD-CGD	2.38	118.72	113.60
10	A	302	HEM	CMC-C2C-C1C	2.30	128.77	124.71
10	C	302	HEM	O1D-CGD-CBD	-2.24	115.89	123.08
10	A	302	HEM	CMC-C2C-C3C	-2.22	122.86	128.30
10	A	302	HEM	CHD-C1D-ND	-2.13	122.10	124.42
10	A	302	HEM	O2D-CGD-CBD	2.09	120.76	114.03
10	A	302	HEM	CMA-C3A-C4A	2.08	128.54	125.37
10	C	302	HEM	O2A-CGA-CBA	2.06	120.65	114.03

There are no chirality outliers.

All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
10	A	302	HEM	C4B-C3B-CAB-CBB
10	A	302	HEM	C4C-C3C-CAC-CBC
10	C	302	HEM	C4C-C3C-CAC-CBC
10	A	302	HEM	CAA-CBA-CGA-O2A
10	A	302	HEM	CAD-CBD-CGD-O1D
10	A	302	HEM	CAA-CBA-CGA-O1A
10	A	302	HEM	CAD-CBD-CGD-O2D
10	C	302	HEM	CAD-CBD-CGD-O1D
10	C	302	HEM	CAD-CBD-CGD-O2D


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
10	C	302	HEM	1	0
10	A	302	HEM	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	100/114 (87%)	-0.27	1 (1%) 79 80	24, 41, 63, 90	0
1	C	97/114 (85%)	0.06	0 100 100	29, 52, 84, 93	0
2	B	464/467 (99%)	-0.21	3 (0%) 85 88	23, 44, 74, 119	0
3	E	69/73 (94%)	0.72	5 (7%) 21 20	59, 77, 98, 104	0
3	F	70/73 (95%)	1.93	28 (40%) 1 0	109, 135, 155, 158	0
4	D	456/467 (97%)	0.45	13 (2%) 53 55	33, 70, 105, 126	0
All	All	1256/1308 (96%)	0.22	50 (3%) 42 43	23, 55, 120, 158	0

All (50) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	F	81	ILE	6.3
3	F	39	LEU	4.5
3	F	31	VAL	4.2
3	F	38	VAL	3.8
3	F	57	LEU	3.7
3	F	43	ALA	3.6
3	F	30	LYS	3.6
2	B	715	ASN	3.6
3	F	35	ASN	3.6
4	D	521	PRO	3.5
1	A	271	ALA	3.3
3	E	31	VAL	3.1
3	F	32	TYR	3.1
3	F	48	HIS	2.9
3	F	78	LEU	2.9
3	F	40	HIS	2.9
4	D	573	PHE	2.9
4	D	433	LEU	2.8
3	F	84	ILE	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	F	93	PHE	2.8
4	D	435	PRO	2.8
4	D	303	ILE	2.7
4	D	550	LEU	2.7
3	E	38	VAL	2.7
3	F	55	VAL	2.7
3	F	56	LEU	2.6
3	F	64	ALA	2.6
3	F	52	TYR	2.6
3	F	42	ASP	2.6
3	F	85	ILE	2.6
4	D	715	ASN	2.6
3	F	46	LEU	2.5
4	D	532	PHE	2.5
4	D	744	ALA	2.4
3	F	53	ILE	2.3
3	F	69	LEU	2.3
4	D	321	GLY	2.3
3	E	32	TYR	2.3
3	E	41	ASP	2.3
3	F	41	ASP	2.3
3	F	34	GLN	2.2
3	F	95	VAL	2.2
3	F	98	TYR	2.2
3	E	81	ILE	2.1
4	D	473	ARG	2.1
3	F	45	PHE	2.1
2	B	323	ASN	2.1
2	B	744	ALA	2.0
4	D	392	ARG	2.0
4	D	733	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	YCM	B	319	10/11	0.77	0.19	61,82,109,128	0
4	YCM	D	319	10/11	0.80	0.14	60,74,103,105	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	YCM	D	309	10/11	0.82	0.13	79,82,85,85	0
1	YCM	A	180	10/11	0.89	0.15	51,60,88,102	0
1	YCM	C	180	10/11	0.92	0.11	59,73,97,98	0

6.3 Carbohydrates




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
8	BMA	K	3	11/12	0.45	0.17	111,114,117,117	0
6	NAG	L	2	14/15	0.60	0.15	121,131,138,141	0
8	NAG	K	2	14/15	0.64	0.15	103,113,120,126	0
7	NAG	J	5	14/15	0.65	0.20	72,85,94,102	0
6	NAG	I	2	14/15	0.70	0.13	83,95,102,102	0
5	MAN	G	4	11/12	0.72	0.12	61,63,66,68	0
6	NAG	H	2	14/15	0.77	0.11	75,80,85,86	0
6	NAG	L	1	14/15	0.77	0.14	103,109,117,129	0
8	NAG	K	1	14/15	0.84	0.11	76,93,100,107	0
7	MAN	J	6	11/12	0.85	0.09	59,61,67,68	0
6	NAG	I	1	14/15	0.88	0.12	58,66,80,82	0
5	MAN	G	5	11/12	0.89	0.09	45,48,50,53	0
6	NAG	H	1	14/15	0.89	0.08	49,57,61,69	0
5	NAG	G	1	14/15	0.92	0.08	34,39,45,47	0
7	MAN	J	4	11/12	0.93	0.09	48,50,55,67	0
7	NAG	J	1	14/15	0.94	0.07	39,41,45,45	0
5	NAG	G	2	14/15	0.95	0.08	36,38,42,43	0
7	FUC	J	7	10/11	0.95	0.09	45,49,55,56	0
7	BMA	J	3	11/12	0.95	0.08	42,44,49,52	0
5	BMA	G	3	11/12	0.95	0.07	46,48,51,54	0
5	FUC	G	6	10/11	0.95	0.07	32,33,35,35	0
7	NAG	J	2	14/15	0.96	0.07	35,39,41,41	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain G:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain H:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain I:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain L:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain J:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
9	CL	A	301	1/1	0.95	0.06	32,32,32,32	0
9	CL	C	301	1/1	0.95	0.06	42,42,42,42	0
10	HEM	C	302	43/43	0.95	0.10	38,67,77,86	0
10	HEM	A	302	43/43	0.98	0.06	20,28,33,37	0
11	CA	B	801	1/1	0.98	0.02	23,23,23,23	0
11	CA	D	801	1/1	0.99	0.03	46,46,46,46	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around HEM C 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.