



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 11:14 AM UTC

PDB ID : 3A0F / pdb_00003a0f
Title : The crystal structure of Geotrichum sp. M128 xyloglucanase
Authors : Yaoi, K.; Kondo, H.; Hiyoshi, A.; Noro, N.; Sugimoto, H.; Tsuda, S.; Miyazaki, K.
Deposited on : 2009-03-16
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	763	3% 77% 15% 5% ..

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5783 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Xyloglucanase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	753	Total	C	N	O	S	0	0	0
			5676	3589	976	1098	13			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	MET	-	expression tag	UNP Q764N8
A	-5	HIS	-	expression tag	UNP Q764N8
A	-4	HIS	-	expression tag	UNP Q764N8
A	-3	HIS	-	expression tag	UNP Q764N8
A	-2	HIS	-	expression tag	UNP Q764N8
A	-1	HIS	-	expression tag	UNP Q764N8
A	0	HIS	-	expression tag	UNP Q764N8


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	107	Total	O	0	0
			107	107		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Xyloglucanase



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	135.19Å 135.19Å 119.89Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 2.50 20.00 – 2.50	Depositor EDS
% Data completeness (in resolution range)	98.5 (20.00-2.50) 86.2 (20.00-2.50)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.44 (at 2.50Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.236 , 0.276 (Not available) , 0.225	Depositor DCC
R_{free} test set	2192 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	18.4	Xtriage
Anisotropy	0.299	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 25.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.023 for -h,-k,l	Xtriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	5783	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.89	1/5843 (0.0%)	1.04	20/7989 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	327	GLY	N-CA	5.33	1.53	1.45

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	22	HIS	CA-C-N	-8.13	111.41	119.87
1	A	22	HIS	C-N-CA	-8.13	111.41	119.87
1	A	290	ALA	N-CA-C	7.37	121.87	107.57
1	A	10	ILE	N-CA-C	-7.34	105.82	111.62
1	A	283	ILE	CB-CA-C	7.05	118.24	110.62
1	A	234	ILE	CB-CA-C	-6.71	104.21	111.59
1	A	335	ARG	N-CA-C	-6.43	104.75	112.59
1	A	79	ASN	CA-C-N	6.26	125.95	119.56
1	A	79	ASN	C-N-CA	6.26	125.95	119.56
1	A	739	VAL	CB-CA-C	-5.96	101.16	110.69
1	A	499	GLY	N-CA-C	5.92	119.77	111.19
1	A	204	ASP	N-CA-C	-5.85	104.69	112.94
1	A	32	ARG	NE-CZ-NH2	5.81	124.43	119.20
1	A	63	ASN	N-CA-C	5.58	117.45	107.80
1	A	625	TRP	N-CA-C	5.43	117.09	110.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	283	ILE	N-CA-CB	-5.27	106.06	112.33
1	A	325	ALA	N-CA-C	-5.26	107.04	112.93
1	A	583	THR	CB-CA-C	-5.19	99.30	110.45
1	A	61	ALA	N-CA-C	-5.18	103.65	110.33
1	A	553	TRP	CA-CB-CG	5.17	123.41	113.60

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	127	GLY	Peptide
1	A	290	ALA	Peptide
1	A	622	PRO	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5676	0	5394	114	0
2	A	107	0	0	2	0
All	All	5783	0	5394	114	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (114) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:460:GLY:O	1:A:482:THR:HG23	1.72	0.90
1:A:726:ALA:HA	1:A:743:MET:HE3	1.53	0.88
1:A:344:LEU:O	1:A:348:THR:HG23	1.80	0.82
1:A:319:VAL:HG13	1:A:393:ILE:HD12	1.62	0.82
1:A:539:GLY:O	1:A:553:TRP:CZ3	2.34	0.81
1:A:439:ILE:HD11	1:A:742:GLY:C	2.09	0.78
1:A:663:LEU:O	1:A:666:SER:HB2	1.86	0.76

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:690:PHE:HE2	1:A:705:MET:HE2	1.53	0.74
1:A:128:GLY:HA2	2:A:758:HOH:O	1.88	0.73
1:A:171:THR:HG21	2:A:760:HOH:O	1.87	0.73
1:A:171:THR:HG23	1:A:194:HIS:HD2	1.53	0.73
1:A:539:GLY:O	1:A:553:TRP:CH2	2.42	0.73
1:A:61:ALA:O	1:A:62:GLY:O	2.07	0.72
1:A:553:TRP:HD1	1:A:573:TRP:CH2	2.08	0.72
1:A:726:ALA:HA	1:A:743:MET:CE	2.21	0.69
1:A:609:ASP:OD2	1:A:613:SER:OG	2.12	0.67
1:A:34:ASP:OD1	1:A:125:ARG:NH1	2.27	0.67
1:A:645:VAL:HG12	1:A:648:VAL:HG13	1.77	0.66
1:A:283:ILE:HD12	1:A:338:GLY:O	1.95	0.66
1:A:38:THR:HG22	1:A:70:THR:OG1	1.96	0.66
1:A:285:PRO:HD2	1:A:307:TRP:CH2	2.30	0.66
1:A:539:GLY:O	1:A:553:TRP:HZ3	1.79	0.65
1:A:439:ILE:HG13	1:A:742:GLY:HA3	1.78	0.65
1:A:534:LEU:HD21	1:A:553:TRP:CH2	2.32	0.64
1:A:679:ALA:O	1:A:688:THR:HG23	1.97	0.63
1:A:61:ALA:O	1:A:62:GLY:C	2.41	0.63
1:A:553:TRP:HD1	1:A:573:TRP:CZ2	2.15	0.63
1:A:344:LEU:O	1:A:348:THR:CG2	2.46	0.63
1:A:642:TRP:HE1	1:A:709:ASN:HD21	1.46	0.62
1:A:647:GLY:HA2	1:A:671:ASN:OD1	2.00	0.61
1:A:176:ILE:HG23	1:A:191:VAL:HG13	1.82	0.61
1:A:462:MET:HG2	1:A:464:HIS:CE1	2.36	0.60
1:A:622:PRO:HD3	1:A:651:TYR:CZ	2.38	0.59
1:A:439:ILE:CG1	1:A:742:GLY:HA3	2.33	0.59
1:A:485:LEU:HA	1:A:542:ILE:HD11	1.85	0.58
1:A:726:ALA:HB2	1:A:743:MET:HE1	1.85	0.58
1:A:497:ARG:HH21	1:A:524:TRP:CD1	2.23	0.57
1:A:176:ILE:CG2	1:A:191:VAL:HG13	2.34	0.57
1:A:593:ALA:O	1:A:595:THR:HG22	2.05	0.57
1:A:283:ILE:HG23	1:A:341:TRP:HZ2	1.69	0.56
1:A:476:LYS:HA	1:A:505:ASP:OD2	2.05	0.56
1:A:679:ALA:O	1:A:688:THR:CG2	2.54	0.56
1:A:287:ARG:NH2	1:A:290:ALA:HB3	2.22	0.55
1:A:705:MET:HE1	1:A:737:GLY:HA2	1.89	0.54
1:A:226:ILE:HD11	1:A:304:PRO:HD3	1.89	0.54
1:A:439:ILE:HD11	1:A:743:MET:N	2.23	0.53
1:A:705:MET:HE3	1:A:716:LEU:HD21	1.89	0.53
1:A:583:THR:HB	1:A:585:GLY:H	1.73	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:498:VAL:HG22	1:A:553:TRP:CZ2	2.43	0.52
1:A:553:TRP:CD1	1:A:573:TRP:CH2	2.94	0.52
1:A:439:ILE:CD1	1:A:743:MET:N	2.73	0.52
1:A:281:LYS:HD2	1:A:283:ILE:HD11	1.92	0.52
1:A:140:GLU:HA	1:A:153:SER:O	2.11	0.51
1:A:100:LEU:HB3	1:A:109:PHE:CD2	2.46	0.50
1:A:595:THR:HA	1:A:607:SER:O	2.12	0.50
1:A:547:ASP:HB3	1:A:549:LYS:H	1.77	0.50
1:A:680:PRO:HA	1:A:688:THR:HG22	1.94	0.50
1:A:642:TRP:HE1	1:A:709:ASN:ND2	2.10	0.50
1:A:726:ALA:CA	1:A:743:MET:CE	2.90	0.50
1:A:497:ARG:HH21	1:A:524:TRP:CG	2.30	0.49
1:A:498:VAL:HG13	1:A:539:GLY:C	2.38	0.49
1:A:34:ASP:HB3	1:A:125:ARG:HD2	1.94	0.49
1:A:79:ASN:C	1:A:79:ASN:HD22	2.20	0.49
1:A:439:ILE:CD1	1:A:439:ILE:H	2.25	0.49
1:A:38:THR:HG21	1:A:85:LEU:HD22	1.94	0.49
1:A:367:ASN:OD1	1:A:369:THR:HG23	2.13	0.48
1:A:442:LEU:HD22	1:A:452:LEU:HD11	1.96	0.48
1:A:171:THR:CG2	1:A:194:HIS:HD2	2.25	0.48
1:A:439:ILE:HD12	1:A:743:MET:O	2.14	0.48
1:A:494:VAL:HA	1:A:517:THR:O	2.14	0.47
1:A:594:ASN:HD22	1:A:610:GLY:CA	2.27	0.47
1:A:738:ARG:HD3	1:A:752:GLN:OE1	2.14	0.47
1:A:324:ASP:C	1:A:326:ASN:H	2.21	0.47
1:A:539:GLY:O	1:A:553:TRP:HH2	1.95	0.47
1:A:439:ILE:HG21	1:A:748:ILE:HG13	1.97	0.47
1:A:672:PRO:HB3	1:A:691:LEU:HD11	1.97	0.47
1:A:594:ASN:HD22	1:A:610:GLY:HA3	1.79	0.47
1:A:645:VAL:HG12	1:A:648:VAL:CG1	2.45	0.47
1:A:149:GLY:HA2	1:A:176:ILE:CD1	2.45	0.46
1:A:498:VAL:HG13	1:A:539:GLY:O	2.15	0.46
1:A:597:TYR:HD2	1:A:606:VAL:HG22	1.79	0.46
1:A:325:ALA:O	1:A:326:ASN:C	2.58	0.46
1:A:488:ALA:HB3	1:A:491:ALA:O	2.15	0.45
1:A:684:ASN:C	1:A:684:ASN:ND2	2.75	0.45
1:A:645:VAL:CG1	1:A:648:VAL:HG13	2.46	0.45
1:A:431:THR:O	1:A:432:GLU:C	2.60	0.45
1:A:579:LEU:HD21	1:A:598:VAL:HG11	1.98	0.45
1:A:38:THR:HG22	1:A:70:THR:CB	2.47	0.44
1:A:439:ILE:CD1	1:A:439:ILE:N	2.81	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:274:ASP:OD1	1:A:276:THR:HB	2.18	0.44
1:A:439:ILE:O	1:A:439:ILE:HD13	2.18	0.43
1:A:439:ILE:N	1:A:439:ILE:HD13	2.33	0.43
1:A:226:ILE:HG13	1:A:267:TYR:HE2	1.82	0.43
1:A:439:ILE:HD11	1:A:742:GLY:CA	2.48	0.43
1:A:446:PRO:HD3	1:A:487:PHE:CE2	2.53	0.43
1:A:439:ILE:HD13	1:A:439:ILE:C	2.43	0.43
1:A:284:THR:HA	1:A:285:PRO:HD3	1.92	0.43
1:A:497:ARG:HG3	1:A:524:TRP:CH2	2.54	0.43
1:A:179:VAL:HG22	1:A:191:VAL:HG22	2.01	0.43
1:A:37:GLY:HA2	1:A:68:LEU:O	2.18	0.42
1:A:65:ALA:HB2	1:A:95:PRO:HD2	2.00	0.42
1:A:188:THR:HA	1:A:202:THR:O	2.20	0.42
1:A:67:LEU:HD21	1:A:96:TRP:CH2	2.55	0.42
1:A:494:VAL:HG22	1:A:518:THR:HG22	2.02	0.42
1:A:553:TRP:HB2	1:A:573:TRP:CZ3	2.55	0.42
1:A:234:ILE:H	1:A:234:ILE:HG13	1.65	0.41
1:A:308:ASN:HB2	1:A:323:LEU:HG	2.02	0.41
1:A:498:VAL:HG22	1:A:553:TRP:CH2	2.55	0.41
1:A:553:TRP:CD1	1:A:573:TRP:CZ2	3.04	0.41
1:A:723:TYR:CZ	1:A:749:ILE:HG12	2.55	0.41
1:A:32:ARG:NH1	1:A:125:ARG:O	2.53	0.41
1:A:38:THR:CG2	1:A:70:THR:OG1	2.67	0.41
1:A:741:ILE:HD12	1:A:743:MET:HE1	2.02	0.41
1:A:177:ILE:H	1:A:177:ILE:HG13	1.43	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	751/763 (98%)	707 (94%)	40 (5%)	4 (0%)	24	43

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	62	GLY
1	A	458	ASP
1	A	11	SER
1	A	354	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	586/596 (98%)	524 (89%)	62 (11%)	6 14

All (62) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	LEU
1	A	38	THR
1	A	63	ASN
1	A	67	LEU
1	A	68	LEU
1	A	71	GLU
1	A	131	LEU
1	A	171	THR
1	A	177	ILE
1	A	184	LYS
1	A	191	VAL
1	A	204	ASP
1	A	226	ILE
1	A	227	VAL
1	A	232	THR
1	A	234	ILE
1	A	276	THR
1	A	283	ILE
1	A	288	GLU
1	A	302	VAL
1	A	308	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	319	VAL
1	A	323	LEU
1	A	326	ASN
1	A	339	ASN
1	A	348	THR
1	A	369	THR
1	A	379	ARG
1	A	383	ILE
1	A	394	LEU
1	A	402	LEU
1	A	439	ILE
1	A	462	MET
1	A	469	VAL
1	A	482	THR
1	A	485	LEU
1	A	498	VAL
1	A	525	THR
1	A	534	LEU
1	A	542	ILE
1	A	549	LYS
1	A	553	TRP
1	A	577	SER
1	A	583	THR
1	A	595	THR
1	A	628	THR
1	A	648	VAL
1	A	666	SER
1	A	670	LEU
1	A	677	ILE
1	A	684	ASN
1	A	686	THR
1	A	688	THR
1	A	703	LEU
1	A	709	ASN
1	A	714	THR
1	A	728	VAL
1	A	738	ARG
1	A	739	VAL
1	A	743	MET
1	A	749	ILE
1	A	754	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10)

such sidechains are listed below:

Mol	Chain	Res	Type
1	A	79	ASN
1	A	148	GLN
1	A	194	HIS
1	A	218	GLN
1	A	326	ASN
1	A	594	ASN
1	A	635	ASN
1	A	658	ASN
1	A	709	ASN
1	A	744	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	753/763 (98%)	0.30	26 (3%)	47 42	22, 31, 43, 61	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	755	GLY	6.7
1	A	625	TRP	5.4
1	A	623	CYS	5.0
1	A	684	ASN	4.6
1	A	754	LEU	4.4
1	A	353	ALA	4.4
1	A	624	CYS	4.1
1	A	290	ALA	3.4
1	A	289	GLY	3.4
1	A	439	ILE	3.2
1	A	753	ALA	3.0
1	A	62	GLY	3.0
1	A	327	GLY	2.9
1	A	351	ALA	2.9
1	A	547	ASP	2.5
1	A	204	ASP	2.5
1	A	683	PRO	2.5
1	A	553	TRP	2.4
1	A	508	VAL	2.3
1	A	578	GLY	2.2
1	A	536	VAL	2.2
1	A	326	ASN	2.2
1	A	686	THR	2.2
1	A	352	GLY	2.1
1	A	490	ARG	2.1
1	A	184	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.