



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 07:44 AM UTC

PDB ID : 4A0B / pdb_00004a0b
Title : Structure of hsDDB1-drDDB2 bound to a 16 bp CPD-duplex (pyrimidine at D-1 position) at 3.8 Å resolution (CPD 4)
Authors : Scrima, A.; Fischer, E.S.; Iwai, S.; Gut, H.; Thoma, N.H.
Deposited on : 2011-09-08
Resolution : 3.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1065 (3.96-3.64)
Clashscore	190562	1012 (3.94-3.66)
Ramachandran outliers	187476	1048 (3.96-3.64)
Sidechain outliers	187428	1043 (3.96-3.64)
RSRZ outliers	180081	1064 (3.96-3.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1159	
1	C	1159	
2	B	382	
2	D	382	
3	G	15	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	I	15	 60% 20% 7% 13%
4	H	16	 6% 50% 38% 12%
4	J	16	 6% 81% 19%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 23938 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA DAMAGE-BINDING PROTEIN 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1090	Total	C	N	O	S	0	0	0
			8543	5423	1443	1631	46			
1	C	1095	Total	C	N	O	S	0	0	0
			8582	5447	1449	1639	47			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-18	MET	-	expression tag	UNP Q16531
A	-17	HIS	-	expression tag	UNP Q16531
A	-16	HIS	-	expression tag	UNP Q16531
A	-15	HIS	-	expression tag	UNP Q16531
A	-14	HIS	-	expression tag	UNP Q16531
A	-13	HIS	-	expression tag	UNP Q16531
A	-12	HIS	-	expression tag	UNP Q16531
A	-11	VAL	-	expression tag	UNP Q16531
A	-10	ASP	-	expression tag	UNP Q16531
A	-9	GLU	-	expression tag	UNP Q16531
A	-8	ASN	-	expression tag	UNP Q16531
A	-7	LEU	-	expression tag	UNP Q16531
A	-6	TYR	-	expression tag	UNP Q16531
A	-5	PHE	-	expression tag	UNP Q16531
A	-4	GLN	-	expression tag	UNP Q16531
A	-3	GLY	-	expression tag	UNP Q16531
A	-2	GLY	-	expression tag	UNP Q16531
A	-1	GLY	-	expression tag	UNP Q16531
A	0	ARG	-	expression tag	UNP Q16531
A	224	SER	GLU	engineered mutation	UNP Q16531
C	-18	MET	-	expression tag	UNP Q16531
C	-17	HIS	-	expression tag	UNP Q16531
C	-16	HIS	-	expression tag	UNP Q16531
C	-15	HIS	-	expression tag	UNP Q16531
C	-14	HIS	-	expression tag	UNP Q16531

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-13	HIS	-	expression tag	UNP Q16531
C	-12	HIS	-	expression tag	UNP Q16531
C	-11	VAL	-	expression tag	UNP Q16531
C	-10	ASP	-	expression tag	UNP Q16531
C	-9	GLU	-	expression tag	UNP Q16531
C	-8	ASN	-	expression tag	UNP Q16531
C	-7	LEU	-	expression tag	UNP Q16531
C	-6	TYR	-	expression tag	UNP Q16531
C	-5	PHE	-	expression tag	UNP Q16531
C	-4	GLN	-	expression tag	UNP Q16531
C	-3	GLY	-	expression tag	UNP Q16531
C	-2	GLY	-	expression tag	UNP Q16531
C	-1	GLY	-	expression tag	UNP Q16531
C	0	ARG	-	expression tag	UNP Q16531
C	224	SER	GLU	engineered mutation	UNP Q16531

- Molecule 2 is a protein called DNA DAMAGE-BINDING PROTEIN 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	354	Total	C	N	O	S	0	0	0
			2839	1804	498	526	11			
2	D	355	Total	C	N	O	S	0	0	0
			2843	1806	499	527	11			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	76	MET	-	expression tag	UNP Q2YDS1
B	77	HIS	-	expression tag	UNP Q2YDS1
B	78	HIS	-	expression tag	UNP Q2YDS1
B	79	HIS	-	expression tag	UNP Q2YDS1
B	80	HIS	-	expression tag	UNP Q2YDS1
B	81	HIS	-	expression tag	UNP Q2YDS1
B	82	HIS	-	expression tag	UNP Q2YDS1
B	83	ARG	-	expression tag	UNP Q2YDS1
B	84	ARG	-	expression tag	UNP Q2YDS1
B	85	LEU	-	expression tag	UNP Q2YDS1
B	86	VAL	-	expression tag	UNP Q2YDS1
B	87	PRO	-	expression tag	UNP Q2YDS1
B	88	ARG	-	expression tag	UNP Q2YDS1
B	89	GLY	-	expression tag	UNP Q2YDS1
B	90	SER	-	expression tag	UNP Q2YDS1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	91	GLY	-	expression tag	UNP Q2YDS1
B	92	GLY	-	expression tag	UNP Q2YDS1
B	93	ARG	-	expression tag	UNP Q2YDS1
B	180	GLN	LEU	variant	UNP Q2YDS1
B	214	ARG	TRP	variant	UNP Q2YDS1
D	76	MET	-	expression tag	UNP Q2YDS1
D	77	HIS	-	expression tag	UNP Q2YDS1
D	78	HIS	-	expression tag	UNP Q2YDS1
D	79	HIS	-	expression tag	UNP Q2YDS1
D	80	HIS	-	expression tag	UNP Q2YDS1
D	81	HIS	-	expression tag	UNP Q2YDS1
D	82	HIS	-	expression tag	UNP Q2YDS1
D	83	ARG	-	expression tag	UNP Q2YDS1
D	84	ARG	-	expression tag	UNP Q2YDS1
D	85	LEU	-	expression tag	UNP Q2YDS1
D	86	VAL	-	expression tag	UNP Q2YDS1
D	87	PRO	-	expression tag	UNP Q2YDS1
D	88	ARG	-	expression tag	UNP Q2YDS1
D	89	GLY	-	expression tag	UNP Q2YDS1
D	90	SER	-	expression tag	UNP Q2YDS1
D	91	GLY	-	expression tag	UNP Q2YDS1
D	92	GLY	-	expression tag	UNP Q2YDS1
D	93	ARG	-	expression tag	UNP Q2YDS1
D	180	GLN	LEU	variant	UNP Q2YDS1
D	214	ARG	TRP	variant	UNP Q2YDS1

- Molecule 3 is a DNA chain called 5'-D(*DGP*GP*TP*GP*AP*AP*AP*(TTD)P*AP*GP*CP*AP*GP*DGP)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	G	14	Total	C	N	O	P	0	0	0
			312	149	61	88	14			
3	I	13	Total	C	N	O	P	0	0	0
			290	139	56	82	13			

- Molecule 4 is a DNA chain called 5'-D(*CP*CP*TP*GP*CP*TP*CP*CP*TP*TP*TP*CP*AP*CP*CP*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	H	14	Total	C	N	O	P	0	0	0
			274	133	44	84	13			

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	J	13	Total	C	N	O	P	0	0	0
			255	124	41	78	12			

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: DNA DAMAGE-BINDING PROTEIN 1



T1130
K1131
V1132
V1133
E1134
T1137
R1138
R1139
H1140

• Molecule 1: DNA DAMAGE-BINDING PROTEIN 1


Chain C:  65% 27% 6%



• Molecule 2: DNA DAMAGE-BINDING PROTEIN 2


Chain B:  63% 26% 7%





• Molecule 2: DNA DAMAGE-BINDING PROTEIN 2

Chain D: 65% 24% 7%




• Molecule 3: 5'-D(*DGP*GP*TP*GP*AP*AP*AP*(TTD)P*AP*GP*CP*AP*GP*DGP)-3'

Chain G: 60% 27% 7%




• Molecule 3: 5'-D(*DGP*GP*TP*GP*AP*AP*AP*(TTD)P*AP*GP*CP*AP*GP*DGP)-3'

Chain I: 60% 20% 7% 13%





• Molecule 4: 5'-D(*CP*CP*TP*GP*CP*TP*CP*CP*TP*TP*TP*CP*AP*CP*CP*C)-3'

Chain H: 6% 50% 38% 12%



• Molecule 4: 5'-D(*CP*CP*TP*GP*CP*TP*CP*CP*TP*TP*TP*CP*AP*CP*CP*C)-3'



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2 ₁	Depositor
Cell constants a, b, c, α , β , γ	113.10Å 145.90Å 224.44Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.72 – 3.80 47.72 – 3.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (47.72-3.80) 99.7 (47.72-3.80)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.34 (at 3.77Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.243 , 0.319 0.240 , 0.312	Depositor DCC
R_{free} test set	1862 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	102.6	Xtriage
Anisotropy	0.090	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 63.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	23938	wwPDB-VP
Average B, all atoms (Å ²)	126.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 19.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TTD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.48	0/8697	0.79	4/11777 (0.0%)
1	C	0.48	0/8737	0.78	4/11831 (0.0%)
2	B	0.49	0/2913	0.82	0/3957
2	D	0.48	0/2917	0.83	2/3962 (0.1%)
3	G	0.22	0/306	0.76	0/470
3	I	0.20	0/281	0.67	0/431
4	H	0.25	0/304	0.92	0/465
4	J	0.22	0/283	0.79	0/433
All	All	0.47	0/24438	0.80	10/33326 (0.0%)

There are no bond length outliers.

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	1025	GLN	N-CA-C	7.42	119.88	110.24
1	C	785	GLU	N-CA-C	6.11	119.46	110.30
2	D	190	GLY	N-CA-C	5.49	123.53	112.34
1	A	1134	GLU	N-CA-C	5.29	116.73	111.07
2	D	181	ASN	N-CA-C	5.28	117.95	111.82

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8543	0	8538	193	0
1	C	8582	0	8588	183	0
2	B	2839	0	2785	64	0
2	D	2843	0	2788	72	0
3	G	312	0	171	8	0
3	I	290	0	160	4	0
4	H	274	0	160	6	0
4	J	255	0	149	0	0
All	All	23938	0	23339	515	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 515 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:40:GLU:HG2	1:A:54:GLU:HG3	1.44	0.97
1:A:864:LYS:HG3	1:A:899:LEU:HB2	1.51	0.93
1:A:204:LYS:HG2	1:A:205:GLY:H	1.33	0.91
1:C:391:ARG:HH21	1:C:711:HIS:HD2	1.18	0.90
2:D:102:GLN:HG3	2:D:107:HIS:HB3	1.54	0.89

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1076/1159 (93%)	960 (89%)	100 (9%)	16 (2%)	8	36
1	C	1081/1159 (93%)	953 (88%)	114 (10%)	14 (1%)	9	38
2	B	352/382 (92%)	307 (87%)	36 (10%)	9 (3%)	4	27
2	D	353/382 (92%)	318 (90%)	30 (8%)	5 (1%)	9	37

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	2862/3082 (93%)	2538 (89%)	280 (10%)	44 (2%)	8	36

5 of 44 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	118	ILE
2	B	183	THR
2	B	194	ALA
1	C	707	ILE
1	C	855	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	957/1015 (94%)	852 (89%)	105 (11%)	6	24
1	C	962/1015 (95%)	862 (90%)	100 (10%)	7	26
2	B	313/335 (93%)	276 (88%)	37 (12%)	5	22
2	D	313/335 (93%)	285 (91%)	28 (9%)	9	32
All	All	2545/2700 (94%)	2275 (89%)	270 (11%)	6	26

5 of 270 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	1041	THR
1	C	1108	VAL
2	D	365	ILE
1	A	1065	VAL
1	A	1043	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 86 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	524	GLN
1	C	908	ASN
1	C	550	ASN
1	C	731	GLN
1	C	1111	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	TTD	I	9	3	42,45,46	1.60	8 (19%)	61,74,77	2.74	16 (26%)
3	TTD	G	9	3	42,45,46	1.68	8 (19%)	61,74,77	2.45	17 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TTD	I	9	3	-	13/22/109/110	0/5/6/6
3	TTD	G	9	3	-	19/22/109/110	0/5/6/6

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	G	9	TTD	C1'-N1	4.63	1.51	1.45
3	G	9	TTD	C1R-N1T	4.56	1.51	1.45
3	I	9	TTD	C1R-N1T	4.14	1.51	1.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	I	9	TTD	C1'-N1	4.13	1.51	1.45
3	G	9	TTD	C2-N1	3.82	1.43	1.36

The worst 5 of 33 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	I	9	TTD	O4R-C1R-N1T	13.54	124.69	108.65
3	G	9	TTD	O4R-C1R-N1T	8.05	118.19	108.65
3	I	9	TTD	C5T-C4T-N3T	6.90	121.88	116.09
3	I	9	TTD	C5-C4-N3	6.67	121.69	116.09
3	G	9	TTD	N3-C2-N1	6.47	123.56	116.78

There are no chirality outliers.

5 of 32 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	G	9	TTD	C3R-O3R-PB-O4P
3	G	9	TTD	C3R-O3R-PB-O5R
3	G	9	TTD	O4'-C1'-N1-C2
3	G	9	TTD	C5R-O5R-PB-O3R
3	G	9	TTD	C5R-O5R-PB-O5P

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	I	9	TTD	1	0
3	G	9	TTD	5	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1090/1159 (94%)	-0.42	0 100 100	68, 109, 228, 265	0
1	C	1095/1159 (94%)	-0.45	1 (0%) 92 87	70, 106, 215, 247	0
2	B	354/382 (92%)	-0.56	0 100 100	77, 98, 132, 144	0
2	D	355/382 (92%)	-0.49	0 100 100	74, 103, 147, 175	0
3	G	13/15 (86%)	-0.01	0 100 100	128, 164, 217, 229	0
3	I	12/15 (80%)	0.34	0 100 100	167, 206, 232, 286	0
4	H	14/16 (87%)	0.34	1 (7%) 22 19	131, 154, 223, 232	0
4	J	13/16 (81%)	0.79	1 (7%) 19 18	196, 230, 337, 374	0
All	All	2946/3144 (93%)	-0.44	3 (0%) 92 87	68, 106, 219, 374	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	H	2	DC	2.4
1	C	406	GLY	2.2
4	J	2	DC	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	TTD	I	9	40/41	0.86	0.09	164,193,226,236	0
3	TTD	G	9	40/41	0.93	0.08	128,141,161,167	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.