



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:41 PM UTC

PDB ID : 7A03 / pdb_00007a03
Title : The Structure of CHT
Authors : Fernandes, G.; Machado, E.; Pereira, M.; Brear, P.; Lemos, E.
Deposited on : 2020-08-06
Resolution : 1.39 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.40-1.40)
Clashscore	190562	2660 (1.40-1.40)
Ramachandran outliers	187476	2611 (1.40-1.40)
Sidechain outliers	187428	2610 (1.40-1.40)
RSRZ outliers	180081	2561 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	500	16% 93% 6%
1	B	500	19% 93% 6%

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 8775 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called M32 carboxypeptidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	493	Total	C	N	O	S	0	6	0
			4062	2589	696	765	12			
1	B	494	Total	C	N	O	S	0	7	0
			4077	2597	698	771	11			

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 4 is CITRIC ACID (CCD ID: CIT) (formula: $C_6H_8O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			5	2	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	307	Total	O	0	0
			307	307		
5	B	277	Total	O	0	0
			277	277		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: M32 carboxypeptidase



- Molecule 1: M32 carboxypeptidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	93.09Å 110.27Å 113.87Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	71.13 – 1.39 71.13 – 1.39	Depositor EDS
% Data completeness (in resolution range)	96.2 (71.13-1.39) 97.4 (71.13-1.39)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.55 (at 1.39Å)	Xtriage
Refinement program	PHENIX 1.16_3549	Depositor
R, R_{free}	0.245 , 0.266 0.246 , 0.266	Depositor DCC
R_{free} test set	11857 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	14.2	Xtriage
Anisotropy	0.327	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 29.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.013 for -h,l,k	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	8775	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CIT, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.30	0/4172	0.53	0/5663
1	B	0.30	0/4188	0.51	0/5688
All	All	0.30	0/8360	0.52	0/11351

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4062	0	3887	20	0
1	B	4077	0	3893	22	0
2	A	18	0	24	0	0
2	B	18	0	24	3	0
3	A	7	0	10	1	0
4	A	4	0	0	0	0
4	B	5	0	0	0	0
5	A	307	0	0	0	0
5	B	277	0	0	1	0
All	All	8775	0	7838	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 3.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:10:GLN:HG2	1:A:74:LEU:HD21	1.72	0.70
1:A:199:ARG:HG3	1:A:340:ALA:HB1	1.78	0.64
1:B:301:ARG:NH1	2:B:603:GOL:H11	2.14	0.62
1:B:301:ARG:CZ	2:B:603:GOL:H11	2.32	0.59
1:A:10:GLN:H	1:A:10:GLN:CD	2.13	0.56
1:A:155[A]:HIS:CE1	1:A:374:ASP:HB3	2.43	0.54
1:B:143:LYS:O	1:B:147:THR:HG23	2.08	0.54
1:B:127:ASN:HD21	1:B:402:THR:H	1.55	0.53
1:A:97:LYS:HE2	1:A:101:TYR:HE1	1.74	0.52
1:A:137:ALA:HB2	1:A:379:THR:HG21	1.91	0.52
1:B:473[A]:ASN:O	1:B:477:GLN:HG3	2.10	0.52
1:A:125:LYS:N	1:A:125:LYS:HD3	2.27	0.50
1:A:140:VAL:O	1:A:144:ARG:HG3	2.11	0.50
1:B:62:PHE:HD2	1:B:98:ASN:HD22	1.59	0.49
1:B:127:ASN:ND2	1:B:402:THR:H	2.10	0.49
1:A:252:ARG:CZ	1:A:260:ASN:HD21	2.26	0.48
1:B:95:TYR:CZ	1:B:99:LYS:HE3	2.47	0.48
1:B:175:THR:O	1:B:179:GLU:HG3	2.12	0.48
1:A:137:ALA:O	1:A:141:GLU:HG3	2.14	0.48
1:B:10:GLN:HG3	1:B:73:GLU:CD	2.39	0.48
1:B:305:ASN:HD21	2:B:603:GOL:H32	1.80	0.47
1:B:371:GLY:HA3	1:B:377:ILE:HD12	1.97	0.46
1:A:121:ILE:HG23	1:A:125:LYS:HZ3	1.82	0.45
1:B:95:TYR:OH	1:B:99:LYS:HE3	2.17	0.45
1:B:473[B]:ASN:O	1:B:477:GLN:HG3	2.17	0.45
1:A:143:LYS:O	1:A:147:THR:HG23	2.16	0.44
1:B:10:GLN:HG3	1:B:73:GLU:OE2	2.17	0.44
1:B:164:TYR:CE1	1:B:418:TYR:HD2	2.36	0.44
1:B:205:HIS:HD2	5:B:950:HOH:O	2.01	0.44
1:A:120:TRP:CZ2	1:A:124:ARG:HD2	2.53	0.43
1:A:466:PHE:HD1	3:A:604:PEG:H21	1.82	0.43
1:B:97:LYS:HA	1:B:97:LYS:HD3	1.75	0.43
1:A:498:LEU:HD23	1:A:498:LEU:HA	1.79	0.42
1:A:10:GLN:HG2	1:A:74:LEU:CD2	2.47	0.42
1:B:59:HIS:NE2	1:B:98:ASN:OD1	2.52	0.42
1:A:10:GLN:CD	1:A:10:GLN:N	2.77	0.41
1:B:111:GLU:HG3	1:B:115:GLN:OE1	2.21	0.41
1:A:220:ARG:HD3	1:A:226[A]:MET:HE3	2.03	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:232:ASP:O	1:A:250:THR:HA	2.21	0.40
1:B:473[B]:ASN:OD1	1:B:483:PRO:HB3	2.21	0.40
1:A:339:LYS:HA	1:A:339:LYS:HD3	1.88	0.40
1:B:93:GLU:O	1:B:97:LYS:HG2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	497/500 (99%)	484 (97%)	10 (2%)	3 (1%)	21	5
1	B	499/500 (100%)	488 (98%)	10 (2%)	1 (0%)	43	19
All	All	996/1000 (100%)	972 (98%)	20 (2%)	4 (0%)	36	10

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	198[A]	ARG
1	A	198[B]	ARG
1	A	243	ASN
1	B	243	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	432/432 (100%)	432 (100%)	0	100	100
1	B	435/432 (101%)	435 (100%)	0	100	100
All	All	867/864 (100%)	867 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	400	ASN
1	B	35	GLN
1	B	127	ASN
1	B	130	GLN
1	B	305	ASN
1	B	306	ASN
1	B	387	ASN
1	B	482	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	B	601	-	5,5,5	0.91	0	5,5,5	1.04	0
2	GOL	B	603	-	5,5,5	0.97	0	5,5,5	1.24	0
4	CIT	B	604	-	4,4,12	1.13	0	4,4,17	2.78	1 (25%)
2	GOL	B	602	-	5,5,5	0.74	0	5,5,5	1.13	0
3	PEG	A	604	-	6,6,6	0.50	0	5,5,5	0.24	0
4	CIT	A	605	-	3,3,12	0.86	0	3,3,17	1.02	0
2	GOL	A	602	-	5,5,5	0.95	0	5,5,5	1.02	0
2	GOL	A	603	-	5,5,5	0.88	0	5,5,5	1.15	0
2	GOL	A	601	-	5,5,5	0.90	0	5,5,5	1.01	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	601	-	-	1/4/4/4	-
2	GOL	B	603	-	-	2/4/4/4	-
4	CIT	B	604	-	-	2/2/2/16	-
2	GOL	B	602	-	-	0/4/4/4	-
3	PEG	A	604	-	-	2/4/4/4	-
2	GOL	A	602	-	-	0/4/4/4	-
2	GOL	A	603	-	-	1/4/4/4	-
2	GOL	A	601	-	-	0/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	604	CIT	O6-C6-C3	4.90	120.80	111.90

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	603	GOL	C1-C2-C3-O3
4	B	604	CIT	O7-C3-C6-O5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	B	604	CIT	O7-C3-C6-O6
2	A	603	GOL	O1-C1-C2-O2
3	A	604	PEG	C4-C3-O2-C2
2	B	601	GOL	C1-C2-C3-O3
2	B	603	GOL	O2-C2-C3-O3
3	A	604	PEG	C1-C2-O2-C3

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	603	GOL	3	0
3	A	604	PEG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	493/500 (98%)	1.06	80 (16%) 4 4	6, 19, 38, 50	6 (1%)
1	B	494/500 (98%)	1.09	95 (19%) 3 3	7, 19, 39, 60	7 (1%)
All	All	987/1000 (98%)	1.07	175 (17%) 4 4	6, 19, 38, 60	13 (1%)

All (175) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	8	ALA	7.2
1	A	500	GLY	7.0
1	B	76[A]	HIS	7.0
1	A	441	PRO	5.9
1	A	499	ARG	5.8
1	A	197	ALA	5.8
1	A	117	TYR	5.6
1	B	98	ASN	5.3
1	B	8	ALA	5.2
1	B	118	HIS	5.1
1	B	441	PRO	4.9
1	B	77	HIS	4.9
1	A	126	ALA	4.9
1	B	197	ALA	4.7
1	B	99	LYS	4.6
1	B	149	VAL	4.5
1	A	79	GLU	4.5
1	A	199	ARG	4.5
1	B	103	ALA	4.5
1	B	198	ARG	4.5
1	A	76	HIS	4.3
1	A	196	PRO	4.3
1	B	101	TYR	4.3
1	A	125	LYS	4.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	6	SER	4.2
1	B	125	LYS	4.2
1	B	117	TYR	4.1
1	A	10	GLN	4.1
1	B	10	GLN	4.1
1	B	122	LYS	4.1
1	A	97	LYS	4.1
1	A	134	PRO	4.1
1	A	121	ILE	4.0
1	B	7	THR	4.0
1	B	194	GLN	4.0
1	A	131	VAL	3.9
1	B	131	VAL	3.9
1	A	498	LEU	3.9
1	B	97	LYS	3.8
1	A	195	THR	3.8
1	B	123	ALA	3.8
1	B	121	ILE	3.8
1	B	148	THR	3.7
1	B	498	LEU	3.7
1	B	78	PRO	3.6
1	A	380	LYS	3.6
1	A	123	ALA	3.6
1	B	105	LEU	3.6
1	B	163	GLU	3.6
1	A	148	THR	3.6
1	A	114	ASN	3.6
1	B	126	ALA	3.5
1	A	377	ILE	3.5
1	B	195	THR	3.5
1	B	167	GLY	3.5
1	B	409	ILE	3.5
1	A	200	ASP	3.5
1	A	98	ASN	3.4
1	A	132	PHE	3.4
1	B	109	ILE	3.4
1	A	120	TRP	3.4
1	B	107	ALA	3.4
1	B	152	TYR	3.3
1	B	114	ASN	3.3
1	B	79	GLU	3.3
1	B	164	TYR	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	122	LYS	3.2
1	B	193	LYS	3.2
1	B	100	LYS	3.2
1	B	380	LYS	3.2
1	A	101	TYR	3.1
1	A	379	THR	3.1
1	A	393	TYR	3.1
1	A	145	LYS	3.1
1	B	9	GLU	3.1
1	A	116	ALA	3.1
1	B	199	ARG	3.1
1	A	378	SER	3.0
1	A	118	HIS	3.0
1	A	149	VAL	3.0
1	A	135	ALA	3.0
1	A	252	ARG	2.9
1	A	119	ALA	2.9
1	A	440	VAL	2.9
1	B	154	ASP	2.9
1	B	418	TYR	2.9
1	A	77	HIS	2.9
1	A	9	GLU	2.8
1	B	360	PHE	2.8
1	A	187	LEU	2.8
1	A	198[A]	ARG	2.8
1	B	375	GLY	2.8
1	B	111	GLU	2.8
1	A	402	THR	2.8
1	A	381	ASP	2.7
1	B	66	GLU	2.7
1	B	153	GLU	2.7
1	B	196	PRO	2.7
1	A	375	GLY	2.7
1	B	473[A]	ASN	2.7
1	B	490	LEU	2.7
1	A	442	ASP	2.7
1	A	181	LYS	2.7
1	B	145	LYS	2.7
1	B	373	ILE	2.6
1	B	102	PRO	2.6
1	A	472	SER	2.6
1	B	171	ASP	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	120	TRP	2.6
1	A	473	ASN	2.6
1	B	327	GLU	2.6
1	B	127	ASN	2.6
1	B	150	LEU	2.5
1	A	41	GLU	2.5
1	B	176	ILE	2.5
1	A	100	LYS	2.5
1	B	106	VAL	2.5
1	B	377	ILE	2.5
1	A	163	GLU	2.5
1	B	132	PHE	2.4
1	A	65	PRO	2.4
1	A	80	LEU	2.4
1	B	129	TYR	2.4
1	B	403	GLN	2.4
1	A	75	HIS	2.4
1	B	477	GLN	2.4
1	B	393	TYR	2.4
1	A	193	LYS	2.4
1	B	119	ALA	2.4
1	B	497	PHE	2.4
1	B	80	LEU	2.3
1	B	170	VAL	2.3
1	B	124	ARG	2.3
1	B	168	ALA	2.3
1	A	180	VAL	2.3
1	A	382	LEU	2.3
1	B	440	VAL	2.3
1	B	116	ALA	2.2
1	A	111	GLU	2.2
1	A	385	THR	2.2
1	B	226	MET	2.2
1	A	345	GLN	2.2
1	A	129	TYR	2.2
1	B	140	VAL	2.2
1	B	135	ALA	2.2
1	B	402	THR	2.2
1	B	151	GLY	2.2
1	B	172	MET	2.2
1	B	110	SER	2.2
1	A	78	PRO	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	208	ARG	2.1
1	A	406	LEU	2.1
1	B	75	HIS	2.1
1	B	499	ARG	2.1
1	A	128	ASP	2.1
1	B	442	ASP	2.1
1	B	471	THR	2.1
1	A	153	GLU	2.1
1	B	235	GLU	2.1
1	A	127	ASN	2.1
1	B	191	ILE	2.1
1	A	141	GLU	2.1
1	A	102	PRO	2.1
1	B	481	GLY	2.1
1	A	383	ASN	2.1
1	A	400	ASN	2.1
1	B	324	LEU	2.1
1	A	115	GLN	2.1
1	A	374	ASP	2.1
1	A	409	ILE	2.1
1	A	110	SER	2.0
1	B	166	LYS	2.0
1	B	469	PHE	2.0
1	B	352	GLU	2.0
1	A	203	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GOL	B	603	6/6	0.69	0.23	26,38,42,44	0
2	GOL	A	603	6/6	0.75	0.18	37,41,46,50	0
4	CIT	B	604	5/13	0.81	0.16	33,34,37,38	0
3	PEG	A	604	7/7	0.83	0.15	33,34,37,39	0
4	CIT	A	605	4/13	0.87	0.16	24,30,30,35	0
2	GOL	A	601	6/6	0.89	0.11	27,29,33,36	0
2	GOL	B	602	6/6	0.93	0.10	20,23,30,32	0
2	GOL	B	601	6/6	0.97	0.07	14,20,21,23	0
2	GOL	A	602	6/6	0.97	0.06	17,19,20,21	0

6.5 Other polymers [i](#)

There are no such residues in this entry.