



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 05:45 PM UTC

PDB ID : 8A1P / pdb_00008a1p
Title : HIV-1 Integrase Catalytic Core Domain and C-Terminal Domain in Complex with Allosteric Integrase Inhibitor BI-D
Authors : Singer, M.R.; Pye, V.E.; Cook, N.J.; Cherepanov, P.
Deposited on : 2022-06-01
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	233	3% 21% 78%
1	B	233	3% 54% 8% 39%
1	C	233	4% 18% 79%
1	D	233	58% 39%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 3372 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Integrase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	52	Total	C	N	O		0	0	0
			427	272	81	74				
1	B	143	Total	C	N	O	S	0	3	0
			1132	725	198	205	4			
1	C	50	Total	C	N	O		0	0	0
			410	263	76	71				
1	D	141	Total	C	N	O	S	0	1	0
			1102	704	192	201	5			

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	219	SER	-	expression tag	UNP P12497
A	243	GLU	TRP	engineered mutation	UNP P12497
A	424	LYS	PHE	engineered mutation	UNP P12497
B	-20	SER	-	expression tag	UNP P12497
B	4	GLU	TRP	engineered mutation	UNP P12497
B	185	LYS	PHE	engineered mutation	UNP P12497
C	219	SER	-	expression tag	UNP P12497
C	243	GLU	TRP	engineered mutation	UNP P12497
C	424	LYS	PHE	engineered mutation	UNP P12497
D	-20	SER	-	expression tag	UNP P12497
D	4	GLU	TRP	engineered mutation	UNP P12497
D	185	LYS	PHE	engineered mutation	UNP P12497

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			7	4	3		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	D	1	Total	C	O	0	0
			7	4	3		
3	D	1	Total	C	O	0	0
			7	4	3		
3	D	1	Total	C	O	0	0
			7	4	3		
3	D	1	Total	C	O	0	0
			7	4	3		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		
4	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is (2S)-tert-butoxy[4-(3,4-dihydro-2H-chromen-6-yl)-2-methylquinolin-3-yl]ethanoic acid (CCD ID: LF0) (formula: $C_{25}H_{27}NO_4$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	N	O	0	0
			30	25	1	4		
5	D	1	Total	C	N	O	0	0
			30	25	1	4		



- Molecule 6 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Mg	0	0
			1	1		
6	D	1	Total	Mg	0	0
			1	1		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	15	Total	O	0	0
			15	15		
7	B	79	Total	O	0	0
			79	79		
7	C	9	Total	O	0	0
			9	9		
7	D	77	Total	O	0	0
			77	77		


- Molecule 1: Integrase



VAL	LYS	ALA	ALA	ALA	CYS	TRP	TRP	ALA	ALA	GLY	ILE	ILE	LYS	GLN	GLU	PHE	GLY	ILE	PRO	TYR	ASN	PRO	GLN	SER	SER	GLN	GLY	GLY	VAL	ILE	ILE	GLU	GLU	LEU	LYS	LYS	ILE	ILE	GLN	GLY	GLN	VAL	ARG	ASP	GLN	ALA	ALA	GLU	LEU	LEU	LYS	THR	VAL	PHE	ILE	ILE	HIS	LYS	ASN	LYS
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

LYS ARG LYS GLY GLY ILE GLY GLY TYR SER ALA GLY GLU ARG ILE VAL ASP ILE ILE ALA THR ASP ILE GLN THR LYS GLU

- Molecule 1: Integrase

Chain D:  58% . 39%

SER	ILE	GLN	ASN	PHE	ARG	VAL	TYR	TYR	ARG	ASP	SER	ARG	ASP	PRO	VAL	VAL	TRP	LYS	GLY	GLY	PRO	ALA	ALA	LEU	LEU	GLU	GLY	GLY	GLY	ALA	VAL	VAL	ILE	ILE	GLN	ASP	ASN	SER	SER	ASP	ASP	ILE	ILE	LYS	VAL	VAL	PRO	ARG	ARG	LYS	LYS	ALA	ALA	ILE	ILE	ARG	ASP	TYR	GLY	GLN	GLN	MET	ALA	ALA	GLY	ASP
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

ASP	CYS	VAL	ALA	SER	ARG	GLN	ASP	GLU	MET	HIS	GLY	GLN	VAL	ASP	ES7	G140	ILE	PRO	TYR	ASN	PRO	GLN	SER	GLN	G149	K173	R187	K188	GLY	GLY	ILE	GLY	G193	I194	S195	V201	A205	I208	GLN	THR	LYS	GLU
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	-----	-----	-----	-----	------	------	------	------	------	------	-----	-----	-----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	53.55Å 65.21Å 69.67Å 90.00° 100.17° 90.00°	Depositor
Resolution (Å)	52.71 – 1.80 52.71 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.3 (52.71-1.80) 99.4 (52.71-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.04 (at 1.80Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.181 , 0.209 0.181 , 0.209	Depositor DCC
R_{free} test set	2177 reflections (3.97%)	wwPDB-VP
Wilson B-factor (Å ²)	43.2	Xtriage
Anisotropy	0.364	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 48.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3372	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, EDO, PEG, GOL, LF0

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.17	0/435	0.39	0/583
1	B	0.16	0/1160	0.33	0/1567
1	C	0.11	0/417	0.32	0/558
1	D	0.17	0/1123	0.35	0/1517
All	All	0.16	0/3135	0.34	0/4225

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	427	0	444	3	0
1	B	1132	0	1148	16	0
1	C	410	0	425	4	0
1	D	1102	0	1109	6	0
2	A	4	0	6	0	0
2	B	8	0	12	1	0
3	B	7	0	10	1	0
3	D	28	0	40	1	0
4	B	6	0	8	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	6	0	8	0	0
5	B	30	0	26	0	0
5	D	30	0	26	1	0
6	B	1	0	0	0	0
6	D	1	0	0	0	0
7	A	15	0	0	0	0
7	B	79	0	0	1	0
7	C	9	0	0	0	0
7	D	77	0	0	1	0
All	All	3372	0	3262	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:77[A]:VAL:HG22	1:B:84:ILE:HG22	1.72	0.71
1:B:95:GLN:NE2	7:B:703:HOH:O	2.27	0.67
1:B:95:GLN:HG2	5:D:304:LF0:H21	1.82	0.61
1:B:114[B]:HIS:NE2	1:B:140:GLY:O	2.32	0.61
1:C:245:GLY:O	1:C:262:ARG:NH1	2.32	0.60
1:A:224:ARG:HG3	1:B:131:TRP:CE2	2.37	0.59
1:B:114[A]:HIS:NE2	1:B:138:GLU:OE1	2.37	0.57
1:D:195:SER:OG	3:D:306:PEG:H31	2.05	0.57
1:A:268:ILE:HD13	1:B:128:ALA:HB1	1.87	0.55
3:B:601:PEG:H31	1:D:173:LYS:HZ2	1.72	0.54
1:C:249:VAL:HG11	1:C:267:ILE:HD11	1.91	0.52
1:C:235:TRP:CH2	1:C:266:LYS:HE2	2.49	0.48
1:B:205:ALA:HB2	1:D:201:VAL:HG12	1.96	0.47
1:B:211:LYS:NZ	2:B:602:EDO:O1	2.49	0.46
1:B:188:LYS:HG3	1:B:193:GLY:HA2	1.99	0.45
1:B:85:GLU:OE1	1:B:107:ARG:NH1	2.52	0.43
1:B:201:VAL:HG12	1:D:205:ALA:HB2	2.01	0.42
1:D:187:ARG:NH1	1:D:193:GLY:O	2.53	0.42
1:A:270:ASP:HB3	1:B:131:TRP:CE2	2.56	0.41
1:D:87:GLU:OE1	7:D:401:HOH:O	2.22	0.41
1:B:123:SER:O	1:B:127:LYS:HG3	2.20	0.41
1:C:273:LYS:HB2	1:C:273:LYS:HE2	1.87	0.41
1:B:62:GLN:HG2	1:B:114[A]:HIS:HB2	2.04	0.40
1:B:62:GLN:HG2	1:B:114[B]:HIS:HB3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	50/233 (22%)	49 (98%)	1 (2%)	0	100	100
1	B	140/233 (60%)	140 (100%)	0	0	100	100
1	C	46/233 (20%)	43 (94%)	3 (6%)	0	100	100
1	D	136/233 (58%)	136 (100%)	0	0	100	100
All	All	372/932 (40%)	368 (99%)	4 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	45/193 (23%)	44 (98%)	1 (2%)	45	34
1	B	120/193 (62%)	120 (100%)	0	100	100
1	C	43/193 (22%)	43 (100%)	0	100	100
1	D	116/193 (60%)	116 (100%)	0	100	100
All	All	324/772 (42%)	323 (100%)	1 (0%)	86	86

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	273	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	95	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 14 ligands modelled in this entry, 2 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	D	301	-	5,5,5	0.94	0	5,5,5	1.06	0
3	PEG	D	302	-	6,6,6	0.12	0	5,5,5	0.06	0
5	LF0	D	304	-	31,33,33	1.78	8 (25%)	41,49,49	1.31	3 (7%)
3	PEG	D	305	-	6,6,6	0.11	0	5,5,5	0.10	0
3	PEG	D	306	-	6,6,6	0.10	0	5,5,5	0.12	0
2	EDO	A	901	-	3,3,3	0.49	0	2,2,2	0.30	0
3	PEG	B	601	-	6,6,6	0.15	0	5,5,5	0.08	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	LF0	B	604	-	31,33,33	1.65	7 (22%)	41,49,49	1.16	2 (4%)
3	PEG	D	303	-	6,6,6	0.16	0	5,5,5	0.06	0
2	EDO	B	602	-	3,3,3	0.44	0	2,2,2	0.39	0
4	GOL	B	603	-	5,5,5	0.92	0	5,5,5	1.06	0
2	EDO	B	605	-	3,3,3	0.45	0	2,2,2	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	D	301	-	-	0/4/4/4	-
3	PEG	D	302	-	-	3/4/4/4	-
5	LF0	D	304	-	-	2/17/24/24	0/4/4/4
3	PEG	D	305	-	-	2/4/4/4	-
3	PEG	D	306	-	-	2/4/4/4	-
2	EDO	A	901	-	-	1/1/1/1	-
3	PEG	B	601	-	-	3/4/4/4	-
5	LF0	B	604	-	-	2/17/24/24	0/4/4/4
3	PEG	D	303	-	-	4/4/4/4	-
2	EDO	B	602	-	-	0/1/1/1	-
4	GOL	B	603	-	-	1/4/4/4	-
2	EDO	B	605	-	-	0/1/1/1	-

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	D	304	LF0	O27-C10	5.58	1.43	1.37
5	B	604	LF0	O27-C10	4.76	1.43	1.37
5	D	304	LF0	C1-C7	3.32	1.53	1.49
5	B	604	LF0	C6-C5	-3.26	1.37	1.42
5	D	304	LF0	C21-C11	3.23	1.56	1.51
5	B	604	LF0	C21-C11	3.20	1.56	1.51
5	D	304	LF0	C6-C5	-3.13	1.37	1.42
5	B	604	LF0	C1-C7	2.69	1.53	1.49
5	D	304	LF0	C17-C3	2.17	1.53	1.50
5	D	304	LF0	O22-C23	-2.15	1.40	1.45
5	B	604	LF0	C1-C2	2.12	1.41	1.38
5	B	604	LF0	C17-C3	2.06	1.53	1.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	604	LF0	O27-C4	-2.04	1.39	1.43
5	D	304	LF0	O27-C4	-2.03	1.39	1.43
5	D	304	LF0	C1-C2	2.03	1.41	1.38

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	D	304	LF0	C3-N1-C5	4.33	122.40	118.42
5	B	604	LF0	C3-N1-C5	3.54	121.67	118.42
5	D	304	LF0	C7-C1-C2	-3.09	118.22	121.56
5	D	304	LF0	C8-C7-C1	-2.54	117.01	120.75
5	B	604	LF0	C8-C7-C1	-2.18	117.54	120.75

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	604	LF0	C2-C18-C19-O20
5	B	604	LF0	C2-C18-C19-O21
5	D	304	LF0	C2-C18-C19-O20
5	D	304	LF0	C2-C18-C19-O21
3	D	306	PEG	O2-C3-C4-O4
3	D	302	PEG	O1-C1-C2-O2
3	D	303	PEG	O2-C3-C4-O4
3	D	305	PEG	O1-C1-C2-O2
3	D	306	PEG	O1-C1-C2-O2
3	B	601	PEG	O1-C1-C2-O2
3	B	601	PEG	O2-C3-C4-O4
2	A	901	EDO	O1-C1-C2-O2
3	D	303	PEG	C4-C3-O2-C2
3	D	305	PEG	C1-C2-O2-C3
3	D	303	PEG	O1-C1-C2-O2
3	D	302	PEG	C1-C2-O2-C3
3	B	601	PEG	C1-C2-O2-C3
3	D	303	PEG	C1-C2-O2-C3
4	B	603	GOL	O2-C2-C3-O3
3	D	302	PEG	O2-C3-C4-O4


There are no ring outliers.


4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	304	LF0	1	0
3	D	306	PEG	1	0
3	B	601	PEG	1	0
2	B	602	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand LF0 D 304





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	52/233 (22%)	0.85	7 (13%) 7 5	45, 62, 109, 129	0
1	B	143/233 (61%)	0.24	6 (4%) 40 40	22, 46, 70, 95	3 (2%)
1	C	50/233 (21%)	1.38	9 (18%) 3 3	61, 78, 110, 135	0
1	D	141/233 (60%)	0.13	1 (0%) 84 85	24, 45, 72, 94	1 (0%)
All	All	386/932 (41%)	0.43	23 (5%) 27 26	22, 49, 94, 135	4 (1%)

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	271	TYR	3.0
1	A	254	ASN	2.8
1	B	194	TYR	2.8
1	B	193	GLY	2.7
1	C	272	GLY	2.7
1	A	234	VAL	2.6
1	B	149	GLY	2.4
1	C	271	TYR	2.4
1	C	253	ASP	2.4
1	B	171[A]	HIS	2.3
1	D	149	GLY	2.2
1	C	232	ASP	2.2
1	A	273	LYS	2.2
1	A	230	SER	2.2
1	B	140	GLY	2.2
1	C	229	ASP	2.2
1	C	248	ALA	2.2
1	A	229	ASP	2.1
1	C	273	LYS	2.1
1	C	226	TYR	2.1
1	C	267	ILE	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	211	LYS	2.0
1	A	257	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PEG	D	303	7/7	0.74	0.19	58,64,72,76	0
3	PEG	D	306	7/7	0.76	0.21	59,65,69,69	0
2	EDO	B	605	4/4	0.77	0.19	62,71,73,74	0
3	PEG	B	601	7/7	0.77	0.20	57,69,72,79	0
3	PEG	D	305	7/7	0.80	0.14	65,67,74,83	0
3	PEG	D	302	7/7	0.82	0.15	50,68,71,72	0
2	EDO	A	901	4/4	0.83	0.15	56,61,65,66	0
2	EDO	B	602	4/4	0.84	0.14	67,71,73,76	0
4	GOL	D	301	6/6	0.84	0.14	53,61,71,74	0
4	GOL	B	603	6/6	0.88	0.10	57,62,67,76	0
5	LF0	B	604	30/30	0.95	0.08	35,41,47,50	0
5	LF0	D	304	30/30	0.95	0.08	37,41,46,49	0
6	MG	D	307	1/1	0.99	0.03	41,41,41,41	0
6	MG	B	606	1/1	1.00	0.03	50,50,50,50	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around LF0 B 604:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around LF0 D 304:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.