



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 02:13 PM UTC

PDB ID : 1A2Z / pdb_00001a2z
Title : PYRROLIDONE CARBOXYL PEPTIDASE FROM THERMOCOCCUS LITORALIS
Authors : Singleton, M.R.; Isupov, M.N.; Littlechild, J.A.
Deposited on : 1998-01-13
Resolution : 1.73 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.73 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1207 (1.74-1.74)
Ramachandran outliers	187476	1200 (1.74-1.74)
Sidechain outliers	187428	1200 (1.74-1.74)
RSRZ outliers	180081	1188 (1.74-1.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	220	 5% 81% 16% .
1	B	220	 3% 85% 11% .
1	C	220	 3% 82% 16% .
1	D	220	 7% 76% 20% ..

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 7666 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PYRROLIDONE CARBOXYL PEPTIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	220	Total	C	N	O	S	0	2	0
			1746	1130	284	324	8			
1	B	220	Total	C	N	O	S	0	2	0
			1746	1130	284	324	8			
1	C	220	Total	C	N	O	S	0	3	0
			1747	1130	284	325	8			
1	D	220	Total	C	N	O	S	0	3	0
			1747	1130	284	325	8			

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	C	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	171	Total	O	0	0
			171	171		
3	B	152	Total	O	0	0
			152	152		
3	C	172	Total	O	0	0
			172	172		
3	D	165	Total	O	0	0
			165	165		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: PYRROLIDONE CARBOXYL PEPTIDASE




• Molecule 1: PYRROLIDONE CARBOXYL PEPTIDASE



• Molecule 1: PYRROLIDONE CARBOXYL PEPTIDASE



• Molecule 1: PYRROLIDONE CARBOXYL PEPTIDASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	91.64Å 147.01Å 71.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.73 20.00 – 1.73	Depositor EDS
% Data completeness (in resolution range)	97.0 (20.00-1.73) 98.7 (20.00-1.73)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.50 (at 1.70Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.180 , 0.220 0.172 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	16.6	Xtriage
Anisotropy	0.528	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 47.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7666	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.94% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.87	1/1792 (0.1%)	1.65	18/2429 (0.7%)
1	B	0.83	0/1792	1.57	16/2429 (0.7%)
1	C	0.91	0/1798	1.61	16/2437 (0.7%)
1	D	0.86	0/1798	1.58	12/2437 (0.5%)
All	All	0.87	1/7180 (0.0%)	1.60	62/9732 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	164	GLY	N-CA	5.06	1.51	1.45

All (62) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	12	PHE	CA-CB-CG	12.49	126.29	113.80
1	A	192	GLU	CB-CG-CD	10.85	131.04	112.60
1	A	29	ARG	CD-NE-CZ	9.99	138.38	124.40
1	D	48	ARG	CD-NE-CZ	8.56	136.39	124.40
1	B	213	ARG	CD-NE-CZ	8.48	136.27	124.40
1	A	214	ASP	CA-CB-CG	8.17	120.77	112.60
1	D	61	LYS	CA-C-O	8.08	126.70	120.48
1	A	214	ASP	CA-C-O	-7.93	112.91	121.56

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	180	PHE	CA-CB-CG	7.64	121.44	113.80
1	C	107	ASP	CA-CB-CG	-7.02	105.58	112.60
1	D	12	PHE	CB-CA-C	-6.93	97.20	110.24
1	A	180	PHE	CA-CB-CG	6.93	120.73	113.80
1	D	141	TYR	N-CA-CB	-6.91	100.46	108.96
1	A	101	ASP	N-CA-CB	-6.75	101.72	111.70
1	A	31	GLN	OE1-CD-NE2	-6.74	115.86	122.60
1	B	90	ARG	N-CA-C	-6.72	104.72	113.12
1	B	22	GLN	CB-CG-CD	6.63	123.88	112.60
1	D	60	ILE	CA-C-N	6.54	129.90	122.83
1	D	60	ILE	C-N-CA	6.54	129.90	122.83
1	B	180	PHE	CA-CB-CG	6.27	120.07	113.80
1	B	119	ARG	CD-NE-CZ	6.23	133.12	124.40
1	A	119	ARG	CG-CD-NE	-6.20	98.36	112.00
1	C	34	ASN	OD1-CG-ND2	6.11	128.71	122.60
1	A	213	ARG	CA-C-N	6.08	130.48	120.94
1	A	213	ARG	C-N-CA	6.08	130.48	120.94
1	C	90	ARG	CD-NE-CZ	6.03	132.84	124.40
1	A	90	ARG	N-CA-C	-5.97	104.34	111.69
1	A	61	LYS	CA-C-O	5.84	124.50	120.47
1	A	38	TYR	CA-C-O	-5.79	114.11	120.30
1	B	192	GLU	CB-CG-CD	5.73	122.34	112.60
1	C	165	PHE	O-C-N	5.71	130.24	123.27
1	B	109	PRO	O-C-N	5.69	129.91	122.86
1	C	194	GLU	O-C-N	5.69	128.63	122.15
1	B	175	VAL	CB-CA-C	-5.63	104.65	112.36
1	B	22	GLN	OE1-CD-NE2	-5.60	117.00	122.60
1	B	214	ASP	CA-C-O	-5.60	114.67	121.05
1	A	44	VAL	CA-C-O	5.58	127.76	120.78
1	B	142	LEU	CA-C-O	-5.58	114.50	120.42
1	C	38	TYR	CA-C-O	-5.58	114.54	120.40
1	C	67	ASN	CA-C-O	-5.54	114.55	120.92
1	C	38	TYR	CA-C-N	-5.54	117.58	122.73
1	C	38	TYR	C-N-CA	-5.54	117.58	122.73
1	B	57	LEU	O-C-N	5.46	127.69	122.07
1	B	22	GLN	CG-CD-NE2	5.45	124.57	116.40
1	C	101	ASP	CA-CB-CG	-5.42	107.18	112.60
1	A	132	ALA	O-C-N	-5.31	117.54	123.48
1	D	13	GLY	N-CA-C	5.30	125.74	113.18
1	D	191	LEU	CA-C-O	-5.28	114.95	120.55
1	B	67	ASN	CA-C-O	-5.26	115.05	121.05
1	C	5	LEU	O-C-N	5.26	129.21	123.22

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	90	ARG	N-CA-C	-5.23	105.49	111.14
1	B	108	ALA	CA-C-N	5.21	125.22	120.21
1	B	108	ALA	C-N-CA	5.21	125.22	120.21
1	D	15	ASP	O-C-N	-5.19	117.13	123.10
1	C	133	THR	O-C-N	5.16	129.41	123.17
1	C	70	LEU	O-C-N	-5.13	116.96	123.01
1	A	78	THR	N-CA-CB	5.13	119.67	110.80
1	A	169	PRO	CB-CA-C	-5.08	103.54	111.22
1	D	158	GLY	O-C-N	5.06	129.05	122.42
1	C	50	THR	CB-CA-C	-5.05	102.41	110.79
1	C	108	ALA	O-C-N	5.05	126.14	121.80
1	A	101	ASP	CA-CB-CG	5.03	117.63	112.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	12	PHE	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1746	0	1803	22	0
1	B	1746	0	1803	18	0
1	C	1747	0	1804	25	0
1	D	1747	0	1804	36	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	1	0
2	D	5	0	0	1	0
3	A	171	0	0	2	0
3	B	152	0	0	0	0
3	C	172	0	0	2	0
3	D	165	0	0	1	0
All	All	7666	0	7214	98	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (98) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:10:GLU:HG3	1:D:44:VAL:HG23	1.44	0.96
1:A:64:ILE:HD12	1:A:206:LEU:HD22	1.47	0.95
1:C:85:ASN:HD21	1:C:105:GLU:H	1.13	0.90
1:D:85:ASN:HD21	1:D:105:GLU:H	1.16	0.90
1:A:190[B]:CYS:HG	1:C:190[B]:CYS:HG	0.87	0.82
1:D:211:LYS:HD3	1:D:213:ARG:HH21	1.50	0.76
1:A:214:ASP:HB3	3:A:757:HOH:O	1.88	0.73
1:B:213:ARG:HH22	1:B:216:ILE:HD12	1.60	0.66
1:C:90:ARG:HH12	1:C:140:THR:H	1.43	0.66
1:B:44:VAL:HG13	1:B:142:LEU:HD12	1.76	0.66
1:C:55:ARG:HB3	1:C:55:ARG:HH11	1.61	0.65
1:D:35:ALA:HB2	1:D:206:LEU:HD11	1.78	0.64
1:C:35:ALA:HB2	1:C:206:LEU:HD21	1.81	0.61
1:D:85:ASN:ND2	1:D:105:GLU:H	1.94	0.61
1:D:216:ILE:HB	1:D:218:ILE:HD12	1.83	0.61
1:C:50:THR:HG23	1:C:149:LYS:HD2	1.84	0.60
1:D:9:PHE:HB3	1:D:12:PHE:HE1	1.65	0.60
1:B:213:ARG:HG2	1:B:213:ARG:HH11	1.68	0.59
1:D:85:ASN:ND2	1:D:104:ILE:H	2.01	0.58
1:D:90:ARG:HB2	1:D:91:ILE:HD12	1.86	0.58
1:A:64:ILE:CD1	1:A:206:LEU:HD22	2.28	0.58
1:C:27:PHE:HB3	1:C:37:VAL:HG11	1.86	0.57
1:D:10:GLU:CG	1:D:44:VAL:HG23	2.27	0.57
1:D:119:ARG:HD3	1:D:218:ILE:O	2.05	0.57
1:D:64:ILE:HD12	1:D:206:LEU:HD13	1.87	0.55
1:D:206:LEU:O	1:D:210:GLU:HG2	2.07	0.55
1:A:10:GLU:HB3	1:A:11:PRO:HD2	1.89	0.55
1:A:12:PHE:HE1	1:A:70:LEU:HD23	1.72	0.55
1:A:12:PHE:CE1	1:A:70:LEU:HD23	2.42	0.55
1:D:85:ASN:HD22	1:D:104:ILE:H	1.55	0.55
1:D:112:TYR:OH	1:D:152:HIS:HD2	1.90	0.55
1:B:123:LYS:HD2	1:B:218:ILE:CD1	2.37	0.54
1:C:85:ASN:ND2	1:C:104:ILE:H	2.06	0.53
1:C:17:LYS:HE3	3:C:685:HOH:O	2.10	0.52
1:D:34:ASN:ND2	2:D:624:SO4:O2	2.40	0.52
1:D:119:ARG:HG3	1:D:220:LEU:CD1	2.40	0.51
1:A:35:ALA:HB2	1:A:206:LEU:HD21	1.91	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:44:VAL:HG13	1:A:142:LEU:HD12	1.92	0.51
1:B:119:ARG:HD3	1:B:216:ILE:O	2.11	0.51
1:B:12:PHE:HB2	1:B:44:VAL:HG21	1.93	0.50
1:B:123:LYS:HD2	1:B:218:ILE:HD13	1.94	0.50
1:D:10:GLU:HB2	1:D:11:PRO:CD	2.42	0.50
1:B:211:LYS:O	1:B:212:ASP:C	2.55	0.49
1:A:51:ILE:HD13	3:A:788:HOH:O	2.12	0.49
1:C:1:MET:N	2:C:623:SO4:O3	2.45	0.49
1:B:214:ASP:O	1:B:215:ASP:C	2.56	0.48
1:C:125:LEU:CD2	1:C:197:ALA:HA	2.43	0.48
1:A:10:GLU:HB3	1:A:11:PRO:CD	2.44	0.48
1:C:85:ASN:ND2	1:C:105:GLU:H	1.96	0.47
1:D:10:GLU:HG2	1:D:43:PRO:HA	1.97	0.47
1:B:213:ARG:HG2	1:B:213:ARG:NH1	2.31	0.46
1:C:143:CYS:SG	3:C:690:HOH:O	2.61	0.46
1:D:211:LYS:HD3	1:D:213:ARG:NH2	2.25	0.46
1:D:159:TYR:HA	1:D:160:PRO:C	2.40	0.46
1:D:211:LYS:O	1:D:212:ASP:C	2.57	0.46
1:A:12:PHE:HB3	1:A:44:VAL:HG21	1.98	0.46
1:A:61:LYS:HA	1:A:159:TYR:CE2	2.51	0.45
1:B:27:PHE:CE1	1:B:32:ILE:HD11	2.51	0.45
1:C:90:ARG:NH1	1:C:140:THR:H	2.12	0.45
1:A:44:VAL:HA	1:A:142:LEU:CD1	2.46	0.45
1:C:61:LYS:HA	1:C:159:TYR:CE2	2.51	0.45
1:A:90:ARG:HD2	1:B:220:LEU:OXT	2.17	0.45
1:B:143:CYS:HB3	1:B:165:PHE:CZ	2.52	0.45
1:C:153:PHE:CZ	1:C:157:GLU:HG3	2.52	0.45
1:D:10:GLU:HG3	1:D:44:VAL:CG2	2.31	0.44
1:C:211:LYS:HB2	1:C:213:ARG:HG3	1.98	0.44
1:D:92:PRO:HB3	1:D:98:GLN:HB2	2.00	0.44
1:C:85:ASN:ND2	1:C:104:ILE:HB	2.32	0.44
1:D:85:ASN:ND2	1:D:104:ILE:N	2.65	0.44
1:C:217:LYS:O	1:D:100:ILE:HG21	2.18	0.43
1:C:85:ASN:HD22	1:C:104:ILE:H	1.66	0.43
1:A:87:ILE:HG12	1:A:99:PRO:HG2	2.01	0.43
1:D:12:PHE:CD2	1:D:141:TYR:HE2	2.36	0.43
1:D:87:ILE:HG12	1:D:99:PRO:HG2	1.99	0.43
1:A:151:LEU:O	1:A:155:LYS:HG2	2.18	0.43
1:B:27:PHE:CD1	1:B:32:ILE:HD11	2.54	0.43
1:A:159:TYR:HA	1:A:160:PRO:C	2.44	0.42
1:D:215:ASP:N	3:D:669:HOH:O	2.52	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:35:ALA:HB3	1:C:206:LEU:HD11	2.01	0.42
1:D:207:ASP:O	1:D:211:LYS:HD2	2.19	0.41
1:B:12:PHE:CD1	1:B:44:VAL:HG11	2.56	0.41
1:C:61:LYS:N	1:C:62:PRO:CD	2.84	0.41
1:D:218:ILE:HG22	1:D:219:PRO:O	2.20	0.41
1:A:153:PHE:CZ	1:A:157:GLU:HG3	2.55	0.41
1:B:159:TYR:HA	1:B:160:PRO:C	2.46	0.41
1:B:67:ASN:HB2	1:B:165:PHE:CD2	2.56	0.41
1:C:10:GLU:HB2	1:C:11:PRO:CD	2.51	0.41
1:C:125:LEU:HD21	1:C:197:ALA:HA	2.02	0.41
1:D:96:GLY:O	1:D:97:TYR:C	2.64	0.41
1:A:67:ASN:O	1:A:165:PHE:HA	2.22	0.40
1:A:65:VAL:O	1:A:163:ALA:HA	2.22	0.40
1:B:105:GLU:OE2	1:B:149:LYS:NZ	2.49	0.40
1:D:6:ILE:HD12	1:D:37:VAL:HG11	2.03	0.40
1:D:143:CYS:HB3	1:D:165:PHE:CZ	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	220/220 (100%)	212 (96%)	7 (3%)	1 (0%)	24	12
1	B	220/220 (100%)	215 (98%)	4 (2%)	1 (0%)	24	12
1	C	221/220 (100%)	217 (98%)	4 (2%)	0	100	100
1	D	221/220 (100%)	210 (95%)	9 (4%)	2 (1%)	14	3
All	All	882/880 (100%)	854 (97%)	24 (3%)	4 (0%)	24	12

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	13	GLY
1	B	215	ASP
1	D	13	GLY
1	D	215	ASP

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	194/192 (101%)	189 (97%)	5 (3%)	40	17
1	B	194/192 (101%)	189 (97%)	5 (3%)	40	17
1	C	195/192 (102%)	192 (98%)	3 (2%)	57	38
1	D	195/192 (102%)	191 (98%)	4 (2%)	47	24
All	All	778/768 (101%)	761 (98%)	17 (2%)	45	22

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	156	ILE
1	A	191	LEU
1	A	192	GLU
1	A	206	LEU
1	A	214	ASP
1	B	22	GLN
1	B	54	LYS
1	B	161	LEU
1	B	213	ARG
1	B	216	ILE
1	C	34	ASN
1	C	50	THR
1	C	55	ARG
1	D	90	ARG
1	D	206	LEU
1	D	209	LEU
1	D	213	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	144	ASN
1	A	185	ASN
1	B	22	GLN
1	B	34	ASN
1	C	34	ASN
1	C	85	ASN
1	D	85	ASN
1	D	94	ASN
1	D	144	ASN
1	D	152	HIS
1	D	185	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	B	622	-	4,4,4	0.71	0	6,6,6	0.09	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	D	624	-	4,4,4	0.69	0	6,6,6	0.23	0
2	SO4	A	621	-	4,4,4	0.70	0	6,6,6	0.15	0
2	SO4	C	623	-	4,4,4	0.68	0	6,6,6	0.15	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	624	SO4	1	0
2	C	623	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	220/220 (100%)	-0.08	10 (4%) 38 47	10, 17, 39, 61	2 (0%)
1	B	220/220 (100%)	-0.15	7 (3%) 50 58	11, 17, 38, 49	2 (0%)
1	C	220/220 (100%)	-0.12	7 (3%) 50 58	11, 18, 34, 52	3 (1%)
1	D	220/220 (100%)	0.05	16 (7%) 21 26	11, 17, 45, 69	3 (1%)
All	All	880/880 (100%)	-0.08	40 (4%) 38 47	10, 17, 40, 69	10 (1%)

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	91	ILE	5.4
1	A	12	PHE	4.9
1	A	214	ASP	4.6
1	D	12	PHE	4.4
1	B	214	ASP	4.4
1	D	96	GLY	3.9
1	D	214	ASP	3.9
1	D	220	LEU	3.9
1	D	44	VAL	3.7
1	D	218	ILE	3.6
1	C	218	ILE	3.3
1	B	12	PHE	3.3
1	C	86	ILE	3.2
1	A	13	GLY	3.0
1	D	92	PRO	2.8
1	D	89	ALA	2.8
1	D	90	ARG	2.8
1	A	96	GLY	2.8
1	D	213	ARG	2.7
1	C	90	ARG	2.7
1	A	91	ILE	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	220	LEU	2.5
1	B	107	ASP	2.4
1	B	210	GLU	2.4
1	C	13	GLY	2.4
1	B	90	ARG	2.4
1	A	92	PRO	2.4
1	B	213	ARG	2.4
1	A	14	GLY	2.4
1	D	107	ASP	2.4
1	D	141	TYR	2.4
1	D	100	ILE	2.4
1	A	218	ILE	2.3
1	B	91	ILE	2.2
1	D	219	PRO	2.2
1	D	210	GLU	2.2
1	A	213	ARG	2.2
1	C	213	ARG	2.1
1	C	55	ARG	2.1
1	A	94	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SO4	B	622	5/5	0.68	0.15	105,106,106,107	0
2	SO4	A	621	5/5	0.70	0.20	101,102,102,103	0
2	SO4	C	623	5/5	0.81	0.18	154,154,155,155	0
2	SO4	D	624	5/5	0.82	0.18	71,71,75,76	0

6.5 Other polymers [i](#)

There are no such residues in this entry.