



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 12:29 AM UTC

PDB ID : 4A2D / pdb\_00004a2d  
Title : Coriolopsis gallica Laccase T2 Copper Depleted at pH 4.5  
Authors : De La Mora, E.; Valderrama, B.; Horjales, E.; Rudino-Pinera, E.  
Deposited on : 2011-09-26  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	496	
2	B	2	
2	C	2	

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4133 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LACCASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	496	3727	2368	631	719	9	0	0	0

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	39	ASP	TYR	SEE REMARK 999	UNP Q1W6B1
A	151	ASN	GLN	SEE REMARK 999	UNP Q1W6B1
A	178	LYS	ARG	SEE REMARK 999	UNP Q1W6B1
A	182	PRO	ALA	SEE REMARK 999	UNP Q1W6B1
A	183	VAL	ILE	SEE REMARK 999	UNP Q1W6B1
A	229	TYR	HIS	SEE REMARK 999	UNP Q1W6B1
A	256	LEU	ILE	SEE REMARK 999	UNP Q1W6B1
A	287	THR	ASN	SEE REMARK 999	UNP Q1W6B1
A	288	GLN	THR	SEE REMARK 999	UNP Q1W6B1
A	294	THR	VAL	SEE REMARK 999	UNP Q1W6B1
A	314	THR	ALA	SEE REMARK 999	UNP Q1W6B1
A	329	LYS	GLU	SEE REMARK 999	UNP Q1W6B1
A	356	ASN	ARG	SEE REMARK 999	UNP Q1W6B1
A	358	THR	SER	SEE REMARK 999	UNP Q1W6B1
A	423	VAL	ALA	SEE REMARK 999	UNP Q1W6B1

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	2	28	16	2	10	0	0	0

*Continued on next page...*


Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is COPPER (II) ION (CCD ID: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	Cu	0	0
			3	3		

- Molecule 4 is beta-D-mannopyranose (CCD ID: BMA) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			12	6	6		
4	A	1	Total	C	O	0	0
			12	6	6		
4	A	1	Total	C	O	0	0
			12	6	6		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	311	Total	O	0	0
			311	311		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: LACCASE



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	56.02Å 85.39Å 151.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	75.76 – 2.30 75.76 – 2.30	Depositor EDS
% Data completeness (in resolution range)	90.5 (75.76-2.30) 90.5 (75.76-2.30)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.71 (at 2.29Å)	Xtrriage
Refinement program	REFMAC 5.5.0110	Depositor
R, $R_{free}$	0.193 , 0.228 0.196 , 0.232	Depositor DCC
$R_{free}$ test set	1512 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	28.5	Xtrriage
Anisotropy	0.194	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 34.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4133	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.73% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CU, NAG, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.83	2/3838 (0.1%)	0.85	2/5277 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	343	ASP	CG-OD1	28.31	1.79	1.25
1	A	343	ASP	CB-CG	8.95	1.74	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	343	ASP	CA-CB-CG	-11.09	101.51	112.60
1	A	103	VAL	N-CA-C	-5.79	106.58	111.56

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3727	0	3541	38	0
2	B	28	0	25	1	0
2	C	28	0	25	0	0
3	A	3	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	36	0	36	2	0
5	A	311	0	0	1	0
All	All	4133	0	3627	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:343:ASP:OD1	1:A:343:ASP:CG	1.79	1.25
1:A:422:HIS:HE1	1:A:487:GLU:OE1	1.51	0.91
1:A:276:ASN:HD21	1:A:300:ARG:HE	1.21	0.84
1:A:276:ASN:ND2	1:A:300:ARG:HE	1.78	0.81
1:A:22:ALA:HB3	5:A:2002:HOH:O	1.88	0.73
1:A:329:LYS:H	1:A:329:LYS:HE2	1.54	0.73
1:A:422:HIS:CE1	1:A:487:GLU:OE1	2.41	0.68
1:A:242:GLU:HB3	1:A:265:SER:HB2	1.80	0.62
1:A:33:ASN:ND2	1:A:74:THR:H	2.01	0.58
1:A:226:ASP:HB3	1:A:227:PRO:HD3	1.89	0.55
1:A:226:ASP:HB3	1:A:227:PRO:CD	2.38	0.54
1:A:44:GLN:HG3	2:B:1:NAG:H2	1.89	0.54
1:A:439:ASN:HD22	1:A:439:ASN:N	2.05	0.53
1:A:334:PRO:HD3	1:A:442:TYR:OH	2.09	0.52
1:A:447:SER:O	1:A:455:VAL:HG21	2.11	0.50
1:A:228:ASN:H	1:A:284:ASN:ND2	2.08	0.50
1:A:387:PRO:HG2	1:A:390:SER:HB2	1.94	0.50
1:A:284:ASN:H	1:A:284:ASN:HD22	1.60	0.50
1:A:470:HIS:HD1	1:A:477:LEU:HD13	1.77	0.50
1:A:439:ASN:HD22	1:A:439:ASN:H	1.60	0.49
1:A:33:ASN:HD22	1:A:74:THR:H	1.61	0.48
1:A:302:ASP:C	1:A:302:ASP:N	2.72	0.48
1:A:210:VAL:O	1:A:302:ASP:N	2.47	0.47
1:A:276:ASN:HD21	1:A:300:ARG:NE	2.02	0.47
1:A:227:PRO:HA	1:A:284:ASN:HD21	1.79	0.47
1:A:87:HIS:C	1:A:87:HIS:CD2	2.94	0.46
1:A:135:THR:HA	1:A:474:ASP:OD2	2.16	0.46
1:A:448:THR:HA	1:A:455:VAL:HG21	1.98	0.46
1:A:328:LEU:HB3	1:A:329:LYS:HD3	1.98	0.45
1:A:226:ASP:HA	1:A:473:ILE:HG23	1.98	0.45
1:A:83:SER:O	1:A:131:SER:HA	2.17	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:1525:BMA:H1	4:A:1526:BMA:H61	2.00	0.44
1:A:470:HIS:ND1	1:A:477:LEU:HD13	2.33	0.43
1:A:57:ILE:HB	1:A:145:ILE:HG12	2.00	0.43
1:A:400:ILE:O	1:A:458:ARG:HA	2.20	0.41
1:A:290:PHE:HA	1:A:294:THR:HG22	2.03	0.41
1:A:100:PRO:HB2	1:A:103:VAL:HB	2.02	0.41
1:A:275:ASP:HB3	1:A:277:TYR:CE1	2.56	0.41
4:A:1525:BMA:H1	4:A:1526:BMA:H4	2.03	0.41
1:A:282:LEU:HD22	1:A:283:PRO:HD2	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	493/496 (99%)	472 (96%)	19 (4%)	2 (0%)	30 38

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	226	ASP
1	A	79	LEU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	400/401 (100%)	391 (98%)	9 (2%)	44 63

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	141	LEU
1	A	237	SER
1	A	242	GLU
1	A	282	LEU
1	A	284	ASN
1	A	294	THR
1	A	439	ASN
1	A	468	PHE
1	A	481	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	33	ASN
1	A	91	GLN
1	A	92	HIS
1	A	251	HIS
1	A	276	ASN
1	A	284	ASN
1	A	318	ASN
1	A	356	ASN
1	A	380	GLN
1	A	422	HIS
1	A	439	ASN
1	A	500	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates i

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	B	1	1,2	14,14,15	0.72	0	17,19,21	1.08	1 (5%)
2	NAG	B	2	2	14,14,15	0.65	0	17,19,21	1.26	2 (11%)
2	NAG	C	1	1,2	14,14,15	0.46	0	17,19,21	0.73	0
2	NAG	C	2	2	14,14,15	0.65	0	17,19,21	1.45	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	2/6/23/26	0/1/1/1
2	NAG	C	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	4/6/23/26	0/1/1/1

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	2	NAG	O5-C1-C2	-2.96	106.72	111.29
2	C	2	NAG	C4-C3-C2	2.84	115.17	111.02
2	C	2	NAG	C3-C4-C5	2.52	114.80	110.23
2	B	1	NAG	C2-N2-C7	-2.50	119.55	122.90
2	B	2	NAG	O5-C1-C2	-2.38	107.61	111.29
2	B	2	NAG	O5-C5-C6	2.31	112.16	107.66

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	2	NAG	C8-C7-N2-C2
2	C	2	NAG	O7-C7-N2-C2
2	C	2	NAG	C4-C5-C6-O6
2	C	2	NAG	O5-C5-C6-O6
2	B	2	NAG	C4-C5-C6-O6
2	B	2	NAG	O5-C5-C6-O6


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 3 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	BMA	A	1527	-	12,12,12	0.51	0	17,17,17	0.72	0
4	BMA	A	1526	-	12,12,12	0.61	0	17,17,17	0.81	0
4	BMA	A	1525	-	12,12,12	0.54	0	17,17,17	1.01	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	BMA	A	1527	-	-	0/2/22/22	0/1/1/1
4	BMA	A	1526	-	-	0/2/22/22	0/1/1/1
4	BMA	A	1525	-	-	2/2/22/22	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1525	BMA	C3-C4-C5	2.49	114.75	110.23

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1525	BMA	O5-C5-C6-O6
4	A	1525	BMA	C4-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1526	BMA	2	0
4	A	1525	BMA	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	496/496 (100%)	-0.15	8 (1%) 70 72	17, 25, 40, 56	1 (0%)

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	476	HIS	3.1
1	A	337	PRO	2.8
1	A	473	ILE	2.7
1	A	474	ASP	2.6
1	A	477	LEU	2.6
1	A	380	GLN	2.3
1	A	182	PRO	2.3
1	A	183	VAL	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.


### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	C	2	14/15	0.88	0.12	38,41,44,45	0
2	NAG	C	1	14/15	0.94	0.08	28,30,35,35	0
2	NAG	B	2	14/15	0.94	0.10	25,29,34,36	0
2	NAG	B	1	14/15	0.95	0.08	24,26,27,30	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	BMA	A	1525	12/12	0.65	0.23	116,117,117,117	0
4	BMA	A	1527	12/12	0.71	0.19	98,99,99,99	0
4	BMA	A	1526	12/12	0.78	0.21	98,99,99,100	0
3	CU	A	1518	1/1	0.91	0.11	43,43,43,43	1
3	CU	A	1520	1/1	0.98	0.04	36,36,36,36	1
3	CU	A	1519	1/1	0.98	0.04	34,34,34,34	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.