



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 06:10 AM UTC

PDB ID : 4A2F / pdb_00004a2f
Title : Coriolopsis gallica laccase collected at 12.65 keV
Authors : De la Mora, E.; Rudino-Pinera, E.
Deposited on : 2011-09-26
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	497	2% 78% 19% .
2	B	2	100%
2	C	2	100%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4273 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LACCASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	496	Total	C	N	O	S	0	10	0
			3799	2411	640	739	9			

There are 27 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	39	ASP	TYR	conflict	UNP Q1W6B1
A	151	ASN	GLN	conflict	UNP Q1W6B1
A	155	ALA	LYS	conflict	UNP Q1W6B1
A	178	LYS	ARG	conflict	UNP Q1W6B1
A	181	ALA	PRO	conflict	UNP Q1W6B1
A	182	PRO	ALA	conflict	UNP Q1W6B1
A	183	VAL	ILE	conflict	UNP Q1W6B1
A	198	ALA	ILE	conflict	UNP Q1W6B1
A	199	ALA	ASN	conflict	UNP Q1W6B1
A	202	ALA	ASN	conflict	UNP Q1W6B1
A	229	TYR	HIS	conflict	UNP Q1W6B1
A	256	LEU	ILE	conflict	UNP Q1W6B1
A	287	THR	ASN	conflict	UNP Q1W6B1
A	288	GLN	THR	conflict	UNP Q1W6B1
A	291	ALA	ASP	conflict	UNP Q1W6B1
A	294	THR	VAL	conflict	UNP Q1W6B1
A	314	THR	ALA	conflict	UNP Q1W6B1
A	329	LYS	GLU	conflict	UNP Q1W6B1
A	356	ASN	ARG	conflict	UNP Q1W6B1
A	358	THR	SER	conflict	UNP Q1W6B1
A	383	ALA	GLN	conflict	UNP Q1W6B1
A	388	ALA	THR	conflict	UNP Q1W6B1
A	408	ALA	THR	conflict	UNP Q1W6B1
A	423	VAL	ALA	conflict	UNP Q1W6B1
A	438	ALA	GLU	conflict	UNP Q1W6B1
A	450	ALA	SER	conflict	UNP Q1W6B1
A	518	UNK	-	expression tag	UNP Q1W6B1

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is COPPER (II) ION (CCD ID: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	Cu	0	0
			3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	415	Total	O	0	0
			415	415		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: LACCASE



• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	56.05Å 85.50Å 151.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	75.80 – 1.90 75.80 – 1.90	Depositor EDS
% Data completeness (in resolution range)	85.2 (75.80-1.90) 85.2 (75.80-1.90)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.42 (at 1.91Å)	Xtriage
Refinement program	REFMAC 5.5.0110	Depositor
R, R_{free}	0.186 , 0.224 0.191 , 0.231	Depositor DCC
R_{free} test set	2528 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	20.7	Xtriage
Anisotropy	0.138	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 45.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4273	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.34% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CU, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.47	24/3924 (0.6%)	1.27	14/5398 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	517	GLN	C-O	14.47	1.52	1.23
1	A	233	ILE	C-O	6.93	1.31	1.24
1	A	416	PRO	CA-C	6.31	1.58	1.52
1	A	423	VAL	CA-C	6.26	1.60	1.52
1	A	108	ILE	CA-CB	-6.14	1.47	1.54
1	A	445	VAL	N-CA	5.86	1.53	1.46
1	A	243	ALA	C-O	5.83	1.30	1.24
1	A	35	ALA	CA-CB	5.82	1.62	1.53
1	A	508	THR	N-CA	5.73	1.53	1.46
1	A	183	VAL	CA-CB	5.72	1.61	1.54
1	A	244	ASP	N-CA	5.70	1.53	1.46
1	A	69	VAL	CA-CB	5.60	1.60	1.53
1	A	388	ALA	CA-CB	5.53	1.62	1.53
1	A	258	ILE	CA-CB	5.47	1.61	1.54
1	A	82	THR	CA-CB	5.46	1.63	1.53
1	A	70	ILE	CA-CB	5.27	1.60	1.53
1	A	125	GLY	N-CA	5.24	1.51	1.44
1	A	253	VAL	C-O	5.22	1.29	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	45	ALA	CA-CB	5.14	1.60	1.54
1	A	298	ILE	CA-CB	5.14	1.60	1.54
1	A	166	VAL	N-CA	-5.11	1.40	1.46
1	A	265	SER	C-O	5.09	1.29	1.23
1	A	484	VAL	N-CA	5.05	1.52	1.46
1	A	241	ILE	CA-CB	-5.02	1.47	1.54

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	415	HIS	CA-C-N	7.12	128.98	120.23
1	A	415	HIS	C-N-CA	7.12	128.98	120.23
1	A	369	VAL	CA-C-N	-6.84	112.60	119.92
1	A	369	VAL	C-N-CA	-6.84	112.60	119.92
1	A	74	THR	N-CA-C	-5.63	106.22	114.39
1	A	498	VAL	CA-C-N	-5.58	114.50	120.03
1	A	498	VAL	C-N-CA	-5.58	114.50	120.03
1	A	181	ALA	CA-C-N	-5.54	113.96	119.56
1	A	181	ALA	C-N-CA	-5.54	113.96	119.56
1	A	122	ASP	CB-CA-C	-5.53	102.36	111.36
1	A	157	LEU	CA-C-N	-5.26	113.69	122.21
1	A	157	LEU	C-N-CA	-5.26	113.69	122.21
1	A	470	HIS	N-CA-C	5.16	115.16	107.88
1	A	122	ASP	N-CA-C	5.03	119.00	112.86

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	472	HIS	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3799	0	3601	57	0
2	B	28	0	25	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	28	0	25	0	0
3	A	3	0	0	0	0
4	A	415	0	0	3	0
All	All	4273	0	3651	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:418[B]:HIS:CE1	1:A:472:HIS:CD2	2.61	0.88
1:A:329:LYS:H	1:A:329:LYS:HE2	1.39	0.87
1:A:276:ASN:HD21	1:A:300:ARG:HE	1.31	0.78
1:A:284:ASN:H	1:A:284:ASN:HD22	1.34	0.75
1:A:22:ALA:HB2	1:A:163:ASP:OD2	1.85	0.75
1:A:418[B]:HIS:NE2	4:A:2147:HOH:O	2.20	0.75
1:A:135:THR:HA	1:A:474:ASP:OD2	1.91	0.71
1:A:418[B]:HIS:ND1	1:A:472:HIS:CD2	2.61	0.68
1:A:418[B]:HIS:HB3	1:A:445:VAL:HG22	1.75	0.68
1:A:33:ASN:ND2	1:A:74:THR:H	1.92	0.67
1:A:418[B]:HIS:ND1	1:A:472:HIS:HD2	1.93	0.67
1:A:418[B]:HIS:CE1	1:A:470:HIS:NE2	2.66	0.64
1:A:33:ASN:HD22	1:A:74:THR:H	1.48	0.62
1:A:276:ASN:ND2	1:A:300:ARG:HE	1.97	0.61
1:A:418[B]:HIS:CE1	1:A:470:HIS:CE1	2.89	0.60
1:A:322:GLU:OE2	1:A:443:ARG:NH1	2.35	0.58
1:A:227:PRO:HA	1:A:284:ASN:HD21	1.70	0.57
1:A:342:VAL:HB	1:A:399[B]:ASP:O	2.04	0.57
1:A:356:ASN:HB3	1:A:363:SER:OG	2.08	0.54
1:A:242:GLU:HB3	1:A:265:SER:HB2	1.90	0.53
1:A:39:ASP:HA	1:A:195:GLY:O	2.09	0.52
1:A:422:HIS:HE1	1:A:487:GLU:OE1	1.92	0.52
1:A:359:ILE:HD12	1:A:483:VAL:HG13	1.92	0.50
1:A:424:PHE:O	1:A:442:TYR:HA	2.11	0.50
1:A:329:LYS:HE2	1:A:329:LYS:N	2.18	0.50
1:A:418[B]:HIS:CB	1:A:445:VAL:HG22	2.42	0.49
1:A:238:LEU:HD22	1:A:266:PHE:CE2	2.48	0.48
1:A:422:HIS:CE1	1:A:487:GLU:OE1	2.67	0.48
1:A:57:ILE:HB	1:A:145:ILE:HG12	1.95	0.47
1:A:22:ALA:HB3	4:A:2002:HOH:O	2.14	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:352:PHE:CZ	1:A:355:GLY:HA2	2.50	0.47
1:A:470:HIS:HB3	1:A:482:ALA:CB	2.44	0.47
1:A:100:PRO:HD2	1:A:104:ASN:ND2	2.30	0.47
1:A:162:ASP:HB2	4:A:2184:HOH:O	2.14	0.46
1:A:87:HIS:C	1:A:87:HIS:CD2	2.93	0.46
1:A:216:TYR:O	1:A:267:VAL:HA	2.16	0.46
1:A:85:HIS:NE2	1:A:87:HIS:HA	2.31	0.45
1:A:418[B]:HIS:CE1	1:A:472:HIS:HD2	2.22	0.45
1:A:415:HIS:HE1	1:A:476:HIS:CE1	2.33	0.45
1:A:33:ASN:ND2	1:A:75:ASN:H	2.15	0.44
1:A:242:GLU:HG2	1:A:243:ALA:N	2.32	0.44
1:A:256:LEU:HD12	1:A:256:LEU:C	2.44	0.43
1:A:465:GLY:HA2	1:A:492:VAL:HG22	2.00	0.43
1:A:67:LEU:HD13	1:A:86:TRP:CZ3	2.54	0.42
1:A:375:ILE:HG23	1:A:489:ILE:HG23	2.02	0.42
1:A:226:ASP:HB3	1:A:227:PRO:CD	2.49	0.42
1:A:396:ALA:HB2	1:A:488:ASP:CG	2.45	0.42
1:A:336:SER:HA	1:A:337:PRO:HD3	1.90	0.41
1:A:228:ASN:H	1:A:284:ASN:ND2	2.19	0.41
1:A:284:ASN:HD22	1:A:284:ASN:N	2.08	0.41
1:A:415:HIS:CE1	1:A:476:HIS:CE1	3.07	0.41
1:A:87:HIS:CE1	1:A:418[A]:HIS:CE1	3.08	0.41
1:A:365[A]:THR:O	1:A:366:PRO:C	2.63	0.41
1:A:100:PRO:HD2	1:A:104:ASN:HD22	1.85	0.41
1:A:57:ILE:O	1:A:145:ILE:HA	2.21	0.41
1:A:130:HIS:HB2	1:A:137:TYR:HB2	2.03	0.40
1:A:436:ASN:HD21	1:A:439:ASN:ND2	2.19	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	504/497 (101%)	484 (96%)	18 (4%)	2 (0%)	30 22

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	226	ASP
1	A	79	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	411/401 (102%)	396 (96%)	15 (4%)	31 23

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	SER
1	A	221	VAL
1	A	245	SER
1	A	282	LEU
1	A	284	ASN
1	A	329	LYS
1	A	363	SER
1	A	404	LEU
1	A	407	THR
1	A	439	ASN
1	A	468	PHE
1	A	477	LEU
1	A	481	PHE
1	A	490	PRO
1	A	513	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	33	ASN
1	A	91	GLN
1	A	136	GLN
1	A	251	HIS
1	A	269	ASN
1	A	276	ASN
1	A	284	ASN
1	A	313	GLN
1	A	318	ASN
1	A	422	HIS
1	A	439	ASN
1	A	517	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	B	1	1,2	14,14,15	0.89	0	17,19,21	1.73	5 (29%)
2	NAG	B	2	2	14,14,15	1.18	1 (7%)	17,19,21	1.43	3 (17%)
2	NAG	C	1	1,2	14,14,15	0.59	0	17,19,21	1.50	4 (23%)
2	NAG	C	2	2	14,14,15	0.97	1 (7%)	17,19,21	1.30	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	2/6/23/26	0/1/1/1
2	NAG	C	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	0/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2	NAG	C2-N2	3.11	1.51	1.46
2	C	2	NAG	C3-C2	2.19	1.57	1.52

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	NAG	C3-C4-C5	-3.53	103.84	110.23
2	C	2	NAG	C4-C3-C2	3.18	115.68	111.02
2	B	2	NAG	O5-C5-C6	3.16	113.81	107.66
2	B	1	NAG	O7-C7-N2	3.05	127.38	121.98
2	B	1	NAG	O4-C4-C5	-2.82	102.37	109.32
2	B	1	NAG	C2-N2-C7	2.76	126.59	122.90
2	C	1	NAG	O4-C4-C3	-2.48	104.53	110.38
2	B	1	NAG	C4-C3-C2	-2.41	107.49	111.02
2	B	2	NAG	C4-C3-C2	2.41	114.54	111.02
2	C	1	NAG	O4-C4-C5	2.27	114.91	109.32
2	B	2	NAG	C1-O5-C5	2.16	115.08	112.19
2	C	2	NAG	C2-N2-C7	-2.09	120.10	122.90
2	B	1	NAG	O6-C6-C5	-2.07	104.30	111.33
2	C	1	NAG	C4-C3-C2	-2.04	108.03	111.02

There are no chirality outliers.


All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	2	NAG	C4-C5-C6-O6
2	B	2	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	496/497 (99%)	-0.13	12 (2%) 59 63	8, 22, 38, 56	10 (2%)

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	198	ALA	4.0
1	A	477	LEU	3.6
1	A	22	ALA	3.6
1	A	182	PRO	2.9
1	A	479	ALA	2.9
1	A	476	HIS	2.7
1	A	473	ILE	2.7
1	A	183	VAL	2.6
1	A	181	ALA	2.4
1	A	478	GLU	2.2
1	A	472	HIS	2.2
1	A	339	PRO	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NAG	C	2	14/15	0.88	0.12	29,46,51,52	0
2	NAG	B	2	14/15	0.92	0.13	23,28,33,36	0
2	NAG	C	1	14/15	0.94	0.10	23,32,37,38	0
2	NAG	B	1	14/15	0.96	0.08	23,27,32,32	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CU	A	1518	1/1	0.94	0.06	23,23,23,23	1
3	CU	A	1519	1/1	0.99	0.05	27,27,27,27	1
3	CU	A	1520	1/1	0.99	0.03	29,29,29,29	1

6.5 Other polymers [i](#)

There are no such residues in this entry.