



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:17 AM UTC

PDB ID : 6A2L / pdb_00006a2l
Title : Crystal structure of quadruple mutant (N51I+C59R+S108N+I164L) Plasmodium falciparum DHFR-TS complexed with BT1, NADPH, and dUMP
Authors : Chitnumsub, P.; Jaruwat, A.; Tarnchampoo, B.; Yuthavong, Y.
Deposited on : 2018-06-12
Resolution : 2.38 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.38 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7164 (2.40-2.36)
Clashscore	190562	7722 (2.40-2.36)
Ramachandran outliers	187476	7626 (2.40-2.36)
Sidechain outliers	187428	7627 (2.40-2.36)
RSRZ outliers	180081	7170 (2.40-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	608	
1	B	608	

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 9232 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bifunctional dihydrofolate reductase-thymidylate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	529	Total	C	N	O	S	0	0	0
			4409	2853	726	804	26			
1	B	521	Total	C	N	O	S	0	0	0
			4344	2813	714	791	26			

- Molecule 2 is 5-(3-{3-[(2,4-diamino-6-ethylpyrimidin-5-yl)oxy]propoxy}phenyl)-6-ethylpyrimidine-2,4-diamine (CCD ID: 9QO) (formula: $C_{21}H_{28}N_8O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			31	21	8	2		
2	B	1	Total	C	N	O	0	0
			31	21	8	2		

- Molecule 3 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$).


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	158	Total 158	O 158	0	0
5	B	123	Total 123	O 123	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Bifunctional dihydrofolate reductase-thymidylate synthase



- Molecule 1: Bifunctional dihydrofolate reductase-thymidylate synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.54Å 157.17Å 165.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	26.19 – 2.38 26.19 – 2.38	Depositor EDS
% Data completeness (in resolution range)	94.1 (26.19-2.38) 94.1 (26.19-2.38)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.48 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.197 , 0.255 0.193 , 0.251	Depositor DCC
R_{free} test set	2980 reflections (4.78%)	wwPDB-VP
Wilson B-factor (Å ²)	42.6	Xtriage
Anisotropy	0.278	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 31.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.011 for -h,l,k	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9232	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: UMP, NAP, 9QO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.84	1/4510 (0.0%)	0.98	5/6087 (0.1%)
1	B	0.80	1/4445 (0.0%)	0.98	3/6002 (0.0%)
All	All	0.82	2/8955 (0.0%)	0.98	8/12089 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	541	ALA	CA-C	7.77	1.56	1.52
1	B	541	ALA	CA-C	5.21	1.54	1.52

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	156	LEU	N-CA-C	6.65	119.54	109.23
1	A	343	SER	N-CA-C	-6.18	102.56	110.53
1	A	165	GLY	O-C-N	6.04	129.44	123.10
1	B	569	TYR	CA-C-N	-5.83	113.75	119.76
1	B	569	TYR	C-N-CA	-5.83	113.75	119.76
1	A	17	CYS	N-CA-C	5.73	118.24	108.90
1	A	557	ILE	N-CA-C	5.37	116.00	110.36
1	A	336	MET	N-CA-C	5.17	116.91	111.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4409	0	4369	33	0
1	B	4344	0	4295	33	0
2	A	31	0	0	1	0
2	B	31	0	0	2	0
3	A	48	0	25	1	0
3	B	48	0	25	1	0
4	A	20	0	11	1	0
4	B	20	0	11	0	0
5	A	158	0	0	4	0
5	B	123	0	0	0	0
All	All	9232	0	8736	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:315:TYR:HB2	1:A:564:LEU:O	1.83	0.78
1:A:210:VAL:HB	5:A:844:HOH:O	1.87	0.74
1:A:318:LEU:HB2	1:A:321:LYS:HD2	1.74	0.69
1:B:163:ILE:HD13	1:B:163:ILE:N	2.08	0.68
1:A:328:TYR:CZ	1:A:332:ILE:HD11	2.34	0.63
2:A:701:9QO:C9	2:A:701:9QO:C16	2.78	0.61
1:B:208:ILE:HG12	1:B:227:LYS:HB2	1.86	0.56
1:A:96:LYS:HA	5:A:941:HOH:O	2.07	0.55
1:B:332:ILE:HD13	1:B:560:LEU:HD22	1.87	0.55
1:A:165:GLY:HA2	1:A:170:TYR:CZ	2.41	0.55
1:A:502:LYS:HG2	5:A:895:HOH:O	2.06	0.54
1:B:373:LYS:HG3	1:B:598:HIS:CE1	2.42	0.54
1:A:102:VAL:HB	1:A:164:LEU:HG	1.90	0.53
1:B:11:ILE:HB	1:B:178:LEU:O	2.08	0.53
1:A:173:PHE:HB3	1:A:179:ILE:HD11	1.92	0.52
1:B:182:ILE:HD13	1:B:200:ILE:HD13	1.93	0.51
1:A:41:GLY:HA2	1:A:47:PRO:HD3	1.93	0.50
1:A:336:MET:HE3	1:A:557:ILE:HG23	1.94	0.49
1:A:109:TRP:CE2	1:A:117:LYS:HD2	2.48	0.49
1:B:493:LEU:HD12	1:B:493:LEU:C	2.38	0.49
1:A:318:LEU:CB	1:A:321:LYS:HD2	2.43	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:491:HIS:CE1	4:A:703:UMP:O4	2.66	0.49
1:B:494:CYS:HA	1:B:506:ILE:O	2.13	0.48
1:A:109:TRP:CZ2	1:A:117:LYS:HD2	2.49	0.48
1:B:6:CYS:HB3	1:B:178:LEU:O	2.14	0.48
1:B:7:ASP:HA	1:B:180:LYS:HE3	1.96	0.48
1:A:214:TYR:O	1:A:220:THR:HA	2.15	0.47
1:A:342:GLN:NE2	1:A:352:SER:OG	2.48	0.46
1:A:350:VAL:HG12	1:A:553:TYR:CD1	2.51	0.46
1:B:487:LEU:N	1:B:487:LEU:HD23	2.31	0.46
1:A:12:TYR:HD1	1:A:181:LYS:HB2	1.79	0.46
1:A:171:GLN:HG2	1:A:175:GLU:OE2	2.16	0.46
1:A:485:MET:SD	1:A:489:PRO:HD3	2.56	0.46
1:B:151:VAL:HG12	1:B:155:LYS:HE2	1.99	0.45
1:B:51:ILE:HD13	1:B:187:ILE:HD12	1.98	0.45
1:B:328:TYR:CZ	1:B:332:ILE:HD11	2.51	0.45
1:B:117:LYS:HA	1:B:118:PRO:HA	1.82	0.45
1:B:162:PHE:C	1:B:163:ILE:HD13	2.41	0.45
1:A:376:LEU:HD13	1:A:527:ILE:HD11	1.98	0.45
1:A:493:LEU:C	1:A:493:LEU:HD12	2.42	0.45
1:B:149:LEU:O	1:B:153:LEU:HG	2.16	0.45
1:A:21:GLU:HG2	1:A:190:THR:HG22	1.98	0.44
1:B:166:GLY:H	1:B:169:VAL:HB	1.82	0.44
1:B:421:ARG:HD2	1:B:425:ASP:HB3	2.00	0.44
1:A:491:HIS:ND1	5:A:802:HOH:O	2.32	0.43
1:B:150:ILE:H	1:B:150:ILE:HG13	1.62	0.43
1:B:165:GLY:HA3	1:B:169:VAL:HG11	1.99	0.43
1:A:556:HIS:CD2	1:A:604:MET:HB3	2.54	0.43
1:B:19:LYS:HG2	1:B:36:THR:HG22	2.00	0.43
1:A:102:VAL:HG21	1:A:119:LEU:HD12	2.01	0.43
1:A:428:PRO:HG2	1:A:481:ASP:HB3	2.00	0.42
1:B:41:GLY:HA2	1:B:47:PRO:HD3	2.01	0.42
1:A:354:PHE:CE2	1:B:506:ILE:HG13	2.54	0.42
1:B:100:ASN:OD1	1:B:159:TYR:HB3	2.19	0.42
2:B:701:9QO:C9	2:B:701:9QO:C16	2.98	0.42
1:B:421:ARG:CD	1:B:425:ASP:HB3	2.50	0.41
1:B:436:HIS:O	1:B:437:PHE:C	2.63	0.41
1:A:42:ASN:HA	1:A:194:ASP:OD1	2.20	0.41
1:B:214:TYR:O	1:B:220:THR:HA	2.20	0.41
1:B:3:GLU:H	1:B:3:GLU:HG3	1.70	0.41
1:B:195:VAL:HG21	3:B:702:NAP:H4D	2.03	0.41
1:B:12:TYR:CD1	1:B:181:LYS:HB2	2.56	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:42:ASN:HB2	1:B:193:CYS:HA	2.02	0.41
2:B:701:9QO:C12	2:B:701:9QO:N8	2.84	0.41
1:A:12:TYR:CD1	1:A:181:LYS:HB2	2.56	0.40
1:B:11:ILE:HG22	1:B:179:ILE:HD13	2.02	0.40
1:A:166:GLY:HA3	3:A:702:NAP:PA	2.61	0.40
1:A:380:ILE:O	1:A:384:LEU:HG	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	519/608 (85%)	494 (95%)	25 (5%)	0	100	100
1	B	511/608 (84%)	477 (93%)	27 (5%)	7 (1%)	9	11
All	All	1030/1216 (85%)	971 (94%)	52 (5%)	7 (1%)	18	26

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	132	LYS
1	B	165	GLY
1	B	230	ASN
1	B	141	TYR
1	B	140	VAL
1	B	68	SER
1	B	429	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	495/570 (87%)	479 (97%)	16 (3%)	34	53
1	B	487/570 (85%)	454 (93%)	33 (7%)	14	23
All	All	982/1140 (86%)	933 (95%)	49 (5%)	22	35

All (49) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	2	MET
1	A	3	GLU
1	A	72	LYS
1	A	124	ASN
1	A	133	LYS
1	A	164	LEU
1	A	189	SER
1	A	202	GLU
1	A	286	GLU
1	A	321	LYS
1	A	344	ASP
1	A	379	ILE
1	A	402	ARG
1	A	592	THR
1	A	600	GLU
1	B	3	GLU
1	B	11	ILE
1	B	40	LEU
1	B	72	LYS
1	B	96	LYS
1	B	107	THR
1	B	114	LYS
1	B	115	LYS
1	B	120	SER
1	B	127	LEU
1	B	131	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	134	GLU
1	B	140	VAL
1	B	143	ILE
1	B	146	VAL
1	B	149	LEU
1	B	150	ILE
1	B	152	LEU
1	B	163	ILE
1	B	164	LEU
1	B	174	LEU
1	B	182	ILE
1	B	189	SER
1	B	195	VAL
1	B	200	ILE
1	B	208	ILE
1	B	218	ASN
1	B	229	THR
1	B	284	ASP
1	B	310	ASN
1	B	457	LYS
1	B	516	LEU
1	B	603	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	99	GLN
1	A	296	ASN
1	A	338	ASN
1	A	340	ASN
1	A	342	GLN
1	A	407	ASN
1	A	542	GLN
1	B	66	ASN
1	B	108	ASN
1	B	217	ASN
1	B	231	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	NAP	A	702	-	50,52,52	1.34	7 (14%)	71,80,80	1.61	15 (21%)
2	9QO	A	701	-	33,33,33	0.71	0	42,45,45	2.75	15 (35%)
4	UMP	B	703	-	21,21,21	1.43	3 (14%)	30,31,31	1.80	9 (30%)
3	NAP	B	702	-	50,52,52	1.26	7 (14%)	71,80,80	1.63	13 (18%)
2	9QO	B	701	-	33,33,33	0.79	0	42,45,45	2.57	13 (30%)
4	UMP	A	703	-	21,21,21	1.45	4 (19%)	30,31,31	1.93	9 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAP	A	702	-	-	3/35/67/67	0/5/5/5
2	9QO	A	701	-	-	8/16/16/16	0/3/3/3
4	UMP	B	703	-	-	0/10/22/22	0/2/2/2
3	NAP	B	702	-	-	10/35/67/67	0/5/5/5
2	9QO	B	701	-	-	8/16/16/16	0/3/3/3
4	UMP	A	703	-	-	0/10/22/22	0/2/2/2

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	702	NAP	C5A-C4A	5.22	1.48	1.39
3	B	702	NAP	C5A-C4A	4.88	1.47	1.39
4	A	703	UMP	C2-N1	4.18	1.45	1.38
4	B	703	UMP	C2-N1	3.27	1.43	1.38
3	A	702	NAP	O4D-C1D	3.23	1.45	1.40
3	A	702	NAP	C5A-C6A	2.98	1.49	1.41
3	B	702	NAP	C5A-C6A	2.92	1.49	1.41
3	A	702	NAP	C5A-N7A	-2.70	1.34	1.39
3	B	702	NAP	PN-O3	2.62	1.62	1.59
3	B	702	NAP	PA-O3	2.56	1.62	1.59
4	A	703	UMP	C2-N3	-2.50	1.33	1.38
3	A	702	NAP	C4A-N9A	-2.47	1.32	1.37
4	B	703	UMP	O2-C2	2.44	1.27	1.23
4	A	703	UMP	C4-N3	-2.37	1.34	1.38
4	A	703	UMP	C6-C5	2.30	1.40	1.35
3	B	702	NAP	C8A-N7A	2.23	1.36	1.31
3	A	702	NAP	C8A-N7A	2.18	1.35	1.31
3	B	702	NAP	C5A-N7A	-2.17	1.35	1.39
3	B	702	NAP	O4D-C1D	2.07	1.43	1.40
3	A	702	NAP	PN-O3	2.02	1.61	1.59
4	B	703	UMP	C5-C4	-2.02	1.39	1.43

All (74) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	701	9QO	C25-N24-C23	8.11	122.18	116.26
2	A	701	9QO	C25-N24-C23	7.98	122.08	116.26
2	B	701	9QO	C2-N1-C6	7.48	121.72	116.26
2	A	701	9QO	C2-N1-C6	7.45	121.69	116.26
3	B	702	NAP	C5A-C4A-N3A	-5.34	119.36	126.72
3	A	702	NAP	C5A-C4A-N3A	-5.23	119.51	126.72
2	A	701	9QO	N26-C25-N24	-5.09	117.67	125.48
2	A	701	9QO	C25-N26-C27	5.09	123.38	117.28
2	B	701	9QO	C5-C4-N3	-4.58	119.89	122.55
2	A	701	9QO	C20-O21-C22	4.50	127.73	114.22
3	A	702	NAP	C4A-C5A-N7A	-4.39	105.57	110.58
3	B	702	NAP	C2B-C1B-N9A	-4.34	106.61	113.75
4	A	703	UMP	C5-C4-N3	4.25	120.75	114.80
3	B	702	NAP	N3A-C4A-N9A	4.24	134.39	127.17
2	B	701	9QO	N26-C25-N24	-4.07	119.24	125.48
2	B	701	9QO	N3-C2-N1	-4.05	119.27	125.48
2	B	701	9QO	C5-C6-N1	-3.92	118.98	123.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	703	UMP	C4-N3-C2	-3.81	121.89	126.61
2	A	701	9QO	C5-C6-N1	-3.72	119.21	123.47
4	B	703	UMP	O4-C4-C5	-3.68	118.81	125.16
4	B	703	UMP	C1'-N1-C2	3.67	124.83	117.66
3	B	702	NAP	C2A-N3A-C4A	3.67	120.78	111.83
3	A	702	NAP	N3A-C4A-N9A	3.64	133.36	127.17
2	B	701	9QO	C4-C5-C6	3.63	118.59	115.97
2	A	701	9QO	C31-C30-C23	-3.55	105.01	114.59
2	B	701	9QO	C2-N3-C4	3.54	121.53	117.28
3	B	702	NAP	N3A-C2A-N1A	-3.53	123.24	128.58
3	B	702	NAP	C4A-C5A-N7A	-3.51	106.57	110.58
2	A	701	9QO	N3-C2-N1	-3.42	120.24	125.48
4	A	703	UMP	N3-C2-N1	3.36	119.27	114.89
3	A	702	NAP	O2B-C2B-C1B	-3.34	98.32	110.05
4	B	703	UMP	C5-C4-N3	3.28	119.40	114.80
4	B	703	UMP	OP3-P-OP2	3.21	119.82	107.80
2	B	701	9QO	C25-N26-C27	3.18	121.10	117.28
2	A	701	9QO	C11-C5-C6	-3.15	119.50	123.30
4	A	703	UMP	OP3-P-OP2	3.09	119.41	107.80
3	A	702	NAP	C2A-N3A-C4A	3.04	119.26	111.83
4	A	703	UMP	OP2-P-O5'	-3.00	98.85	106.67
3	A	702	NAP	C5A-N7A-C8A	2.95	108.08	103.45
4	A	703	UMP	O2-C2-N3	-2.89	116.15	121.49
4	B	703	UMP	C1'-N1-C6	-2.84	115.94	121.53
2	A	701	9QO	C4-C5-C6	2.81	118.00	115.97
4	B	703	UMP	C4-N3-C2	-2.72	123.23	126.61
2	B	701	9QO	N7-C2-N1	2.72	121.30	117.22
3	B	702	NAP	C4A-N9A-C8A	2.71	108.58	105.74
2	A	701	9QO	N29-C25-N26	2.70	121.27	117.22
2	A	701	9QO	C5-C4-N3	-2.69	120.99	122.55
3	A	702	NAP	C5N-C4N-C3N	-2.66	117.75	120.36
3	B	702	NAP	C5A-N7A-C8A	2.59	107.52	103.45
3	A	702	NAP	C6A-C5A-N7A	2.56	137.03	132.09
4	A	703	UMP	C1'-N1-C2	2.56	122.67	117.66
2	A	701	9QO	N29-C25-N24	2.56	121.06	117.22
2	B	701	9QO	C22-C23-N24	-2.53	119.32	122.46
4	A	703	UMP	O4-C4-C5	-2.45	120.94	125.16
2	A	701	9QO	C20-C19-C18	-2.43	105.82	113.67
2	B	701	9QO	N29-C25-N26	2.42	120.85	117.22
3	B	702	NAP	C5N-C4N-C3N	-2.40	118.01	120.36
3	B	702	NAP	C6A-C5A-N7A	2.39	136.70	132.09
3	A	702	NAP	N3A-C2A-N1A	-2.35	125.02	128.58

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	702	NAP	O3X-P2B-O2X	2.29	116.38	107.80
3	B	702	NAP	C2A-N1A-C6A	2.25	122.43	118.73
3	A	702	NAP	O2A-PA-O1A	2.23	122.83	112.44
3	B	702	NAP	O2A-PA-O1A	2.23	122.81	112.44
3	A	702	NAP	C6N-N1N-C2N	-2.23	119.98	121.88
3	A	702	NAP	C4A-N9A-C8A	2.20	108.05	105.74
2	B	701	9QO	C18-O17-C15	2.19	123.63	117.93
4	B	703	UMP	OP2-P-O5'	-2.19	100.97	106.67
4	B	703	UMP	N3-C2-N1	2.19	117.74	114.89
2	A	701	9QO	C2-N3-C4	2.12	119.83	117.28
4	B	703	UMP	O4'-C1'-N1	2.11	111.61	107.86
3	A	702	NAP	C6N-N1N-C1D	2.03	123.70	119.73
3	B	702	NAP	N9A-C8A-N7A	-2.02	111.07	113.94
4	A	703	UMP	C1'-N1-C6	-2.01	117.57	121.53
3	A	702	NAP	C2A-N1A-C6A	2.00	122.02	118.73

There are no chirality outliers.

All (29) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	701	9QO	C12-C11-C5-C4
3	A	702	NAP	O4D-C1D-N1N-C2N
3	A	702	NAP	O4D-C1D-N1N-C6N
3	A	702	NAP	C2D-C1D-N1N-C2N
3	B	702	NAP	C5D-O5D-PN-O1N
3	B	702	NAP	O4D-C1D-N1N-C2N
3	B	702	NAP	O4D-C1D-N1N-C6N
3	B	702	NAP	C2D-C1D-N1N-C2N
3	B	702	NAP	C2D-C1D-N1N-C6N
2	B	701	9QO	C14-C15-O17-C18
2	B	701	9QO	C16-C15-O17-C18
2	B	701	9QO	C18-C19-C20-O21
2	A	701	9QO	O17-C18-C19-C20
2	A	701	9QO	C12-C11-C5-C6
2	A	701	9QO	C16-C11-C5-C4
2	A	701	9QO	C16-C11-C5-C6
2	B	701	9QO	C12-C11-C5-C4
2	B	701	9QO	C12-C11-C5-C6
2	A	701	9QO	C18-C19-C20-O21
3	B	702	NAP	C5D-O5D-PN-O3
3	B	702	NAP	C4N-C3N-C7N-O7N
2	B	701	9QO	C16-C11-C5-C6

Continued on next page...

Continued from previous page...



Mol	Chain	Res	Type	Atoms
2	A	701	9QO	C16-C15-O17-C18
3	B	702	NAP	C4N-C3N-C7N-N7N
2	B	701	9QO	C16-C11-C5-C4
2	A	701	9QO	C14-C15-O17-C18
3	B	702	NAP	C2N-C3N-C7N-O7N
2	B	701	9QO	N1-C6-C9-C10
3	B	702	NAP	C2N-C3N-C7N-N7N



There are no ring outliers.


5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	702	NAP	1	0
2	A	701	9QO	1	0
3	B	702	NAP	1	0
2	B	701	9QO	2	0
4	A	703	UMP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	529/608 (87%)	-0.21	13 (2%) 58 58	23, 39, 90, 120	0
1	B	521/608 (85%)	0.09	22 (4%) 40 40	23, 45, 115, 120	0
All	All	1050/1216 (86%)	-0.06	35 (3%) 49 49	23, 41, 111, 120	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	146	VAL	4.1
1	A	375	PHE	3.9
1	B	165	GLY	3.9
1	B	136	PHE	3.6
1	B	2	MET	3.5
1	B	285	GLU	3.4
1	A	85	THR	3.4
1	B	80	TYR	3.3
1	A	2	MET	3.2
1	B	131	LEU	3.2
1	B	75	TYR	3.1
1	B	143	ILE	2.8
1	A	605	ASP	2.5
1	A	22	SER	2.5
1	A	75	TYR	2.5
1	A	114	LYS	2.4
1	B	141	TYR	2.4
1	B	140	VAL	2.4
1	B	73	LEU	2.3
1	A	231	ASN	2.3
1	B	29	ASN	2.3
1	A	5	VAL	2.3
1	B	168	VAL	2.3
1	B	178	LEU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	344	ASP	2.2
1	B	105	GLY	2.2
1	B	173	PHE	2.1
1	B	35	TYR	2.1
1	B	11	ILE	2.1
1	B	151	VAL	2.1
1	A	28	LYS	2.1
1	A	81	LEU	2.1
1	B	139	ASP	2.1
1	A	9	PHE	2.1
1	A	286	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	9QO	B	701	31/31	0.84	0.13	55,78,98,104	0
3	NAP	B	702	48/48	0.86	0.12	83,107,120,120	0
2	9QO	A	701	31/31	0.90	0.12	28,58,72,78	0
3	NAP	A	702	48/48	0.96	0.07	41,52,58,63	0
4	UMP	A	703	20/20	0.96	0.09	35,44,58,63	0
4	UMP	B	703	20/20	0.97	0.08	34,50,61,67	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 9QO B 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAP B 702:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 9QO A 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAP A 702:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around UMP A 703:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around UMP B 703:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.