



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 08:02 PM UTC

PDB ID : 4A3I / pdb_00004a3i
Title : RNA Polymerase II binary complex with DNA
Authors : Cheung, A.C.M.; Sainsbury, S.; Cramer, P.
Deposited on : 2011-09-30
Resolution : 3.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 3.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1065 (3.96-3.64)
Clashscore	190562	1012 (3.94-3.66)
Ramachandran outliers	187476	1048 (3.96-3.64)
Sidechain outliers	187428	1043 (3.96-3.64)
RSRZ outliers	180081	1064 (3.96-3.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1732	2% 48%26%6%18%
2	B	1224	2% 57%27%7%9%
3	C	318	46%30%8%16%
4	D	221	2% 44%27%9%19%
5	E	215	58%34%7%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
6	F	155	
7	G	171	
8	H	146	
9	I	122	
10	J	70	
11	K	120	
12	L	70	
13	N	15	
14	T	27	

2 Entry composition

There are 16 unique types of molecules in this entry. The entry contains 31768 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1422	Total	C	N	O	S	0	0	0
			11174	7037	1954	2121	62			

- Molecule 2 is a protein called DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	1115	Total	C	N	O	S	0	0	0
			8859	5609	1554	1641	55			

- Molecule 3 is a protein called DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	266	Total	C	N	O	S	0	0	0
			2095	1317	348	417	13			

- Molecule 4 is a protein called DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	178	Total	C	N	O	S	0	0	0
			1434	887	257	288	2			

- Molecule 5 is a protein called DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	214	Total	C	N	O	S	0	0	0
			1752	1111	309	321	11			

- Molecule 6 is a protein called DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	84	Total	C	N	O	S	0	0	0
			679	434	115	127	3			

- Molecule 7 is a protein called RPB7, DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	G	171	Total	C	N	O	S	0	0	0
			1340	861	222	249	8			

- Molecule 8 is a protein called DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	H	133	Total	C	N	O	S	0	0	0
			1068	673	180	211	4			

- Molecule 9 is a protein called DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	I	119	Total	C	N	O	S	0	0	0
			971	596	179	186	10			

- Molecule 10 is a protein called DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	J	65	Total	C	N	O	S	0	0	0
			532	339	93	94	6			

- Molecule 11 is a protein called DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
11	K	115	Total	C	N	O	S	0	0	1
			920	590	157	171	2			

- Molecule 12 is a protein called DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
12	L	46	Total	C	N	O	S	0	0	0
			363	224	72	63	4			

- Molecule 13 is a DNA chain called 5'-D(*GP*GP*CP*AP*CP*AP*AP*CP*TP*GP*CP*GP*GP*CP*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
13	N	11	Total	C	N	O	P	0	0	0
			222	106	44	62	10			

- Molecule 14 is a DNA chain called TEMPLATE DNA 27-MER.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
14	T	17	Total	Br	C	N	O	P	0	0
			350	1	166	61	105	17		

- Molecule 15 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
15	A	2	Total	Zn	0	0
			2	2		
15	B	1	Total	Zn	0	0
			1	1		
15	C	1	Total	Zn	0	0
			1	1		
15	I	2	Total	Zn	0	0
			2	2		
15	J	1	Total	Zn	0	0
			1	1		
15	L	1	Total	Zn	0	0
			1	1		


- Molecule 16 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
16	A	1	Total	Mg	0	0
			1	1		

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB1






● Molecule 2: DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB2






• Molecule 3: DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB3


Chain C: 46% 30% 8% 16%




• Molecule 4: DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB4




• Molecule 5: DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 1




• Molecule 6: DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 2




• Molecule 7: RPB7, DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB7




• Molecule 8: DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 3




• Molecule 9: DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB9




• Molecule 10: DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 5



• Molecule 11: DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB11



• Molecule 12: DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 4




- Molecule 13: 5'-D(*GP*GP*CP*AP*CP*AP*AP*CP*TP*GP*CP*GP*GP*CP*T)-3'

Chain N:  40% 33% 27%



- Molecule 14: TEMPLATE DNA 27-MER

Chain T:  48% 11% 37%



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	221.14Å 393.18Å 282.48Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.80 – 3.80 49.80 – 3.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.80-3.80) 99.9 (49.80-3.80)	Depositor EDS
R_{merge}	0.89	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.92 (at 3.77Å)	Xtriage
Refinement program	BUSTER 2.11.2	Depositor
R, R_{free}	0.159 , 0.191 0.183 , 0.210	Depositor DCC
R_{free} test set	2395 reflections (1.99%)	wwPDB-VP
Wilson B-factor (Å ²)	111.3	Xtriage
Anisotropy	0.409	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 138.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.026 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.034 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	31768	wwPDB-VP
Average B, all atoms (Å ²)	131.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN, BRU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.91	6/11374 (0.1%)	1.60	161/15383 (1.0%)
2	B	0.89	1/9029 (0.0%)	1.51	100/12171 (0.8%)
3	C	0.83	0/2133	1.52	22/2891 (0.8%)
4	D	0.87	1/1444 (0.1%)	1.63	27/1935 (1.4%)
5	E	0.83	0/1788	1.50	24/2406 (1.0%)
6	F	0.95	0/691	1.46	8/933 (0.9%)
7	G	0.89	1/1368 (0.1%)	1.45	11/1844 (0.6%)
8	H	0.86	0/1086	1.52	21/1470 (1.4%)
9	I	0.78	0/989	1.37	7/1331 (0.5%)
10	J	0.95	1/541 (0.2%)	1.65	10/727 (1.4%)
11	K	0.79	0/938	1.46	6/1267 (0.5%)
12	L	0.98	0/365	1.55	5/485 (1.0%)
13	N	0.48	0/249	0.54	0/382
14	T	0.49	0/369	0.57	0/568
All	All	0.88	10/32364 (0.0%)	1.53	402/43793 (0.9%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	867	ILE	CG1-CD1	7.74	1.81	1.51
1	A	507	VAL	CA-CB	7.37	1.57	1.54
7	G	1	MET	C-N	6.78	1.43	1.33
4	D	18	VAL	CA-C	6.48	1.60	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	57	ARG	CA-C	5.79	1.60	1.52

The worst 5 of 402 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	26	THR	N-CA-C	-12.45	98.53	112.72
3	C	39	ALA	N-CA-C	10.44	122.42	111.14
5	E	5	ASN	N-CA-C	-9.99	101.22	113.41
7	G	34	VAL	N-CA-C	9.42	119.39	110.53
1	A	56	PRO	CA-C-N	9.34	139.38	121.54

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	34	LYS	Peptide,Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	11174	0	11233	265	0
2	B	8859	0	8901	180	0
3	C	2095	0	2051	64	0
4	D	1434	0	1460	38	0
5	E	1752	0	1776	41	0
6	F	679	0	701	23	0
7	G	1340	0	1357	47	0
8	H	1068	0	1040	26	0
9	I	971	0	927	17	0
10	J	532	0	542	24	0
11	K	920	0	929	28	0
12	L	363	0	386	9	0
13	N	222	0	124	4	0
14	T	350	0	191	3	0
15	A	2	0	0	0	0
15	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
15	C	1	0	0	0	0
15	I	2	0	0	0	0
15	J	1	0	0	0	0
15	L	1	0	0	0	0
16	A	1	0	0	0	0
All	All	31768	0	31618	676	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 676 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:867:ILE:CG1	1:A:867:ILE:CD1	1.81	1.53
1:A:37:PHE:HD1	1:A:52:GLY:HA3	1.26	1.01
1:A:768:GLN:HG2	1:A:816:HIS:HA	1.49	0.94
1:A:35:ILE:HG22	1:A:84:ILE:HG22	1.51	0.92
2:B:114:PRO:HG3	2:B:181:LEU:HD11	1.52	0.90

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1414/1732 (82%)	1207 (85%)	137 (10%)	70 (5%)	1	17
2	B	1095/1224 (90%)	929 (85%)	122 (11%)	44 (4%)	2	20
3	C	264/318 (83%)	231 (88%)	25 (10%)	8 (3%)	3	25
4	D	174/221 (79%)	152 (87%)	13 (8%)	9 (5%)	1	17
5	E	212/215 (99%)	186 (88%)	19 (9%)	7 (3%)	3	24

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
6	F	82/155 (53%)	75 (92%)	6 (7%)	1 (1%)	10	40
7	G	169/171 (99%)	149 (88%)	17 (10%)	3 (2%)	6	33
8	H	129/146 (88%)	102 (79%)	16 (12%)	11 (8%)	0	9
9	I	117/122 (96%)	91 (78%)	23 (20%)	3 (3%)	4	27
10	J	63/70 (90%)	52 (82%)	6 (10%)	5 (8%)	1	10
11	K	113/120 (94%)	107 (95%)	6 (5%)	0	100	100
12	L	44/70 (63%)	26 (59%)	10 (23%)	8 (18%)	0	2
All	All	3876/4564 (85%)	3307 (85%)	400 (10%)	169 (4%)	2	19

5 of 169 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	43	GLU
1	A	48	ALA
1	A	54	ASN
1	A	57	ARG
1	A	58	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1240/1519 (82%)	1013 (82%)	227 (18%)	2	11
2	B	966/1061 (91%)	795 (82%)	171 (18%)	2	12
3	C	234/274 (85%)	191 (82%)	43 (18%)	1	11
4	D	160/200 (80%)	131 (82%)	29 (18%)	2	12
5	E	196/197 (100%)	171 (87%)	25 (13%)	4	21
6	F	74/137 (54%)	60 (81%)	14 (19%)	1	10
7	G	152/152 (100%)	129 (85%)	23 (15%)	3	16
8	H	117/128 (91%)	99 (85%)	18 (15%)	2	15
9	I	113/116 (97%)	102 (90%)	11 (10%)	8	29

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
10	J	60/65 (92%)	44 (73%)	16 (27%)	0	3
11	K	99/102 (97%)	83 (84%)	16 (16%)	2	14
12	L	40/57 (70%)	27 (68%)	13 (32%)	0	1
All	All	3451/4008 (86%)	2845 (82%)	606 (18%)	2	12

5 of 606 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	D	182	SER
10	J	41	LEU
5	E	92	THR
4	D	164	ILE
7	G	108	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 74 such sidechains are listed below:

Mol	Chain	Res	Type
5	E	99	HIS
11	K	29	ASN
5	E	174	GLN
8	H	131	ASN
1	A	1378	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
14	BRU	T	22	14	18,21,22	1.60	4 (22%)	25,30,33	2.64	10 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
14	BRU	T	22	14	-	2/7/21/22	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
14	T	22	BRU	C6-C5	3.72	1.41	1.34
14	T	22	BRU	BR-C5	3.31	1.96	1.88
14	T	22	BRU	C6-N1	2.79	1.42	1.38
14	T	22	BRU	C4-C5	2.57	1.50	1.45

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
14	T	22	BRU	BR-C5-C4	7.08	126.17	118.02
14	T	22	BRU	N3-C2-N1	4.54	120.80	114.89
14	T	22	BRU	O4'-C1'-N1	4.41	115.69	107.86
14	T	22	BRU	C6-C5-C4	-4.19	116.42	120.67
14	T	22	BRU	C5-C4-N3	4.05	118.00	113.34

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
14	T	22	BRU	C3'-C4'-C5'-O5'
14	T	22	BRU	O4'-C4'-C5'-O5'

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
14	T	22	BRU	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 9 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	B	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	934:LYS	C	935:ARG	N	5.74
1	B	351:TYR	C	352:ALA	N	3.07

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1422/1732 (82%)	-0.27	15 (1%) 78 56	64, 116, 174, 247	0
2	B	1115/1224 (91%)	-0.10	22 (1%) 65 44	67, 130, 191, 222	0
3	C	266/318 (83%)	-0.38	0 100 100	90, 119, 163, 183	0
4	D	178/221 (80%)	-0.00	4 (2%) 62 44	102, 135, 182, 198	0
5	E	214/215 (99%)	-0.14	0 100 100	90, 151, 199, 208	0
6	F	84/155 (54%)	-0.63	1 (1%) 76 54	70, 95, 126, 149	0
7	G	171/171 (100%)	-0.26	1 (0%) 85 68	87, 116, 155, 179	0
8	H	133/146 (91%)	0.26	7 (5%) 32 24	122, 161, 195, 205	0
9	I	119/122 (97%)	-0.07	1 (0%) 82 62	123, 158, 192, 214	0
10	J	65/70 (92%)	-0.31	1 (1%) 72 50	97, 115, 153, 166	0
11	K	115/120 (95%)	-0.47	1 (0%) 81 60	83, 115, 163, 181	0
12	L	46/70 (65%)	0.46	3 (6%) 25 21	103, 159, 184, 191	0
13	N	11/15 (73%)	0.50	0 100 100	203, 219, 272, 273	0
14	T	16/27 (59%)	0.57	0 100 100	170, 216, 266, 270	0
All	All	3955/4606 (85%)	-0.19	56 (1%) 73 51	64, 125, 188, 273	0

The worst 5 of 56 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	446	LEU	5.0
8	H	63	LEU	4.4
12	L	27	LEU	4.3
1	A	251	SER	4.3
2	B	134	LYS	4.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
14	BRU	T	22	20/21	0.74	0.14	221,231,236,237	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
16	MG	A	2458	1/1	0.93	0.05	106,106,106,106	0
15	ZN	I	1122	1/1	0.99	0.03	197,197,197,197	0
15	ZN	B	2225	1/1	1.00	0.05	92,92,92,92	0
15	ZN	C	1269	1/1	1.00	0.03	88,88,88,88	0
15	ZN	I	1121	1/1	1.00	0.02	126,126,126,126	0
15	ZN	A	2456	1/1	1.00	0.01	146,146,146,146	0
15	ZN	J	1066	1/1	1.00	0.03	90,90,90,90	0
15	ZN	L	1071	1/1	1.00	0.02	164,164,164,164	0
15	ZN	A	2457	1/1	1.00	0.03	89,89,89,89	0

6.5 Other polymers [i](#)

There are no such residues in this entry.