



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 12:43 PM UTC

PDB ID : 4A3Q / pdb_00004a3q
Title : The 2.15 Angstrom resolution crystal structure of Staphylococcus aureus alanine racemase
Authors : Scaletti, E.R.; Luckner, S.R.; Krause, K.L.
Deposited on : 2011-10-04
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2057 (2.16-2.16)
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	382	4% 83% 10% 7%
1	B	382	5% 82% 11% 7%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SO4	B	1385	-	-	X	-
4	ACT	A	1389	-	-	X	-

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 6001 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ALANINE RACEMASE 1.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	356	Total	C	N	O	S	0	0	0
			2819	1795	486	522	16			
1	B	356	Total	C	N	O	S	0	0	0
			2813	1793	486	520	14			

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (CCD ID: PLP) (formula: $C_8H_{10}NO_6P$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	B	1	Total	C	N	O	P	0	0
			15	8	1	5	1		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	159	Total	O	0	0
			159	159		
5	B	107	Total	O	0	0
			107	107		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ALANINE RACEMASE 1



• Molecule 1: ALANINE RACEMASE 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	65.14Å 113.94Å 125.99Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.41 – 2.15 39.41 – 2.15	Depositor EDS
% Data completeness (in resolution range)	95.6 (39.41-2.15) 95.6 (39.41-2.15)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.34 (at 2.16Å)	Xtriage
Refinement program	REFMAC 5.5.0110	Depositor
R, R_{free}	0.188 , 0.236 0.186 , 0.232	Depositor DCC
R_{free} test set	2515 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	23.1	Xtriage
Anisotropy	0.082	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 34.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6001	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, PLP, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.92	8/2870 (0.3%)	0.94	8/3881 (0.2%)
1	B	0.87	7/2865 (0.2%)	0.91	8/3878 (0.2%)
All	All	0.89	15/5735 (0.3%)	0.92	16/7759 (0.2%)

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	314	GLN	CD-OE1	5.91	1.34	1.23
1	A	182	GLN	CD-OE1	5.75	1.34	1.23
1	A	16	ASN	CG-OD1	5.59	1.34	1.23
1	A	53	HIS	ND1-CE1	5.53	1.38	1.32
1	B	369	HIS	ND1-CE1	5.48	1.38	1.32
1	B	129	HIS	ND1-CE1	5.43	1.38	1.32
1	B	53	HIS	ND1-CE1	5.37	1.38	1.32
1	A	129	HIS	ND1-CE1	5.37	1.38	1.32
1	A	369	HIS	ND1-CE1	5.31	1.37	1.32
1	B	314	GLN	CD-OE1	5.28	1.33	1.23
1	B	12	ASN	CG-OD1	5.17	1.33	1.23
1	A	109	GLN	CD-OE1	5.16	1.33	1.23
1	A	76	HIS	ND1-CE1	5.10	1.37	1.32
1	B	57	ASN	CG-OD1	5.10	1.33	1.23
1	B	53	HIS	CD2-NE2	-5.02	1.32	1.37

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	140	GLY	N-CA-C	7.51	120.86	111.85
1	A	369	HIS	CB-CG-CD2	-6.72	122.47	131.20
1	A	76	HIS	CB-CG-CD2	-6.59	122.63	131.20
1	A	53	HIS	CB-CG-CD2	-6.59	122.64	131.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	369	HIS	CB-CG-CD2	-6.49	122.77	131.20
1	B	53	HIS	CB-CG-CD2	-6.42	122.86	131.20
1	A	129	HIS	CB-CG-CD2	-6.40	122.87	131.20
1	B	129	HIS	CB-CG-CD2	-6.34	122.96	131.20
1	B	369	HIS	CB-CG-ND1	5.71	131.26	122.70
1	B	319	VAL	N-CA-C	5.60	114.06	107.84
1	A	76	HIS	CB-CG-ND1	5.56	131.04	122.70
1	B	129	HIS	CB-CG-ND1	5.56	131.04	122.70
1	B	53	HIS	CB-CG-ND1	5.54	131.01	122.70
1	A	53	HIS	CB-CG-ND1	5.54	131.01	122.70
1	A	369	HIS	CB-CG-ND1	5.46	130.89	122.70
1	A	129	HIS	CB-CG-ND1	5.40	130.81	122.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2819	0	2840	16	0
1	B	2813	0	2835	19	0
2	A	15	0	6	1	0
2	B	15	0	6	2	0
3	A	30	0	0	1	0
3	B	35	0	0	5	0
4	A	4	0	3	2	0
4	B	4	0	3	0	0
5	A	159	0	0	2	0
5	B	107	0	0	0	0
All	All	6001	0	5693	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:362:ARG:HB3	3:B:1385:SO4:O2	1.82	0.80
1:B:361:SER:HA	3:B:1385:SO4:O3	1.85	0.77
1:B:32:LYS:HE2	1:B:215:CYS:O	1.86	0.76
1:A:377:THR:HG21	5:A:2160:HOH:O	1.94	0.67
1:B:52:ARG:NH2	3:B:1384:SO4:O3	2.29	0.65
1:B:95:LYS:HE2	1:B:98:GLN:OE1	1.98	0.64
4:A:1389:ACT:H1	2:B:1039:PLP:C4A	2.29	0.62
1:A:337:GLU:N	3:A:1386:SO4:O4	2.29	0.56
4:A:1389:ACT:H2	5:A:2135:HOH:O	2.08	0.54
1:A:233:VAL:O	1:A:237:VAL:HG13	2.09	0.53
1:B:298:ASN:HB3	1:B:330:ILE:HB	1.92	0.52
1:B:29:HIS:HB3	1:B:32:LYS:HG3	1.93	0.51
1:A:39:LYS:NZ	2:A:1039:PLP:O3	2.43	0.50
1:A:111:LEU:HD22	1:A:150:VAL:HG13	1.94	0.49
1:B:130:ILE:HD12	1:B:162:PHE:CE1	2.49	0.48
1:A:145:ASN:O	1:A:149:GLU:HG3	2.12	0.48
1:A:201:CYS:HG	1:A:215:CYS:HG	1.58	0.46
1:A:228:TYR:CE2	1:A:237:VAL:HG21	2.51	0.46
1:B:290:ARG:HA	1:B:310:VAL:HB	1.98	0.46
1:A:66:THR:CG2	1:B:7:ARG:HD3	2.46	0.45
1:A:222:ILE:HG12	1:A:227:TYR:HB2	1.98	0.44
1:B:166:PHE:HB2	1:B:200:HIS:O	2.18	0.44
1:A:232:TYR:CZ	1:A:236:LYS:HG3	2.52	0.44
1:B:133:ASP:OD2	1:B:137:GLY:HA2	2.18	0.43
1:B:362:ARG:CB	3:B:1385:SO4:O2	2.59	0.43
1:B:67:LEU:HB2	1:B:88:LEU:HD13	2.01	0.42
1:B:285:ALA:HB2	1:B:312:MET:HG3	2.02	0.42
1:A:228:TYR:HE2	1:A:237:VAL:HG21	1.85	0.42
1:B:368:TYR:O	1:B:374:ARG:HA	2.20	0.41
1:B:39:LYS:NZ	2:B:1039:PLP:O3	2.54	0.41
1:A:304:CYS:HB3	1:A:318:LYS:O	2.20	0.41
1:A:66:THR:HG22	1:B:7:ARG:HD3	2.02	0.41
1:A:131:LYS:HD3	1:A:166:PHE:CE2	2.56	0.40
1:A:209:LEU:HD11	1:A:233:VAL:HG22	2.03	0.40
1:B:168:HIS:HB3	3:B:1389:SO4:O4	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	350/382 (92%)	339 (97%)	11 (3%)	0	100	100
1	B	350/382 (92%)	335 (96%)	15 (4%)	0	100	100
All	All	700/764 (92%)	674 (96%)	26 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	310/330 (94%)	299 (96%)	11 (4%)	32	32
1	B	309/330 (94%)	299 (97%)	10 (3%)	34	36
All	All	619/660 (94%)	598 (97%)	21 (3%)	32	33

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	91	LYS
1	A	113	GLU
1	A	122	GLN
1	A	158	GLU
1	A	178	MET
1	A	179	THR
1	A	238	LYS
1	A	275	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	279	LEU
1	A	361	SER
1	A	377	THR
1	B	95	LYS
1	B	119	SER
1	B	179	THR
1	B	194	ILE
1	B	213	GLN
1	B	275	THR
1	B	300	ASN
1	B	361	SER
1	B	372	ASP
1	B	379	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	57	ASN
1	A	109	GLN
1	A	159	GLN
1	A	250	ASN
1	B	122	GLN
1	B	155	GLN
1	B	202	GLN
1	B	293	GLN
1	B	303	GLN
1	B	340	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

17 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	A	1387	-	4,4,4	0.37	0	6,6,6	0.56	0
3	SO4	B	1387	-	4,4,4	0.13	0	6,6,6	0.32	0
3	SO4	A	1385	-	4,4,4	0.26	0	6,6,6	0.23	0
3	SO4	B	1385	-	4,4,4	0.25	0	6,6,6	0.65	0
3	SO4	B	1386	-	4,4,4	0.26	0	6,6,6	0.35	0
3	SO4	B	1389	-	4,4,4	0.27	0	6,6,6	0.23	0
3	SO4	A	1384	-	4,4,4	0.39	0	6,6,6	0.42	0
3	SO4	A	1386	-	4,4,4	0.39	0	6,6,6	0.45	0
3	SO4	B	1388	-	4,4,4	0.24	0	6,6,6	0.30	0
3	SO4	A	1388	-	4,4,4	0.31	0	6,6,6	0.30	0
4	ACT	B	1390	-	3,3,3	0.62	0	3,3,3	1.51	1 (33%)
2	PLP	B	1039	1	15,15,16	1.93	3 (20%)	21,22,23	1.37	3 (14%)
4	ACT	A	1389	-	3,3,3	0.80	0	3,3,3	1.22	0
2	PLP	A	1039	1	15,15,16	1.99	3 (20%)	21,22,23	1.19	2 (9%)
3	SO4	A	1383	-	4,4,4	0.35	0	6,6,6	0.43	0
3	SO4	B	1384	-	4,4,4	0.28	0	6,6,6	0.29	0
3	SO4	B	1383	-	4,4,4	0.38	0	6,6,6	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	B	1039	1	-	0/6/6/8	0/1/1/1
2	PLP	A	1039	1	-	0/6/6/8	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1039	PLP	O3-C3	-5.46	1.24	1.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1039	PLP	O3-C3	-5.29	1.24	1.36
2	A	1039	PLP	C2-N1	3.11	1.39	1.33
2	B	1039	PLP	C2-N1	3.09	1.39	1.33
2	B	1039	PLP	C6-N1	2.32	1.39	1.34
2	A	1039	PLP	C6-N1	2.30	1.39	1.34

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1039	PLP	C5-C6-N1	-3.17	118.67	123.83
2	B	1039	PLP	O4P-C5A-C5	2.82	114.64	109.36
2	A	1039	PLP	C5-C6-N1	-2.41	119.91	123.83
2	B	1039	PLP	C6-C5-C4	2.29	119.97	118.10
2	A	1039	PLP	O4P-C5A-C5	2.16	113.41	109.36
4	B	1390	ACT	OXT-C-CH3	2.07	123.73	115.05

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

7 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1385	SO4	3	0
3	B	1389	SO4	1	0
3	A	1386	SO4	1	0
2	B	1039	PLP	2	0
4	A	1389	ACT	2	0
2	A	1039	PLP	1	0
3	B	1384	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	356/382 (93%)	-0.19	16 (4%) 38 43	8, 17, 40, 54	0
1	B	356/382 (93%)	-0.06	21 (5%) 28 32	10, 20, 41, 53	0
All	All	712/764 (93%)	-0.12	37 (5%) 33 36	8, 19, 41, 54	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	274	PRO	4.8
1	A	256	THR	3.8
1	A	178	MET	3.7
1	A	177	ASP	3.6
1	B	119	SER	3.2
1	B	256	THR	3.2
1	A	213	GLN	3.2
1	B	137	GLY	3.1
1	B	210	MET	3.1
1	A	211	ASP	3.1
1	A	122	GLN	2.9
1	A	181	GLU	2.9
1	B	179	THR	2.9
1	B	136	MET	2.8
1	B	139	LEU	2.8
1	B	257	LEU	2.6
1	A	2	SER	2.6
1	B	169	PHE	2.5
1	B	373	GLN	2.5
1	A	121	GLU	2.5
1	A	323	VAL	2.5
1	A	300	ASN	2.5
1	B	372	ASP	2.4
1	B	275	THR	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	2	SER	2.4
1	A	169	PHE	2.3
1	B	181	GLU	2.3
1	B	145	ASN	2.3
1	A	372	ASP	2.2
1	B	211	ASP	2.2
1	B	121	GLU	2.2
1	A	275	THR	2.2
1	A	324	LYS	2.1
1	B	300	ASN	2.1
1	B	212	CYS	2.1
1	A	302	HIS	2.0
1	B	168	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	B	1387	5/5	0.56	0.21	68,68,69,69	0
3	SO4	B	1389	5/5	0.72	0.19	87,87,88,88	0
3	SO4	B	1386	5/5	0.80	0.21	53,55,56,58	0
3	SO4	B	1384	5/5	0.82	0.16	78,78,79,79	0
3	SO4	A	1386	5/5	0.82	0.28	53,54,56,56	0
4	ACT	A	1389	4/4	0.84	0.17	46,46,46,47	0
3	SO4	A	1385	5/5	0.85	0.21	61,62,63,63	0
3	SO4	B	1385	5/5	0.85	0.28	45,49,51,53	0
3	SO4	B	1388	5/5	0.89	0.12	47,49,49,51	0
3	SO4	A	1388	5/5	0.90	0.13	49,50,50,51	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PLP	B	1039	15/16	0.91	0.14	17,36,40,41	0
3	SO4	A	1387	5/5	0.94	0.14	29,31,36,36	0
4	ACT	B	1390	4/4	0.94	0.10	28,28,28,28	0
2	PLP	A	1039	15/16	0.95	0.12	12,24,29,29	0
3	SO4	A	1383	5/5	0.97	0.06	21,23,24,24	0
3	SO4	A	1384	5/5	0.97	0.09	23,24,25,25	0
3	SO4	B	1383	5/5	0.98	0.08	19,21,25,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.