



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:42 PM UTC

PDB ID : 6A39 / pdb_00006a39
Title : The crystal structure of Mandelate oxidase Y128F with C4a-Malic acid-monooxide-FMN adduct
Authors : Li, T.L.; Lin, K.H.
Deposited on : 2018-06-15
Resolution : 1.89 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	377	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2769 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 4-hydroxymandelate oxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	330	2464	1549	446	458	11	0	1	0

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP O52792
A	-18	GLY	-	expression tag	UNP O52792
A	-17	SER	-	expression tag	UNP O52792
A	-16	SER	-	expression tag	UNP O52792
A	-15	HIS	-	expression tag	UNP O52792
A	-14	HIS	-	expression tag	UNP O52792
A	-13	HIS	-	expression tag	UNP O52792
A	-12	HIS	-	expression tag	UNP O52792
A	-11	HIS	-	expression tag	UNP O52792
A	-10	HIS	-	expression tag	UNP O52792
A	-9	SER	-	expression tag	UNP O52792
A	-8	SER	-	expression tag	UNP O52792
A	-7	GLY	-	expression tag	UNP O52792
A	-6	LEU	-	expression tag	UNP O52792
A	-5	VAL	-	expression tag	UNP O52792
A	-4	PRO	-	expression tag	UNP O52792
A	-3	ARG	-	expression tag	UNP O52792
A	-2	GLY	-	expression tag	UNP O52792
A	-1	SER	-	expression tag	UNP O52792
A	0	HIS	-	expression tag	UNP O52792
A	128	PHE	TYR	engineered mutation	UNP O52792

- Molecule 2 is ({E})-2-[[(4 {a} {S})-7,8-dimethyl-2,4-bis(oxidanylidene)-10-[(2 {S} ,3 {S} ,4 {S})-2,3,4-tris(oxidanyl)-5-phosphonooxy-pentyl]-5 {H} -benzo[g]pteridin-4 {a} -yl]oxy]-3-oxidanyl-but-2-enedioic acid (CCD ID: B8C) (formula: C₂₁H₂₅N₄O₁₅P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			41	21	4	15	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	264	Total O 264 264	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 4-hydroxymandelate oxidase



4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, α , β , γ	138.44Å 138.44Å 109.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.24 – 1.89 29.24 – 1.89	Depositor EDS
% Data completeness (in resolution range)	96.0 (29.24-1.89) 96.1 (29.24-1.89)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.55 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.190 , 0.235 0.199 , 0.243	Depositor DCC
R_{free} test set	2113 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	18.8	Xtriage
Anisotropy	0.055	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 49.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2769	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: B8C

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.65	29/2506 (1.2%)	1.71	28/3402 (0.8%)

All (29) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	302	ALA	C-O	10.65	1.36	1.23
1	A	253	GLY	C-O	9.83	1.36	1.24
1	A	260	ALA	C-O	7.01	1.32	1.23
1	A	142	ARG	C-O	6.77	1.31	1.24
1	A	351	ASN	C-O	6.75	1.32	1.23
1	A	58	VAL	C-O	-6.71	1.16	1.24
1	A	156	ASP	C-O	6.51	1.33	1.24
1	A	159	TRP	C-O	6.12	1.31	1.23
1	A	215	GLU	C-O	6.08	1.31	1.24
1	A	213	SER	CA-CB	-5.99	1.44	1.53
1	A	341	CYS	C-O	-5.92	1.17	1.24
1	A	280	VAL	C-O	-5.89	1.17	1.24
1	A	85	HIS	CE1-NE2	5.86	1.38	1.32
1	A	161	GLY	C-O	5.56	1.31	1.23
1	A	234	GLU	CD-OE1	5.55	1.35	1.25
1	A	222	ASP	C-O	-5.51	1.16	1.23
1	A	109	SER	CA-CB	-5.48	1.44	1.53
1	A	140	VAL	C-O	-5.43	1.18	1.24
1	A	63	ASP	C-O	5.41	1.30	1.24
1	A	100	GLY	CA-C	5.35	1.59	1.51
1	A	264	ILE	C-O	5.33	1.30	1.24
1	A	296	THR	CB-OG1	-5.27	1.35	1.43
1	A	130	LEU	C-O	5.25	1.30	1.23
1	A	227	LEU	C-O	5.19	1.30	1.24
1	A	82	ARG	C-O	-5.17	1.17	1.24
1	A	266	MET	C-O	-5.11	1.17	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	331	GLU	CD-OE2	-5.11	1.15	1.25
1	A	275	SER	CA-CB	-5.03	1.45	1.53
1	A	35	LEU	C-O	5.02	1.29	1.24

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	158	PRO	N-CA-C	-7.43	103.83	113.57
1	A	234	GLU	CB-CG-CD	7.15	124.75	112.60
1	A	251	ASN	N-CA-C	-6.45	105.37	112.72
1	A	136	SER	O-C-N	6.21	128.80	122.09
1	A	23	ASP	CA-C-O	-6.05	114.47	120.70
1	A	296	THR	OG1-CB-CG2	-5.95	97.39	109.30
1	A	297	ALA	CA-C-O	-5.85	113.49	120.10
1	A	145	ASP	CA-C-O	-5.81	114.39	120.55
1	A	224	PRO	N-CA-CB	5.80	108.49	103.27
1	A	138	ASP	N-CA-C	-5.77	104.97	112.23
1	A	265	GLU	CB-CG-CD	5.76	122.38	112.60
1	A	10	GLU	N-CA-C	-5.72	104.95	111.07
1	A	91	ALA	N-CA-C	-5.48	105.20	111.07
1	A	296	THR	CA-C-O	-5.40	114.83	120.55
1	A	136	SER	CA-C-O	-5.33	115.21	120.70
1	A	267	LEU	CA-C-O	-5.31	115.25	120.82
1	A	291	ASP	N-CA-C	-5.30	105.58	111.36
1	A	101	VAL	N-CA-CB	-5.23	106.01	111.61
1	A	166	ASP	CA-CB-CG	5.19	117.79	112.60
1	A	319	ASP	CA-C-N	5.18	125.69	119.94
1	A	319	ASP	C-N-CA	5.18	125.69	119.94
1	A	54	ASP	CA-CB-CG	-5.15	107.45	112.60
1	A	138	ASP	CA-CB-CG	5.13	117.73	112.60
1	A	49	PRO	CB-CA-C	5.09	117.62	110.60
1	A	337	GLY	CA-C-O	-5.08	115.52	120.75
1	A	106	CYS	CB-CA-C	-5.05	100.17	109.46
1	A	348	ARG	CA-C-O	-5.04	114.17	119.97
1	A	297	ALA	O-C-N	5.02	128.09	122.22

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2464	0	2482	40	0
2	A	41	0	0	12	0
3	A	264	0	0	12	1
All	All	2769	0	2482	47	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:402:B8C:CAQ	2:A:402:B8C:OBF	1.66	1.41
1:A:53:ARG:HD3	3:A:693:HOH:O	1.70	0.90
1:A:183:ASP:OD1	1:A:183:ASP:C	2.28	0.76
2:A:402:B8C:OBF	2:A:402:B8C:CAO	2.33	0.75
1:A:183:ASP:HB2	3:A:658:HOH:O	1.87	0.74
1:A:163:ARG:NH1	2:A:402:B8C:OBN	2.26	0.68
1:A:252:HIS:NE2	2:A:402:B8C:OBM	2.26	0.68
2:A:402:B8C:OBK	2:A:402:B8C:OBO	2.09	0.66
1:A:180:ALA:O	3:A:501:HOH:O	2.15	0.65
1:A:269:GLU:OE2	3:A:502:HOH:O	2.15	0.65
2:A:402:B8C:NAR	2:A:402:B8C:CBH	2.60	0.65
2:A:402:B8C:OBF	2:A:402:B8C:NAR	2.29	0.64
2:A:402:B8C:CAQ	2:A:402:B8C:CBH	2.75	0.64
1:A:132:ASP:N	3:A:506:HOH:O	2.24	0.64
1:A:252:HIS:CD2	2:A:402:B8C:OBM	2.52	0.62
1:A:163:ARG:CD	3:A:512:HOH:O	2.47	0.61
1:A:163:ARG:HD2	3:A:512:HOH:O	2.02	0.58
1:A:6:LEU:HD23	1:A:308:PRO:HB3	1.87	0.56
1:A:73:MET:HE1	1:A:328:LEU:HD13	1.89	0.55
1:A:130:LEU:HA	1:A:206:PHE:HA	1.91	0.53
1:A:255:ARG:HE	2:A:402:B8C:CBJ	2.23	0.52
1:A:255:ARG:NE	2:A:402:B8C:OBN	2.39	0.50
1:A:292:VAL:O	1:A:296:THR:HG23	2.11	0.50
1:A:43:GLU:HG2	3:A:736:HOH:O	2.11	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:67:ARG:HB2	1:A:122:ARG:CZ	2.43	0.49
1:A:131:ARG:HA	1:A:207:ALA:HB2	1.95	0.49
1:A:172:ALA:O	1:A:174:PRO:HD3	2.13	0.49
2:A:402:B8C:OBF	2:A:402:B8C:OBM	2.31	0.49
1:A:163:ARG:NH2	1:A:255:ARG:O	2.46	0.48
1:A:126:GLN:HA	1:A:152:VAL:O	2.13	0.48
1:A:130:LEU:HD23	1:A:206:PHE:CD1	2.49	0.48
1:A:80:TYR:CZ	1:A:173:LEU:HD11	2.49	0.47
1:A:73:MET:HE2	1:A:305:VAL:HG21	1.96	0.47
1:A:73:MET:HG2	1:A:74:ALA:N	2.28	0.46
1:A:130:LEU:HD23	1:A:206:PHE:CE1	2.50	0.46
1:A:158:PRO:HA	1:A:208:PRO:HA	1.97	0.45
1:A:322:ARG:NH1	1:A:326:GLU:HG3	2.32	0.45
1:A:279:GLU:OE1	3:A:505:HOH:O	2.21	0.45
1:A:283:ASP:OD1	1:A:283:ASP:C	2.61	0.44
1:A:348:ARG:HD3	3:A:731:HOH:O	2.17	0.44
1:A:114:GLU:O	3:A:503:HOH:O	2.20	0.43
1:A:6:LEU:CD1	1:A:6:LEU:N	2.81	0.43
1:A:264:ILE:HD12	1:A:264:ILE:HG23	1.74	0.43
1:A:322:ARG:HD2	1:A:322:ARG:HA	1.89	0.42
1:A:20:GLU:HG3	1:A:171:PHE:HD1	1.83	0.42
1:A:61:GLU:OE1	1:A:68:ARG:NH1	2.54	0.41
1:A:129:TRP:O	3:A:504:HOH:O	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:533:HOH:O	3:A:747:HOH:O[6_555]	2.03	0.17

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	327/377 (87%)	314 (96%)	10 (3%)	3 (1%)	14 6

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	19	GLY
1	A	174	PRO
1	A	316	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	246/281 (88%)	239 (97%)	7 (3%)	38 32

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	ARG
1	A	20	GLU
1	A	163	ARG
1	A	174	PRO
1	A	183	ASP
1	A	237	ARG
1	A	356	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	B8C	A	402	-	38,43,43	4.70	18 (47%)	47,66,66	3.19	21 (44%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	B8C	A	402	-	-	13/24/68/68	0/3/3/3

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	402	B8C	CAQ-CAO	-23.49	1.34	1.54
2	A	402	B8C	CAV-CAU	-6.15	1.42	1.53
2	A	402	B8C	CBI-CBH	6.08	1.43	1.34
2	A	402	B8C	CAH-CAG	-5.56	1.40	1.51
2	A	402	B8C	CAA-CAB	-4.90	1.41	1.51
2	A	402	B8C	CAD-NAR	-4.27	1.32	1.39
2	A	402	B8C	CAS-CAT	-4.21	1.46	1.52
2	A	402	B8C	CAW-CAV	-3.94	1.46	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	402	B8C	PAY-OAX	3.78	1.72	1.60
2	A	402	B8C	OAP-CAO	3.43	1.28	1.22
2	A	402	B8C	OBN-CBJ	-3.09	1.22	1.30
2	A	402	B8C	OBE-CAT	-2.91	1.37	1.43
2	A	402	B8C	PAY-OAZ	-2.89	1.44	1.54
2	A	402	B8C	CAB-CAG	-2.73	1.34	1.40
2	A	402	B8C	CAT-CAU	-2.35	1.49	1.53
2	A	402	B8C	CAO-NAN	2.26	1.40	1.37
2	A	402	B8C	CAJ-NAK	2.19	1.37	1.31
2	A	402	B8C	CAC-CAB	2.17	1.42	1.39

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	402	B8C	OBO-CBI-CBJ	12.13	125.54	114.12
2	A	402	B8C	OBF-CBH-CBG	9.65	131.97	116.12
2	A	402	B8C	OBC-CAV-CAW	-5.41	98.06	109.99
2	A	402	B8C	CAT-CAU-CAV	4.88	121.68	113.57
2	A	402	B8C	OBD-CAU-CAV	4.46	119.06	108.93
2	A	402	B8C	OAM-CAL-NAK	4.38	129.07	121.80
2	A	402	B8C	OAX-CAW-CAV	-3.64	99.64	109.36
2	A	402	B8C	OBE-CAT-CAS	3.61	125.05	110.20
2	A	402	B8C	OBC-CAV-CAU	3.35	117.09	109.25
2	A	402	B8C	CAF-CAE-CAD	-3.20	115.48	119.95
2	A	402	B8C	CAC-CAD-CAE	3.09	123.18	119.80
2	A	402	B8C	OBD-CAU-CAT	2.99	115.72	108.93
2	A	402	B8C	CAD-CAE-NAI	2.97	121.37	118.01
2	A	402	B8C	CAE-NAI-CAJ	2.61	123.79	119.73
2	A	402	B8C	NAN-CAL-NAK	-2.60	113.97	119.50
2	A	402	B8C	OBB-PAY-OAX	2.49	113.16	106.67
2	A	402	B8C	CAE-CAD-NAR	-2.21	116.58	119.18
2	A	402	B8C	CAW-CAV-CAU	2.21	116.39	112.22
2	A	402	B8C	OBB-PAY-OAZ	2.18	115.96	107.80
2	A	402	B8C	CAE-CAF-CAG	2.14	123.52	119.22
2	A	402	B8C	OBK-CBG-OBL	-2.04	119.04	123.90

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	402	B8C	CAW-OAX-PAY-OAZ
2	A	402	B8C	OBC-CAV-CAW-OAX

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	402	B8C	CAU-CAV-CAW-OAX
2	A	402	B8C	OBD-CAU-CAV-CAW
2	A	402	B8C	CAT-CAU-CAV-CAW
2	A	402	B8C	OBD-CAU-CAV-OBC
2	A	402	B8C	OBE-CAT-CAU-CAV
2	A	402	B8C	CAS-CAT-CAU-CAV
2	A	402	B8C	CAT-CAU-CAV-OBC
2	A	402	B8C	CAV-CAW-OAX-PAY
2	A	402	B8C	CAW-OAX-PAY-OBA
2	A	402	B8C	OBO-CBI-CBJ-OBN
2	A	402	B8C	CAW-OAX-PAY-OB

There are no ring outliers.

1 monomer is involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	402	B8C	12	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	330/377 (87%)	0.57	48 (14%) 6 6	9, 23, 66, 108	1 (0%)

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	6	LEU	9.0
1	A	7	ALA	6.4
1	A	356	VAL	5.6
1	A	5	SER	5.4
1	A	176	TRP	5.2
1	A	8	ASP	4.6
1	A	164	LEU	4.4
1	A	167	MET	4.4
1	A	171	PHE	3.7
1	A	16	VAL	3.7
1	A	160	MET	3.4
1	A	206	PHE	3.3
1	A	177	VAL	3.2
1	A	178	THR	3.1
1	A	163	ARG	3.1
1	A	21	ILE	3.1
1	A	130	LEU	3.1
1	A	162	ARG	3.1
1	A	168	ARG	3.0
1	A	207	ALA	3.0
1	A	170	GLY	3.0
1	A	183	ASP	2.9
1	A	9	LEU	2.9
1	A	173	LEU	2.9
1	A	12	ALA	2.9
1	A	11	ARG	2.9
1	A	132	ASP	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	169	ASN	2.7
1	A	33	ALA	2.6
1	A	17	LEU	2.6
1	A	319	ASP	2.6
1	A	158	PRO	2.6
1	A	131	ARG	2.6
1	A	20	GLU	2.6
1	A	166	ASP	2.5
1	A	19	GLY	2.5
1	A	24	PHE	2.5
1	A	138	ASP	2.4
1	A	182	PHE	2.4
1	A	14	ARG	2.3
1	A	139	LEU	2.3
1	A	156	ASP	2.2
1	A	22	PHE	2.2
1	A	314	ALA	2.1
1	A	315	ALA	2.1
1	A	159	TRP	2.1
1	A	316	ALA	2.0
1	A	175	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	B8C	A	402	41/41	0.95	0.11	14,28,70,103	41

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.