



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2026 – 07:44 PM UTC

PDB ID : 7A3I / pdb_00007a3i
Title : Crystal structure of DPP8 in complex with a 4-oxo-b-lactam based inhibitor, LMC375
Authors : Ross, B.H.; Huber, R.
Deposited on : 2020-08-18
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	898	5% 87% 10%
1	B	898	5% 87% 10%
1	C	898	6% 88% 10%

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 19856 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dipeptidyl peptidase 8.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	807	Total	C	N	O	S	0	0	0
			6554	4221	1090	1216	27			
1	B	808	Total	C	N	O	S	0	0	0
			6558	4224	1090	1217	27			
1	C	808	Total	C	N	O	S	0	0	0
			6561	4226	1091	1217	27			

- Molecule 2 is trimethylamine oxide (CCD ID: TMO) (formula: C_3H_9NO).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			5	3	1	1		
2	B	1	Total	C	N	O	0	0
			5	3	1	1		
2	C	1	Total	C	N	O	0	0
			5	3	1	1		

- Molecule 3 is {N}-[3,5-bis[[4-[(4-bromophenyl)methyl]piperazin-1-yl]methyl]phenyl]-2-ethyl-2-methanoyl-butanamide (CCD ID: QXH) (formula: C₃₇H₄₇Br₂N₅O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	Br	C	N	O	0	0
			46	2	37	5	2		
3	B	1	Total	Br	C	N	O	0	0
			46	2	37	5	2		
3	C	1	Total	Br	C	N	O	0	0
			46	2	37	5	2		

- Molecule 4 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Na	0	0
			1	1		
4	B	1	Total	Na	0	0
			1	1		
4	C	1	Total	Na	0	0
			1	1		

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Cl	0	0
			1	1		
5	B	1	Total	Cl	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	1	Total	Cl	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	12	Total	O	0	0
			12	12		
6	B	7	Total	O	0	0
			7	7		
6	C	5	Total	O	0	0
			5	5		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Dipeptidyl peptidase 8



- Molecule 1: Dipeptidyl peptidase 8



- Molecule 1: Dipeptidyl peptidase 8



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	163.65Å 252.32Å 259.96Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.33 – 2.80 49.33 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.9 (49.33-2.80) 99.9 (49.33-2.80)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.07 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.8.0155	Depositor
R, R_{free}	0.217 , 0.241 (Not available) , 0.237	Depositor DCC
R_{free} test set	6579 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	66.6	Xtriage
Anisotropy	0.290	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 22.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	19856	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TMO, NA, CL, QXH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.51	0/6739	0.71	0/9142
1	B	0.51	0/6743	0.72	0/9148
1	C	0.51	0/6746	0.71	2/9152 (0.0%)
All	All	0.51	0/20228	0.71	2/27442 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	193	GLY	CA-C-N	5.08	125.11	119.32
1	C	193	GLY	C-N-CA	5.08	125.11	119.32

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6554	0	6363	8	0
1	B	6558	0	6369	8	0
1	C	6561	0	6372	8	0
2	A	5	0	9	0	0
2	B	5	0	9	0	0
2	C	5	0	9	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	46	0	0	2	0
3	B	46	0	0	0	0
3	C	46	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
6	A	12	0	0	0	0
6	B	7	0	0	0	0
6	C	5	0	0	0	0
All	All	19856	0	19131	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:118:ILE:HD12	1:C:599:LEU:CD2	2.40	0.52
1:A:880:LEU:HD23	3:A:902:QXH:BR1	2.66	0.50
1:B:118:ILE:HD12	1:B:599:LEU:CD2	2.42	0.49
1:C:627:LEU:HD13	1:C:630:TYR:HB3	1.96	0.48
1:C:651:TYR:HB2	1:C:699:VAL:HB	1.95	0.47
1:C:273:LEU:HD13	1:C:279:ARG:NH2	2.30	0.47
1:C:376:GLU:HG3	1:C:397:ARG:HB2	1.95	0.47
1:C:787:TYR:CZ	2:C:901:TMO:HABA	2.50	0.47
1:B:273:LEU:HD13	1:B:279:ARG:NH2	2.31	0.46
1:C:118:ILE:HD12	1:C:599:LEU:HD22	1.97	0.46
1:B:376:GLU:HG3	1:B:397:ARG:HB2	1.98	0.46
1:B:189:VAL:HG13	1:B:204:PRO:HA	1.97	0.45
1:A:302:LEU:HD22	1:A:390:ALA:HB1	1.99	0.44
1:B:763:MET:HE1	1:B:808:LEU:O	2.18	0.43
1:A:78:LYS:HB2	1:A:104:MET:HE3	2.01	0.43
1:A:118:ILE:HD12	1:A:599:LEU:CD2	2.48	0.43
1:A:273:LEU:HD13	1:A:279:ARG:NH2	2.34	0.43
1:A:259:MET:HE1	1:A:329:PHE:HA	2.01	0.42
1:B:651:TYR:HB2	1:B:699:VAL:HB	2.01	0.42
1:C:756:TYR:HB2	2:C:901:TMO:HAD	2.02	0.41
1:A:860:TYR:HB3	3:A:902:QXH:C42	2.49	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:782:THR:HA	1:A:811:VAL:HG22	2.01	0.41
1:B:302:LEU:HD22	1:B:390:ALA:HB1	2.02	0.41
1:B:434:THR:HG21	1:B:489:SER:HB2	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	799/898 (89%)	776 (97%)	21 (3%)	2 (0%)	36	66
1	B	800/898 (89%)	778 (97%)	20 (2%)	2 (0%)	36	66
1	C	800/898 (89%)	770 (96%)	30 (4%)	0	100	100
All	All	2399/2694 (89%)	2324 (97%)	71 (3%)	4 (0%)	43	72

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	445	ILE
1	A	561	GLY
1	B	445	ILE
1	B	561	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	714/795 (90%)	704 (99%)	10 (1%)	59	85
1	B	715/795 (90%)	705 (99%)	10 (1%)	59	85
1	C	715/795 (90%)	707 (99%)	8 (1%)	65	87
All	All	2144/2385 (90%)	2116 (99%)	28 (1%)	61	86

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	70	LYS
1	A	214	ILE
1	A	350	ILE
1	A	591	GLN
1	A	592	LYS
1	A	614	LYS
1	A	627	LEU
1	A	656	LEU
1	A	718	PHE
1	A	794	ARG
1	B	131	SER
1	B	170	ASP
1	B	189	VAL
1	B	260	GLU
1	B	386	GLU
1	B	461	GLU
1	B	614	LYS
1	B	655	ASP
1	B	704	ASN
1	B	718	PHE
1	C	170	ASP
1	C	214	ILE
1	C	259	MET
1	C	592	LYS
1	C	614	LYS
1	C	627	LEU
1	C	718	PHE
1	C	846	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (31) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	60	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	72	HIS
1	A	95	HIS
1	A	123	ASN
1	A	173	GLN
1	A	199	GLN
1	A	423	GLN
1	A	579	GLN
1	A	591	GLN
1	A	801	GLN
1	B	123	ASN
1	B	173	GLN
1	B	195	GLN
1	B	253	HIS
1	B	423	GLN
1	B	579	GLN
1	B	654	HIS
1	B	657	GLN
1	B	858	GLN
1	C	123	ASN
1	C	173	GLN
1	C	195	GLN
1	C	199	GLN
1	C	232	HIS
1	C	338	ASN
1	C	550	HIS
1	C	559	ASN
1	C	579	GLN
1	C	801	GLN
1	C	837	HIS
1	C	882	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 12 ligands modelled in this entry, 6 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	TMO	B	901	-	4,4,4	6.36	1 (25%)	6,6,6	0.33	0
3	QXH	C	902	1	48,50,50	0.64	0	63,69,69	1.22	5 (7%)
2	TMO	C	901	-	4,4,4	6.33	1 (25%)	6,6,6	0.28	0
2	TMO	A	901	-	4,4,4	6.33	1 (25%)	6,6,6	0.26	0
3	QXH	B	902	1	48,50,50	0.62	0	63,69,69	1.21	4 (6%)
3	QXH	A	902	1	48,50,50	0.64	0	63,69,69	1.18	6 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	QXH	C	902	1	-	2/32/55/55	0/5/5/5
3	QXH	B	902	1	-	3/32/55/55	0/5/5/5
3	QXH	A	902	1	-	5/32/55/55	0/5/5/5

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	901	TMO	OAE-NAC	-12.60	1.25	1.42
2	A	901	TMO	OAE-NAC	-12.58	1.25	1.42
2	C	901	TMO	OAE-NAC	-12.56	1.25	1.42

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	902	QXH	C1-C6-N9	4.94	122.20	115.68
3	B	902	QXH	C1-C6-N9	4.52	121.64	115.68
3	A	902	QXH	C1-C6-N9	4.04	121.01	115.68
3	C	902	QXH	BR2-C44-C45	3.09	123.51	119.30
3	A	902	QXH	BR2-C44-C45	2.77	123.07	119.30
3	B	902	QXH	BR1-C26-C27	2.75	123.05	119.30
3	B	902	QXH	C2-C1-C6	-2.53	101.82	108.70
3	A	902	QXH	C2-C1-C6	-2.44	102.05	108.70
3	C	902	QXH	C3-C2-C1	2.37	119.34	114.69
3	A	902	QXH	BR1-C26-C27	2.27	122.39	119.30
3	B	902	QXH	C39-C38-N37	-2.21	106.20	110.65
3	A	902	QXH	C9-C8-N9	2.09	126.93	120.13
3	A	902	QXH	C21-C20-N19	-2.08	106.45	110.65
3	C	902	QXH	C2-C1-C6	-2.04	103.14	108.70
3	C	902	QXH	BR1-C26-C25	2.00	122.03	119.30

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	902	QXH	C41-C40-N37-C38
3	A	902	QXH	C4-C1-C6-O10
3	A	902	QXH	C2-C1-C4-C5
3	A	902	QXH	C41-C40-N37-C36
3	A	902	QXH	C4-C1-C6-N9
3	B	902	QXH	C4-C1-C6-N9
3	B	902	QXH	C4-C1-C6-O10
3	C	902	QXH	C4-C1-C6-N9
3	C	902	QXH	C4-C1-C6-O10
3	B	902	QXH	N19-C22-C23-C24


There are no ring outliers.


2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	901	TMO	2	0
3	A	902	QXH	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	807/898 (89%)	0.37	44 (5%)	30	23	45, 65, 109, 163	0
1	B	808/898 (89%)	0.38	45 (5%)	30	23	42, 63, 108, 161	0
1	C	808/898 (89%)	0.47	51 (6%)	26	19	47, 68, 113, 148	0
All	All	2423/2694 (89%)	0.40	140 (5%)	29	22	42, 65, 111, 163	0

All (140) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	870	PRO	8.0
1	B	316	VAL	7.6
1	A	870	PRO	6.5
1	A	259	MET	5.5
1	B	164	VAL	5.4
1	C	870	PRO	5.1
1	B	48	LEU	5.0
1	B	70	LYS	4.8
1	B	139	PHE	4.8
1	A	627	LEU	4.6
1	B	419	VAL	4.5
1	C	164	VAL	4.5
1	C	259	MET	4.5
1	A	138	LEU	4.5
1	B	76	MET	4.4
1	B	872	SER	4.4
1	C	419	VAL	4.4
1	A	628	PRO	4.3
1	B	256	LEU	4.3
1	A	897	VAL	4.3
1	A	831	PHE	4.3
1	C	316	VAL	4.2
1	C	628	PRO	4.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	164	VAL	4.1
1	C	897	VAL	4.1
1	B	259	MET	4.1
1	B	832	LEU	4.0
1	A	76	MET	4.0
1	C	831	PHE	4.0
1	C	256	LEU	4.0
1	B	257	ALA	4.0
1	C	73	GLY	3.9
1	A	75	MET	3.9
1	C	72	HIS	3.9
1	A	260	GLU	3.9
1	B	71	TYR	3.9
1	C	257	ALA	3.9
1	B	74	TYR	3.8
1	A	832	LEU	3.8
1	C	138	LEU	3.8
1	A	315	HIS	3.7
1	A	257	ALA	3.7
1	B	75	MET	3.6
1	A	73	GLY	3.6
1	C	460	HIS	3.6
1	B	355	ARG	3.5
1	C	129	MET	3.5
1	B	831	PHE	3.5
1	B	628	PRO	3.5
1	A	140	GLN	3.5
1	C	76	MET	3.4
1	C	70	LYS	3.4
1	B	862	GLN	3.4
1	C	627	LEU	3.3
1	A	872	SER	3.3
1	A	256	LEU	3.2
1	C	48	LEU	3.2
1	B	73	GLY	3.2
1	B	871	GLU	3.2
1	C	107	GLU	3.2
1	B	420	MET	3.2
1	B	460	HIS	3.2
1	A	419	VAL	3.2
1	C	106	GLY	3.1
1	B	873	GLY	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	524	ARG	3.0
1	B	897	VAL	3.0
1	A	48	LEU	3.0
1	A	655	ASP	3.0
1	C	140	GLN	3.0
1	A	420	MET	3.0
1	B	659	GLY	3.0
1	C	78	LYS	2.9
1	C	74	TYR	2.9
1	B	328	SER	2.9
1	C	75	MET	2.9
1	B	629	ASP	2.9
1	B	261	GLU	2.9
1	A	70	LYS	2.8
1	A	129	MET	2.8
1	C	420	MET	2.8
1	A	71	TYR	2.7
1	A	72	HIS	2.7
1	B	627	LEU	2.7
1	C	871	GLU	2.7
1	A	460	HIS	2.6
1	C	261	GLU	2.6
1	B	830	GLY	2.6
1	A	74	TYR	2.6
1	A	261	GLU	2.6
1	B	72	HIS	2.6
1	A	487	LYS	2.6
1	A	314	ILE	2.5
1	C	369	GLU	2.5
1	B	422	ARG	2.5
1	A	262	ASP	2.5
1	A	629	ASP	2.4
1	A	862	GLN	2.4
1	B	490	LYS	2.4
1	B	315	HIS	2.4
1	C	355	ARG	2.4
1	C	315	HIS	2.4
1	C	626	PRO	2.4
1	B	216	MET	2.4
1	C	258	ASN	2.4
1	C	872	SER	2.3
1	C	895	LEU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	891	ARG	2.3
1	A	873	GLY	2.3
1	A	501	PRO	2.3
1	B	354	GLY	2.3
1	A	871	GLU	2.3
1	C	461	GLU	2.2
1	A	78	LYS	2.2
1	B	834	GLU	2.2
1	C	293	PRO	2.2
1	B	260	GLU	2.2
1	C	314	ILE	2.2
1	A	835	ASN	2.2
1	A	354	GLY	2.2
1	C	91	PRO	2.1
1	C	658	PRO	2.1
1	C	71	TYR	2.1
1	C	624	ALA	2.1
1	C	244	ARG	2.1
1	C	592	LYS	2.1
1	C	835	ASN	2.1
1	B	106	GLY	2.1
1	A	139	PHE	2.1
1	B	891	ARG	2.1
1	A	895	LEU	2.0
1	A	561	GLY	2.0
1	B	110	GLU	2.0
1	B	660	LYS	2.0
1	B	195	GLN	2.0
1	C	832	LEU	2.0
1	C	329	PHE	2.0
1	A	110	GLU	2.0
1	C	278	ASP	2.0
1	C	373	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NA	A	903	1/1	0.71	0.36	77,77,77,77	0
2	TMO	B	901	5/5	0.81	0.23	56,57,58,60	0
3	QXH	B	902	46/46	0.85	0.22	66,88,158,173	0
4	NA	C	903	1/1	0.85	0.21	77,77,77,77	0
3	QXH	A	902	46/46	0.86	0.23	64,90,151,159	0
3	QXH	C	902	46/46	0.86	0.23	73,91,167,183	0
2	TMO	C	901	5/5	0.87	0.24	70,70,70,71	0
4	NA	B	903	1/1	0.90	0.15	70,70,70,70	0
2	TMO	A	901	5/5	0.90	0.21	64,66,67,67	0
5	CL	C	904	1/1	0.91	0.13	82,82,82,82	0
5	CL	A	904	1/1	0.97	0.12	90,90,90,90	0
5	CL	B	904	1/1	0.98	0.17	85,85,85,85	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around QXH B 902:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around QXH A 902:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around QXH C 902:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.