



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:17 PM UTC

PDB ID : 1A4I / pdb_00001a4i
Title : HUMAN TETRAHYDROFOLATE DEHYDROGENASE / CYCLOHYDRO-
LASE
Authors : Allaire, M.; Li, Y.; Mackenzie, R.E.; Cygler, M.
Deposited on : 1998-01-30
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	301	4% 77% 15% • 5%
1	B	301	5% 79% 18% ••

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 4899 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called METHYLENETETRAHYDROFOLATE DEHYDROGENASE / METHENYLTETRAHYDROFOLATE CYCLOHYDROLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	285	Total	C	N	O	S	16	0	0
			2159	1357	374	417	11			
1	B	295	Total	C	N	O	S	4	0	0
			2237	1404	390	432	11			

- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NDP) (formula: $C_{21}H_{30}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	179	Total 179	O 179	0	0
3	B	228	Total 228	O 228	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: METHYLENETETRAHYDROFOLATE DEHYDROGENASE / METHENYLTE TRAHYDROFOLATE CYCLOHYDROLASE



• Molecule 1: METHYLENETETRAHYDROFOLATE DEHYDROGENASE / METHENYLTE TRAHYDROFOLATE CYCLOHYDROLASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	67.52Å 135.84Å 61.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.50 20.00 – 1.50	Depositor EDS
% Data completeness (in resolution range)	90.0 (20.00-1.50) 83.0 (20.00-1.50)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.38 (at 1.50Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.199 , 0.235 0.188 , 0.226	Depositor DCC
R_{free} test set	3818 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	16.5	Xtriage
Anisotropy	0.123	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 36.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4899	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.99	1/2191 (0.0%)	1.81	35/2973 (1.2%)
1	B	1.00	0/2272	1.83	47/3084 (1.5%)
All	All	1.00	1/4463 (0.0%)	1.82	82/6057 (1.4%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	196	HIS	CE1-NE2	5.27	1.37	1.32

All (82) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	251	LYS	CA-CB-CG	-20.54	73.03	114.10
1	B	285	GLN	OE1-CD-NE2	14.29	136.89	122.60
1	A	251	LYS	CB-CG-CD	-11.12	85.72	111.30
1	B	17	ARG	CD-NE-CZ	9.86	138.21	124.40
1	A	139	ASP	CA-CB-CG	9.58	122.18	112.60
1	B	296	GLU	CB-CG-CD	9.31	128.43	112.60
1	B	116	ASN	CA-CB-CG	-8.97	103.63	112.60
1	A	141	ASN	CA-C-N	8.83	137.11	122.54
1	A	141	ASN	C-N-CA	8.83	137.11	122.54
1	A	142	ASP	CB-CA-C	-8.75	95.47	110.09
1	B	289	GLU	O-C-N	8.65	131.29	122.12
1	A	142	ASP	CA-CB-CG	8.54	121.14	112.60
1	B	26	GLN	OE1-CD-NE2	8.23	130.83	122.60
1	A	112	GLU	CB-CG-CD	8.20	126.54	112.60
1	B	131	ASN	CA-CB-CG	-7.91	104.69	112.60
1	A	218	GLN	CA-C-O	7.75	126.99	119.55
1	B	167	HIS	CA-CB-CG	-7.75	106.05	113.80
1	B	290	SER	O-C-N	7.70	130.93	122.15

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	142	ASP	N-CA-C	7.65	122.81	113.17
1	B	48	ASP	CA-CB-CG	7.58	120.18	112.60
1	B	64	GLY	CA-C-O	7.45	127.00	118.96
1	A	107	ASN	CA-CB-CG	-7.17	105.42	112.60
1	B	287	THR	O-C-N	6.98	130.11	122.15
1	A	279	THR	O-C-N	6.82	129.35	122.12
1	A	279	THR	CA-C-O	-6.67	113.48	120.55
1	B	289	GLU	CA-C-O	-6.54	113.62	120.55
1	B	45	ASN	O-C-N	6.46	128.58	122.18
1	B	207	ASN	OD1-CG-ND2	6.41	129.01	122.60
1	B	95	HIS	CA-CB-CG	-6.31	107.49	113.80
1	B	166	ARG	NE-CZ-NH1	-6.31	115.19	121.50
1	B	64	GLY	O-C-N	-6.21	115.61	122.50
1	B	271	VAL	CA-C-O	-6.20	116.72	119.94
1	B	285	GLN	CB-CG-CD	-6.18	102.09	112.60
1	B	129	SER	O-C-N	-6.10	115.66	122.12
1	B	23	GLN	O-C-N	6.02	128.50	122.12
1	A	225	GLU	CA-CB-CG	5.96	126.02	114.10
1	A	64	GLY	CA-C-O	5.93	125.36	118.96
1	A	91	ASP	CA-C-N	5.92	128.80	120.28
1	A	91	ASP	C-N-CA	5.92	128.80	120.28
1	B	66	LYS	CA-C-N	-5.88	114.69	122.99
1	B	66	LYS	C-N-CA	-5.88	114.69	122.99
1	B	37	ARG	O-C-N	5.84	130.48	123.36
1	B	126	GLY	CA-C-N	-5.75	113.05	122.54
1	B	126	GLY	C-N-CA	-5.75	113.05	122.54
1	A	204	GLU	O-C-N	5.73	129.32	122.27
1	A	95	HIS	CA-CB-CG	-5.71	108.08	113.80
1	A	116	ASN	CA-CB-CG	-5.69	106.91	112.60
1	B	42	GLN	CA-C-N	-5.67	116.16	123.19
1	B	42	GLN	C-N-CA	-5.67	116.16	123.19
1	A	100	GLN	OE1-CD-NE2	-5.64	116.96	122.60
1	B	65	ILE	O-C-N	5.50	128.92	122.81
1	A	286	SER	O-C-N	5.50	127.95	122.12
1	A	258	TYR	N-CA-C	5.49	116.94	111.07
1	B	189	ASN	CA-CB-CG	-5.46	107.14	112.60
1	A	97	PHE	O-C-N	5.39	129.30	123.10
1	A	98	LEU	O-C-N	5.37	129.37	123.25
1	A	182	HIS	ND1-CE1-NE2	-5.37	103.03	108.40
1	B	191	THR	CA-C-O	-5.36	114.43	120.43
1	A	35	THR	O-C-N	5.33	127.44	121.32
1	B	194	THR	O-C-N	5.33	129.57	123.29

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	74	ARG	NE-CZ-NH2	5.32	123.98	119.20
1	B	16	ILE	O-C-N	5.30	127.11	121.91
1	B	208	LYS	CA-C-O	5.29	125.47	119.18
1	A	149	PRO	O-C-N	-5.28	115.89	122.23
1	A	194	THR	O-C-N	5.27	129.58	123.30
1	A	290	SER	CA-C-O	-5.25	114.98	120.55
1	B	142	ASP	O-C-N	-5.25	116.37	122.09
1	B	31	VAL	O-C-N	-5.23	115.13	121.10
1	B	61	GLU	CA-C-O	-5.22	113.96	119.97
1	A	119	ALA	CA-C-O	5.22	124.65	119.75
1	B	30	GLN	OE1-CD-NE2	-5.22	117.38	122.60
1	B	45	ASN	CA-C-O	-5.20	116.34	121.02
1	B	20	LEU	O-C-N	5.19	127.62	122.12
1	B	103	LEU	CA-C-O	-5.15	116.09	121.55
1	A	211	ILE	N-CA-C	-5.12	100.83	108.46
1	B	70	ILE	CA-C-O	-5.12	115.05	120.53
1	B	96	GLY	O-C-N	5.11	127.37	123.23
1	A	124	VAL	CA-C-O	-5.09	112.86	119.53
1	A	219	PRO	N-CA-C	5.08	119.17	110.95
1	B	166	ARG	NH1-CZ-NH2	5.08	125.90	119.30
1	B	290	SER	CA-C-O	-5.04	115.08	120.42
1	B	147	CYS	CA-C-O	-5.03	115.22	120.55

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2159	0	2208	20	0
1	B	2237	0	2288	15	0
2	A	48	0	26	0	0
2	B	48	0	26	1	0
3	A	179	0	0	2	1
3	B	228	0	0	3	1
All	All	4899	0	4548	35	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2:ALA:HB3	1:A:3:PRO:HD3	1.61	0.81
1:A:37:ARG:H	1:A:95:HIS:HD2	1.33	0.75
1:B:166:ARG:HH12	1:B:232:ILE:HD12	1.54	0.72
1:A:85:ILE:HD11	1:A:114:VAL:HG13	1.73	0.69
1:A:2:ALA:HB3	1:A:3:PRO:CD	2.26	0.66
1:B:238:ILE:HD11	2:B:302:NDP:N7N	2.11	0.65
1:A:158:GLU:HG2	3:A:464:HOH:O	1.97	0.64
1:A:85:ILE:CD1	1:A:114:VAL:HG13	2.32	0.59
1:B:131:ASN:ND2	3:B:470:HOH:O	2.35	0.58
1:A:27:LEU:HD11	1:A:295:LEU:HD12	1.85	0.57
1:B:166:ARG:HH12	1:B:232:ILE:CD1	2.16	0.57
1:A:74:ARG:NE	1:A:104:ASP:OD1	2.38	0.56
1:A:112:GLU:HG2	1:A:116:ASN:HD22	1.73	0.54
1:A:74:ARG:HE	1:A:104:ASP:CG	2.14	0.54
1:A:37:ARG:H	1:A:95:HIS:CD2	2.20	0.53
1:B:166:ARG:NH1	1:B:232:ILE:HD12	2.25	0.51
1:B:37:ARG:CD	1:B:66:LYS:HD2	2.41	0.50
1:A:121:GLU:H	1:A:121:GLU:CD	2.20	0.49
1:B:58:LYS:HE2	3:B:522:HOH:O	2.12	0.49
1:A:37:ARG:N	1:A:95:HIS:HD2	2.06	0.47
1:B:37:ARG:HD2	1:B:66:LYS:HD2	1.96	0.47
1:B:82:MET:HE1	1:B:117:ALA:HB2	1.97	0.46
1:B:85:ILE:CD1	1:B:114:VAL:HG13	2.45	0.46
1:A:2:ALA:CB	1:A:3:PRO:CD	2.92	0.46
1:A:36:PRO:HA	1:A:95:HIS:CD2	2.53	0.44
1:B:140:LEU:HD11	1:B:184:LEU:HD23	1.99	0.44
1:B:162:PRO:O	1:B:166:ARG:NE	2.51	0.43
3:A:430:HOH:O	1:B:186:LEU:HD13	2.18	0.42
1:B:27:LEU:HD11	1:B:292:LYS:HE2	2.01	0.42
1:B:176:ILE:HG12	3:B:351:HOH:O	2.18	0.42
1:A:37:ARG:HD2	1:A:66:LYS:HE2	2.02	0.41
1:A:216:THR:HG21	1:A:221:MET:HE1	2.03	0.41
1:A:148:THR:HB	1:A:149:PRO:HD3	2.02	0.40
1:A:223:LYS:HA	1:A:223:LYS:HD3	1.85	0.40
1:A:224:GLY:O	1:A:264:ARG:HD2	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:389:HOH:O	3:B:502:HOH:O[3_555]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	281/301 (93%)	278 (99%)	3 (1%)	0	100	100
1	B	293/301 (97%)	287 (98%)	6 (2%)	0	100	100
All	All	574/602 (95%)	565 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	239/253 (94%)	232 (97%)	7 (3%)	37	10
1	B	248/253 (98%)	243 (98%)	5 (2%)	48	20
All	All	487/506 (96%)	475 (98%)	12 (2%)	42	14

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	LYS
1	A	92	SER
1	A	140	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	198	LYS
1	A	221	MET
1	A	225	GLU
1	A	259	ASP
1	B	54	ASN
1	B	74	ARG
1	B	90	GLU
1	B	121	GLU
1	B	247	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	15	GLN
1	A	22	ASN
1	A	95	HIS
1	A	188	ASN
1	B	22	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NDP	A	302	-	51,52,52	1.62	6 (11%)	71,80,80	1.42	12 (16%)
2	NDP	B	302	-	51,52,52	1.59	4 (7%)	71,80,80	1.44	13 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	A	302	-	-	3/34/77/77	0/5/5/5
2	NDP	B	302	-	-	3/34/77/77	0/5/5/5

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	302	NDP	C4N-C3N	-6.32	1.38	1.50
2	B	302	NDP	C4N-C3N	-6.19	1.38	1.50
2	B	302	NDP	C4N-C5N	-5.67	1.34	1.49
2	A	302	NDP	C4N-C5N	-5.37	1.35	1.49
2	B	302	NDP	PA-O3	-3.13	1.56	1.59
2	A	302	NDP	P2B-O2B	-2.96	1.54	1.59
2	A	302	NDP	C2N-C3N	2.73	1.42	1.35
2	B	302	NDP	C2N-C3N	2.40	1.41	1.35
2	A	302	NDP	C5A-C4A	-2.05	1.35	1.39
2	A	302	NDP	C8A-N9A	2.00	1.40	1.37

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	302	NDP	C3N-C2N-N1N	-4.62	116.42	123.20
2	B	302	NDP	C3N-C2N-N1N	-4.49	116.61	123.20
2	A	302	NDP	C1D-N1N-C2N	-3.35	115.63	121.14
2	B	302	NDP	N6A-C6A-N1A	3.13	125.34	118.38
2	B	302	NDP	C2A-N1A-C6A	3.08	123.80	118.73
2	B	302	NDP	N3A-C2A-N1A	-3.03	124.00	128.58
2	B	302	NDP	C1D-N1N-C2N	-2.69	116.71	121.14
2	A	302	NDP	O7N-C7N-N7N	-2.63	116.99	122.89

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	302	NDP	C2D-C1D-N1N	-2.58	106.96	113.31
2	A	302	NDP	C3D-C2D-C1D	-2.56	96.62	101.46
2	B	302	NDP	O7N-C7N-N7N	-2.55	117.17	122.89
2	B	302	NDP	O4D-C1D-C2D	-2.47	101.33	106.62
2	A	302	NDP	O4B-C1B-N9A	2.42	112.74	108.09
2	A	302	NDP	N9A-C8A-N7A	-2.37	110.58	113.94
2	A	302	NDP	C6N-N1N-C2N	2.31	121.80	119.32
2	B	302	NDP	O2B-P2B-O1X	-2.30	101.12	109.33
2	B	302	NDP	C5A-C6A-N6A	-2.25	117.72	123.29
2	A	302	NDP	O4B-C4B-C3B	2.24	109.60	105.15
2	B	302	NDP	O2N-PN-O1N	2.15	122.42	112.44
2	A	302	NDP	O2A-PA-O1A	2.14	122.41	112.44
2	A	302	NDP	O4B-C1B-C2B	2.12	110.24	106.59
2	B	302	NDP	C6N-N1N-C2N	2.11	121.58	119.32
2	B	302	NDP	O4B-C4B-C5B	-2.06	102.72	109.33
2	A	302	NDP	O4D-C1D-C2D	-2.04	102.25	106.62
2	A	302	NDP	O2N-PN-O1N	2.04	121.93	112.44

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	302	NDP	C3B-C2B-O2B-P2B
2	B	302	NDP	C3B-C2B-O2B-P2B
2	A	302	NDP	O4D-C1D-N1N-C2N
2	B	302	NDP	O4D-C1D-N1N-C2N
2	A	302	NDP	C1B-C2B-O2B-P2B
2	B	302	NDP	C1B-C2B-O2B-P2B


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	302	NDP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring

in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	285/301 (94%)	0.30	11 (3%)	43 47	11, 18, 30, 45	4 (1%)
1	B	295/301 (98%)	0.24	15 (5%)	33 36	11, 17, 28, 42	1 (0%)
All	All	580/602 (96%)	0.27	26 (4%)	38 42	11, 18, 29, 45	5 (0%)

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	ALA	5.4
1	A	142	ASP	5.0
1	B	2	ALA	4.3
1	B	166	ARG	4.1
1	B	294	PHE	3.9
1	B	140	LEU	3.8
1	A	296	GLU	3.8
1	A	295	LEU	3.7
1	B	142	ASP	3.6
1	A	240	TYR	3.2
1	B	141	ASN	3.2
1	B	296	GLU	3.1
1	A	33	GLY	3.0
1	B	29	GLU	2.9
1	A	251	LYS	2.9
1	B	90	GLU	2.8
1	A	141	ASN	2.7
1	B	112	GLU	2.6
1	B	47	ASP	2.5
1	B	244	ASP	2.4
1	B	3	PRO	2.3
1	A	221	MET	2.2
1	A	140	LEU	2.1
1	A	27	LEU	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	32	PRO	2.0
1	B	54	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NDP	A	302	48/48	0.97	0.07	13,21,28,32	0
2	NDP	B	302	48/48	0.97	0.06	12,16,24,29	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around NDP A 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NDP B 302:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.