



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 10:16 PM UTC

PDB ID : 8A4U / pdb_00008a4u
Title : Crystal structure of human cathepsin L with CAA0225
Authors : Falke, S.; Lieske, J.; Guenther, S.; Reinke, P.Y.A.; Ewert, W.; Loboda, J.; Karnicar, K.; Usenik, A.; Lindic, N.; Sekirnik, A.; Chapman, H.N.; Hinrichs, W.; Turk, D.; Meents, A.
Deposited on : 2022-06-13
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	220	3% 94% 6%
1	B	220	87% 10% .
1	C	220	89% 7% .
1	D	220	93% . .

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 13801 atoms, of which 6464 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Cathepsin L.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	220	Total	C	H	N	O	S	0	1	0
			3280	1065	1575	282	344	14			
1	B	213	Total	C	H	N	O	S	0	3	0
			3230	1050	1556	277	333	14			
1	C	213	Total	C	H	N	O	S	0	1	0
			3193	1039	1537	274	329	14			
1	D	214	Total	C	H	N	O	S	0	2	0
			3220	1047	1546	277	336	14			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	110	ALA	THR	engineered mutation	UNP P07711
B	110	ALA	THR	engineered mutation	UNP P07711
C	110	ALA	THR	engineered mutation	UNP P07711
D	110	ALA	THR	engineered mutation	UNP P07711

- Molecule 2 is (2S,3S)-N3-[2-(4-hydroxyphenyl)ethyl]-N2-[(2S)-1-oxidanylidene-3-phenyl-1-[(phenylmethyl)amino]propan-2-yl]oxirane-2,3-dicarboxamide (CCD ID: L2X) (formula: C₂₈H₂₉N₃O₅) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	0	0
			65	28	29	3	5		
2	B	1	Total	C	H	N	O	0	0
			65	28	29	3	5		
2	C	1	Total	C	H	N	O	0	0
			65	28	29	3	5		
2	D	1	Total	C	H	N	O	0	0
			65	28	29	3	5		

- Molecule 3 is SODIUM ION (CCD ID: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Na	0	0
			2	2		
3	C	1	Total	Na	0	0
			1	1		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			17	4	10	3		
4	A	1	Total	C	H	O	0	0
			17	4	10	3		
4	A	1	Total	C	H	O	0	0
			17	4	10	3		
4	A	1	Total	C	H	O	0	0
			17	4	10	3		
4	A	1	Total	C	H	O	0	0
			17	4	10	3		
4	B	1	Total	C	H	O	0	0
			17	4	10	3		
4	C	1	Total	C	H	O	0	0
			17	4	10	3		
4	C	1	Total	C	H	O	0	0
			17	4	10	3		
4	D	1	Total	C	H	O	0	0
			17	4	10	3		

- Molecule 5 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: C₆H₁₄O₄).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	H	O	0	0
			24	6	14	4		

- Molecule 6 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	C	1	Total	C	H	O	0	0
			10	2	6	2		
6	D	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 7 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	D	1	Total	C	H	O	0	0
			31	8	18	5		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	90	Total	O	0	0
			90	90		
8	B	93	Total	O	0	0
			93	93		
8	C	90	Total	O	0	0
			90	90		
8	D	114	Total	O	0	0
			114	114		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cathepsin L



- Molecule 1: Cathepsin L



- Molecule 1: Cathepsin L



- Molecule 1: Cathepsin L



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	57.00Å 62.87Å 65.18Å 104.04° 95.83° 116.05°	Depositor
Resolution (Å)	44.14 – 1.90 44.14 – 1.90	Depositor EDS
% Data completeness (in resolution range)	94.0 (44.14-1.90) 94.1 (44.14-1.90)	Depositor EDS
R_{merge}	0.33	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.58 (at 1.89Å)	Xtriage
Refinement program	PHENIX 1.13-2998_9999	Depositor
R, R_{free}	0.178 , 0.209 0.179 , 0.208	Depositor DCC
R_{free} test set	2000 reflections (3.52%)	wwPDB-VP
Wilson B-factor (Å ²)	26.8	Xtriage
Anisotropy	0.169	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 47.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	13801	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, L2X, PEG, CSD, PGE, EDO, PG4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.52	0/1739	0.52	0/2350
1	B	0.49	0/1707	0.49	0/2302
1	C	0.50	0/1689	0.48	0/2279
1	D	0.53	0/1706	0.51	0/2302
All	All	0.51	0/6841	0.50	0/9233

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1705	1575	1577	8	0
1	B	1674	1556	1553	11	0
1	C	1656	1537	1536	9	0
1	D	1674	1546	1543	5	0
2	A	36	29	0	2	0
2	B	36	29	0	1	0
2	C	36	29	0	0	0
2	D	36	29	0	0	0
3	A	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	1	0	0	0	0
4	A	35	50	50	1	0
4	B	7	10	10	1	0
4	C	14	20	20	0	0
4	D	7	10	10	0	0
5	B	10	14	14	0	0
6	C	4	6	6	0	0
6	D	4	6	6	0	0
7	D	13	18	18	0	0
8	A	90	0	0	0	0
8	B	93	0	0	0	0
8	C	90	0	0	0	0
8	D	114	0	0	1	0
All	All	7337	6464	6343	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:93:ALA:O	1:B:94:THR:HG23	2.09	0.53
1:A:195:MET:HE1	1:A:200:LYS:HE3	1.92	0.51
1:D:34:LEU:HD23	1:D:48:LEU:HD13	1.91	0.51
1:B:69:LEU:HD13	2:B:301:L2X:C11	2.41	0.50
1:C:118:GLN:HE22	1:C:121:ALA:HB2	1.78	0.49
1:D:123:MET:HE3	1:D:170:TYR:CE1	2.48	0.48
1:C:118:GLN:NE2	1:C:121:ALA:HB2	2.31	0.46
1:A:181:LYS:NZ	1:A:204:ASP:OD2	2.48	0.45
1:B:124:LYS:NZ	1:D:155:ASP:OD1	2.45	0.45
1:C:7:TRP:CE2	1:C:130:GLY:HA2	2.52	0.44
1:C:25:CSD:OD2	1:C:164:GLY:N	2.46	0.44
1:B:10:LYS:HE3	4:B:302:PEG:H22	2.00	0.44
1:C:58:GLY:N	1:C:59:PRO:CD	2.81	0.44
1:D:118:GLN:NE2	8:D:409:HOH:O	2.51	0.44
1:A:196:GLY:HA3	1:C:196:GLY:HA3	2.01	0.43
1:A:178:ASP:HB3	1:D:179:ASN:O	2.19	0.43
1:C:117:LYS:HD2	1:C:213:SER:O	2.19	0.42
1:A:69:LEU:HD12	1:A:72:TYR:CE2	2.54	0.42
1:B:109:ASP:OD1	1:B:109:ASP:C	2.62	0.42
1:B:185:VAL:HB	1:B:199:VAL:HG13	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:137:ASP:OD2	1:B:140:HIS:HE1	2.03	0.41
1:A:18:ASN:H	4:A:305:PEG:H11	1.85	0.41
1:B:153:GLU:H	1:B:208:HIS:CE1	2.38	0.41
1:A:25:CSD:SG	2:A:301:L2X:C18	3.09	0.41
1:B:82:GLY:HA2	1:B:107:ALA:O	2.20	0.41
1:A:25:CSD:SG	2:A:301:L2X:O3	2.79	0.41
1:C:8:ARG:NH2	1:C:184:LEU:HD21	2.36	0.41
1:B:7:TRP:CE2	1:B:130:GLY:HA2	2.56	0.40
1:B:115:ILE:HD11	1:B:217:TYR:HE1	1.85	0.40
1:C:153:GLU:H	1:C:208:HIS:CE1	2.40	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	218/220 (99%)	212 (97%)	6 (3%)	0	100	100
1	B	211/220 (96%)	204 (97%)	7 (3%)	0	100	100
1	C	209/220 (95%)	204 (98%)	5 (2%)	0	100	100
1	D	211/220 (96%)	203 (96%)	8 (4%)	0	100	100
All	All	849/880 (96%)	823 (97%)	26 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	178/177 (101%)	174 (98%)	4 (2%)	45	42
1	B	174/177 (98%)	171 (98%)	3 (2%)	53	52
1	C	172/177 (97%)	170 (99%)	2 (1%)	63	63
1	D	174/177 (98%)	173 (99%)	1 (1%)	78	81
All	All	698/708 (99%)	688 (99%)	10 (1%)	59	59

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	156	CYS
1	A	174	SER
1	A	176	GLU
1	A	220	VAL
1	B	87	GLU
1	B	103	LYS
1	B	156	CYS
1	C	41	LYS
1	C	156	CYS
1	D	156	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	78	GLN
1	A	108	ASN
1	C	19	GLN
1	C	208	HIS
1	D	108	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	CSD	D	25	1	4,7,8	1.16	1 (25%)	1,8,10	1.61	0
1	CSD	C	25	1	4,7,8	2.43	1 (25%)	1,8,10	1.21	0
1	CSD	A	25	1	4,7,8	1.01	0	1,8,10	2.99	1 (100%)
1	CSD	B	25	1	4,7,8	1.38	1 (25%)	1,8,10	1.94	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSD	D	25	1	-	1/2/6/8	-
1	CSD	C	25	1	-	1/2/6/8	-
1	CSD	A	25	1	-	1/2/6/8	-
1	CSD	B	25	1	-	1/2/6/8	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	25	CSD	OD1-SG	-4.71	1.43	1.47
1	B	25	CSD	OD1-SG	-2.40	1.45	1.47
1	D	25	CSD	OD1-SG	-2.09	1.45	1.47

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	25	CSD	OD1-SG-CB	-2.99	100.10	105.60

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	25	CSD	CA-CB-SG-OD1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
1	B	25	CSD	CA-CB-SG-OD1
1	C	25	CSD	CA-CB-SG-OD1
1	D	25	CSD	CA-CB-SG-OD1

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	C	25	CSD	1	0
1	A	25	CSD	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 20 ligands modelled in this entry, 3 are monoatomic - leaving 17 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	PEG	A	308	-	6,6,6	0.27	0	5,5,5	0.11	0
7	PG4	D	304	-	12,12,12	0.25	0	11,11,11	0.23	0
5	PGE	B	303	-	9,9,9	0.30	0	8,8,8	0.16	0
4	PEG	C	303	-	6,6,6	0.33	0	5,5,5	0.09	0
4	PEG	D	302	-	6,6,6	0.27	0	5,5,5	0.14	0
4	PEG	A	306	-	6,6,6	0.30	0	5,5,5	0.18	0
2	L2X	C	301	-	39,39,39	0.20	0	52,52,52	0.39	0
2	L2X	A	301	-	39,39,39	0.22	0	52,52,52	0.41	0
4	PEG	A	305	-	6,6,6	0.27	0	5,5,5	0.34	0
4	PEG	A	307	-	6,6,6	0.27	0	5,5,5	0.20	0
6	EDO	C	305	-	3,3,3	0.54	0	2,2,2	0.40	0
4	PEG	B	302	-	6,6,6	0.22	0	5,5,5	0.23	0
4	PEG	C	304	-	6,6,6	0.23	0	5,5,5	0.07	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	PEG	A	304	-	6,6,6	0.20	0	5,5,5	0.14	0
2	L2X	B	301	-	39,39,39	0.20	0	52,52,52	0.31	0
6	EDO	D	303	-	3,3,3	0.67	0	2,2,2	0.24	0
2	L2X	D	301	-	39,39,39	0.20	0	52,52,52	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEG	A	308	-	-	3/4/4/4	-
7	PG4	D	304	-	-	5/10/10/10	-
5	PGE	B	303	-	-	3/7/7/7	-
4	PEG	C	303	-	-	0/4/4/4	-
4	PEG	D	302	-	-	1/4/4/4	-
4	PEG	A	306	-	-	2/4/4/4	-
2	L2X	C	301	-	-	10/31/36/36	0/4/4/4
2	L2X	A	301	-	-	9/31/36/36	0/4/4/4
4	PEG	A	305	-	-	3/4/4/4	-
4	PEG	A	307	-	-	2/4/4/4	-
6	EDO	C	305	-	-	0/1/1/1	-
4	PEG	B	302	-	-	1/4/4/4	-
4	PEG	C	304	-	-	2/4/4/4	-
4	PEG	A	304	-	-	2/4/4/4	-
2	L2X	B	301	-	-	8/31/36/36	0/4/4/4
6	EDO	D	303	-	-	1/1/1/1	-
2	L2X	D	301	-	-	6/31/36/36	0/4/4/4

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (58) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	L2X	N1-C16-C17-C18
2	A	301	L2X	O1-C16-C17-C18
2	A	301	L2X	N1-C16-C17-O2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	301	L2X	O1-C16-C17-O2
2	A	301	L2X	O2-C18-C19-N2
2	A	301	L2X	O2-C18-C19-O3
2	A	301	L2X	C18-C19-N2-C20
2	B	301	L2X	N1-C16-C17-C18
2	B	301	L2X	O1-C16-C17-C18
2	B	301	L2X	N1-C16-C17-O2
2	B	301	L2X	O1-C16-C17-O2
2	B	301	L2X	O2-C18-C19-N2
2	B	301	L2X	O2-C18-C19-O3
2	C	301	L2X	N1-C16-C17-C18
2	C	301	L2X	O1-C16-C17-C18
2	C	301	L2X	N1-C16-C17-O2
2	C	301	L2X	O1-C16-C17-O2
2	C	301	L2X	O2-C18-C19-N2
2	C	301	L2X	O2-C18-C19-O3
2	C	301	L2X	C18-C19-N2-C20
2	D	301	L2X	N1-C16-C17-C18
2	D	301	L2X	O1-C16-C17-C18
2	D	301	L2X	N1-C16-C17-O2
2	D	301	L2X	O1-C16-C17-O2
2	D	301	L2X	O2-C18-C19-N2
2	D	301	L2X	O2-C18-C19-O3
4	A	305	PEG	O1-C1-C2-O2
2	A	301	L2X	O3-C19-N2-C20
5	B	303	PGE	O2-C3-C4-O3
4	A	306	PEG	O1-C1-C2-O2
4	A	304	PEG	O2-C3-C4-O4
4	A	307	PEG	O2-C3-C4-O4
4	A	308	PEG	O1-C1-C2-O2
4	B	302	PEG	O2-C3-C4-O4
4	A	308	PEG	O2-C3-C4-O4
5	B	303	PGE	O1-C1-C2-O2
2	C	301	L2X	O3-C19-N2-C20
7	D	304	PG4	O4-C7-C8-O5
7	D	304	PG4	O2-C3-C4-O3
4	A	304	PEG	C1-C2-O2-C3
7	D	304	PG4	C4-C3-O2-C2
2	A	301	L2X	C21-C20-N2-C19
5	B	303	PGE	C3-C4-O3-C5
4	D	302	PEG	O1-C1-C2-O2
4	A	308	PEG	C1-C2-O2-C3

Continued on next page...

Continued from previous page...




Mol	Chain	Res	Type	Atoms
4	A	305	PEG	C1-C2-O2-C3
4	C	304	PEG	C1-C2-O2-C3
4	A	305	PEG	O2-C3-C4-O4
4	A	307	PEG	C4-C3-O2-C2
2	C	301	L2X	C21-C20-N2-C19
4	A	306	PEG	C1-C2-O2-C3
2	B	301	L2X	N1-C7-C8-O
2	B	301	L2X	N1-C7-C8-N
7	D	304	PG4	C3-C4-O3-C5
6	D	303	EDO	O1-C1-C2-O2
2	C	301	L2X	N1-C7-C8-N
7	D	304	PG4	C5-C6-O4-C7
4	C	304	PEG	C4-C3-O2-C2


There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	L2X	2	0
4	A	305	PEG	1	0
4	B	302	PEG	1	0
2	B	301	L2X	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	219/220 (99%)	-0.04	7 (3%) 50 54	14, 32, 60, 82	1 (0%)
1	B	212/220 (96%)	-0.09	0 100 100	15, 32, 58, 81	3 (1%)
1	C	212/220 (96%)	0.06	1 (0%) 87 89	19, 35, 62, 95	1 (0%)
1	D	213/220 (96%)	-0.18	0 100 100	15, 31, 53, 81	2 (0%)
All	All	856/880 (97%)	-0.06	8 (0%) 81 83	14, 32, 60, 95	7 (0%)

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	178	ASP	3.6
1	A	177	SER	2.8
1	A	179	ASN	2.6
1	A	174	SER	2.5
1	A	172	PHE	2.3
1	C	2	PRO	2.1
1	A	180	ASN	2.1
1	A	206	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CSD	A	25	8/9	0.95	0.08	22,30,37,40	0
1	CSD	C	25	8/9	0.96	0.06	23,31,34,41	0
1	CSD	D	25	8/9	0.96	0.07	21,29,35,41	0
1	CSD	B	25	8/9	0.97	0.06	25,32,39,43	0

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	PEG	B	302	7/7	0.79	0.15	63,75,76,76	0
6	EDO	D	303	4/4	0.79	0.15	45,55,56,57	0
4	PEG	A	307	7/7	0.84	0.15	63,75,76,78	0
4	PEG	D	302	7/7	0.84	0.16	63,75,76,76	0
4	PEG	A	308	7/7	0.84	0.18	62,75,76,76	0
5	PGE	B	303	10/10	0.85	0.14	43,52,63,64	0
4	PEG	C	303	7/7	0.86	0.14	51,62,74,75	0
4	PEG	A	305	7/7	0.86	0.12	47,57,61,61	0
2	L2X	B	301	36/36	0.88	0.12	32,47,57,61	0
4	PEG	A	306	7/7	0.88	0.11	51,61,67,68	0
6	EDO	C	305	4/4	0.88	0.24	71,85,90,93	0
2	L2X	D	301	36/36	0.88	0.10	34,42,49,53	0
2	L2X	A	301	36/36	0.89	0.11	31,45,61,61	0
4	PEG	C	304	7/7	0.89	0.11	63,76,78,78	0
3	NA	A	303	1/1	0.90	0.09	51,51,51,51	0
2	L2X	C	301	36/36	0.90	0.10	31,41,56,60	0
7	PG4	D	304	13/13	0.90	0.15	52,62,72,73	0
4	PEG	A	304	7/7	0.93	0.13	55,66,67,67	0
3	NA	C	302	1/1	0.97	0.05	32,32,32,32	0
3	NA	A	302	1/1	0.97	0.04	28,28,28,28	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around L2X B 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around L2X D 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around L2X A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around L2X C 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.