



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 02:57 PM UTC

PDB ID : 4A5B / pdb_00004a5b
Title : Crystal structure of the C258S/C268S variant of Toxoplasma gondii nucleoside triphosphate diphosphohydrolase 1 (NTPDase1)
Authors : Krug, U.; Zebisch, M.; Strater, N.
Deposited on : 2011-10-24
Resolution : 2.48 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7589 (2.50-2.46)
Clashscore	190562	8295 (2.50-2.46)
Ramachandran outliers	187476	8164 (2.50-2.46)
Sidechain outliers	187428	8166 (2.50-2.46)
RSRZ outliers	180081	7593 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	611	
1	B	611	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 9496 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NUCLEOSIDE-TRIPHOSPHATASE 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	592	Total	C	N	O	S	0	0	0
			4619	2902	810	882	25			
1	B	593	Total	C	N	O	S	0	0	0
			4630	2908	815	882	25			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	25	MET	-	expression tag	UNP Q27895
A	629	GLU	-	expression tag	UNP Q27895
A	630	HIS	-	expression tag	UNP Q27895
A	631	HIS	-	expression tag	UNP Q27895
A	632	HIS	-	expression tag	UNP Q27895
A	633	HIS	-	expression tag	UNP Q27895
A	634	HIS	-	expression tag	UNP Q27895
A	635	HIS	-	expression tag	UNP Q27895
A	258	SER	CYS	engineered mutation	UNP Q27895
A	268	SER	CYS	engineered mutation	UNP Q27895
B	25	MET	-	expression tag	UNP Q27895
B	629	GLU	-	expression tag	UNP Q27895
B	630	HIS	-	expression tag	UNP Q27895
B	631	HIS	-	expression tag	UNP Q27895
B	632	HIS	-	expression tag	UNP Q27895
B	633	HIS	-	expression tag	UNP Q27895
B	634	HIS	-	expression tag	UNP Q27895
B	635	HIS	-	expression tag	UNP Q27895
B	258	SER	CYS	engineered mutation	UNP Q27895
B	268	SER	CYS	engineered mutation	UNP Q27895


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	140	Total 140	O 140	0	0
2	B	107	Total 107	O 107	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: NUCLEOSIDE-TRIPHOSPHATASE 2



• Molecule 1: NUCLEOSIDE-TRIPHOSPHATASE 2



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	73.73Å 150.40Å 242.36Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.22 – 2.48 39.22 – 2.48	Depositor EDS
% Data completeness (in resolution range)	(Not available) (39.22-2.48) 99.1 (39.22-2.48)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.51 (at 2.48Å)	Xtriage
Refinement program	BUSTER 2.8.0	Depositor
R, R_{free}	0.176 , 0.226 0.183 , 0.232	Depositor DCC
R_{free} test set	972 reflections (2.04%)	wwPDB-VP
Wilson B-factor (Å ²)	32.2	Xtriage
Anisotropy	0.108	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 46.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9496	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.87	1/4710 (0.0%)	1.41	28/6376 (0.4%)
1	B	0.85	1/4722 (0.0%)	1.43	42/6393 (0.7%)
All	All	0.86	2/9432 (0.0%)	1.42	70/12769 (0.5%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	142	VAL	CA-C	5.82	1.58	1.52
1	A	142	VAL	CA-CB	5.11	1.56	1.54

All (70) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	38	THR	N-CA-C	-9.12	101.92	113.23
1	A	611	PHE	CA-C-N	6.62	130.00	120.79
1	A	611	PHE	C-N-CA	6.62	130.00	120.79
1	A	383	ASP	CA-CB-CG	6.60	119.20	112.60
1	B	141	TYR	CA-C-N	6.60	125.61	120.33
1	B	141	TYR	C-N-CA	6.60	125.61	120.33
1	A	403	ILE	N-CA-C	-6.56	106.58	111.90
1	B	103	ILE	CB-CA-C	6.45	120.36	111.92
1	B	611	PHE	CA-C-N	6.38	131.33	121.19
1	B	611	PHE	C-N-CA	6.38	131.33	121.19
1	A	601	GLN	N-CA-C	6.25	117.76	111.07
1	B	229	ARG	N-CA-C	6.22	113.73	108.13
1	A	39	GLU	N-CA-C	-6.00	105.94	113.20
1	A	343	ASN	CA-CB-CG	5.99	118.58	112.60
1	B	510	ASP	CA-CB-CG	5.95	118.55	112.60
1	A	103	ILE	CB-CA-C	5.91	119.39	111.94
1	B	450	GLU	CA-C-N	5.73	127.89	120.44
1	B	450	GLU	C-N-CA	5.73	127.89	120.44
1	B	189	ALA	N-CA-C	5.64	118.97	111.75
1	A	261	ASP	CA-CB-CG	5.57	118.17	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	601	GLN	N-CA-C	5.50	116.95	111.07
1	B	619	ALA	N-CA-C	5.38	117.22	111.36
1	B	70	ASP	CA-CB-CG	5.38	117.98	112.60
1	A	123	TYR	N-CA-C	5.35	119.87	113.23
1	B	614	GLU	CB-CG-CD	5.35	121.70	112.60
1	A	419	ASP	CA-CB-CG	5.34	117.94	112.60
1	B	179	LEU	N-CA-C	5.32	119.49	112.89
1	B	605	ILE	CA-C-N	5.32	127.41	120.28
1	B	605	ILE	C-N-CA	5.32	127.41	120.28
1	A	403	ILE	CA-C-N	5.31	127.65	120.38
1	A	403	ILE	C-N-CA	5.31	127.65	120.38
1	B	520	GLU	CA-C-N	5.30	127.32	120.44
1	B	520	GLU	C-N-CA	5.30	127.32	120.44
1	B	514	ASP	CA-C-N	5.28	127.41	120.60
1	B	514	ASP	C-N-CA	5.28	127.41	120.60
1	A	254	ASP	CA-CB-CG	5.27	117.87	112.60
1	A	515	VAL	CA-C-N	5.25	127.32	120.28
1	A	515	VAL	C-N-CA	5.25	127.32	120.28
1	B	188	THR	CA-C-N	5.24	128.02	120.38
1	B	188	THR	C-N-CA	5.24	128.02	120.38
1	A	400	ASN	CA-CB-CG	5.23	117.83	112.60
1	B	517	GLU	CA-C-N	5.23	127.24	120.44
1	B	517	GLU	C-N-CA	5.23	127.24	120.44
1	A	514	ASP	CA-C-N	5.23	127.23	120.70
1	A	514	ASP	C-N-CA	5.23	127.23	120.70
1	B	261	ASP	CA-CB-CG	5.22	117.82	112.60
1	A	203	PHE	CA-CB-CG	-5.17	108.63	113.80
1	B	407	VAL	CA-C-N	5.17	129.41	121.19
1	B	407	VAL	C-N-CA	5.17	129.41	121.19
1	A	375	ASP	CA-CB-CG	5.16	117.76	112.60
1	B	454	VAL	N-CA-C	5.16	115.44	110.74
1	B	412	GLN	CA-C-N	5.16	127.46	120.44
1	B	412	GLN	C-N-CA	5.16	127.46	120.44
1	A	54	ASN	CA-C-N	5.15	127.18	120.28
1	A	54	ASN	C-N-CA	5.15	127.18	120.28
1	B	473	SER	CA-C-N	5.12	127.47	120.46
1	B	473	SER	C-N-CA	5.12	127.47	120.46
1	B	46	LYS	CA-C-N	5.10	127.11	120.28
1	B	46	LYS	C-N-CA	5.10	127.11	120.28
1	B	215	PRO	CA-C-N	5.08	127.40	120.54
1	B	215	PRO	C-N-CA	5.08	127.40	120.54
1	B	54	ASN	CA-C-N	5.08	127.34	120.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	54	ASN	C-N-CA	5.08	127.34	120.38
1	B	573	ALA	CA-C-N	5.07	124.92	120.10
1	B	573	ALA	C-N-CA	5.07	124.92	120.10
1	A	413	CYS	CA-C-N	5.03	127.27	120.38
1	A	413	CYS	C-N-CA	5.03	127.27	120.38
1	A	174	VAL	CA-C-N	5.01	126.99	120.28
1	A	174	VAL	C-N-CA	5.01	126.99	120.28
1	A	179	LEU	N-CA-C	5.01	118.65	112.54

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4619	0	4563	32	0
1	B	4630	0	4579	29	0
2	A	140	0	0	3	0
2	B	107	0	0	1	0
All	All	9496	0	9142	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:415:GLU:HG3	1:B:458:LEU:HD22	1.65	0.78
1:A:303:ASN:O	1:A:307:GLU:HG2	1.96	0.66
1:B:253:GLU:HG3	1:B:313:ARG:HG3	1.79	0.63
1:B:104:ARG:HH21	1:B:167:GLN:HE21	1.47	0.62
1:A:335:LEU:HD21	1:A:460:LEU:HD22	1.85	0.58
1:B:184:MET:HG2	1:B:227:TRP:HB3	1.88	0.56
1:A:185:LEU:HB3	1:A:228:THR:HG23	1.85	0.56
1:A:326:GLN:HA	1:A:331:SER:HB3	1.87	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:304:LEU:HD12	1:B:317:ALA:HB3	1.90	0.54
1:A:134:ASP:HB3	1:A:137:LEU:HD12	1.90	0.54
1:A:88:CYS:HB2	1:A:91:LYS:HB2	1.90	0.53
1:A:53:ARG:HA	1:A:56:GLU:HG2	1.89	0.53
1:A:141:TYR:HA	1:A:144:GLN:HE21	1.73	0.53
1:A:479:ARG:HD3	1:B:405:PHE:CE1	2.44	0.52
1:A:85:THR:HG22	1:A:95:ILE:HG12	1.91	0.52
1:B:304:LEU:CD1	1:B:317:ALA:HB3	2.40	0.51
1:A:82:LEU:HD12	1:A:103:ILE:HG21	1.92	0.51
1:A:366:SER:HB3	1:A:380:VAL:HG11	1.94	0.48
1:A:329:MET:HE3	2:A:2074:HOH:O	2.13	0.48
1:A:422:LYS:HB3	1:A:426:GLU:HG3	1.96	0.48
1:A:519:ALA:O	1:A:523:ARG:HB2	2.14	0.47
1:A:415:GLU:HG3	1:A:458:LEU:HD12	1.97	0.47
1:B:110:THR:HG22	1:B:152:LEU:HD13	1.97	0.47
1:B:68:VAL:O	1:B:79:ASN:HB2	2.15	0.47
1:A:184:MET:HG2	1:A:227:TRP:HB3	1.96	0.46
1:A:141:TYR:HA	1:A:144:GLN:NE2	2.31	0.46
1:B:541:LEU:HB3	1:B:545:ARG:HB3	1.98	0.46
1:A:410:GLU:O	1:A:414:ARG:HG3	2.17	0.45
1:B:123:TYR:HB3	1:B:141:TYR:CD2	2.51	0.45
1:A:423:PRO:HD2	1:A:426:GLU:HG2	1.99	0.45
1:A:484:THR:HG23	1:A:576:ASN:HB3	1.98	0.45
1:B:252:GLY:HA3	1:B:257:ARG:HB2	1.99	0.45
1:B:412:GLN:NE2	2:B:2082:HOH:O	2.50	0.44
1:B:185:LEU:HB3	1:B:228:THR:HG23	1.99	0.44
1:B:442:PHE:O	1:B:445:CYS:HB3	2.18	0.44
1:B:514:ASP:HB3	1:B:517:GLU:HB2	2.00	0.44
1:B:88:CYS:HB2	1:B:91:LYS:HB2	1.99	0.44
1:A:419:ASP:OD1	1:A:421:THR:HB	2.18	0.44
1:B:537:PRO:HB2	1:B:550:LEU:HB3	2.00	0.43
1:B:284:GLN:HG3	1:B:323:SER:HB2	2.00	0.43
1:A:302:VAL:HG13	1:A:307:GLU:HG3	2.01	0.43
1:A:338:LYS:HB2	1:A:389:LEU:HD11	2.01	0.43
1:B:341:CYS:HB2	1:B:430:ILE:HG21	2.00	0.43
1:A:362:GLN:HG3	1:A:437:GLU:HG3	2.01	0.42
1:A:525:PHE:HD1	1:A:541:LEU:HD21	1.85	0.42
1:B:305:GLN:OE1	1:B:316:SER:HA	2.20	0.41
1:A:276:VAL:HG11	1:A:567:LEU:HD21	2.01	0.41
1:B:568:LEU:HD22	1:B:581:ILE:HD13	2.02	0.41
1:A:249:ARG:HA	2:A:2056:HOH:O	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:291:GLU:HA	1:B:292:GLY:HA2	1.74	0.41
1:B:361:PHE:HB2	1:B:530:VAL:HG21	2.03	0.41
1:A:245:ASN:HB3	1:A:251:LEU:HB2	2.02	0.41
1:B:63:LEU:HD21	1:B:82:LEU:HD13	2.03	0.41
1:A:468:SER:HA	1:B:464:ILE:HG13	2.03	0.40
1:A:113:ARG:HD3	2:A:2025:HOH:O	2.20	0.40
1:B:185:LEU:O	1:B:228:THR:HA	2.21	0.40
1:B:418:ILE:HG23	1:B:427:ARG:HG2	2.03	0.40
1:B:424:LEU:HG	1:B:428:MET:HE2	2.02	0.40
1:A:368:GLY:HA2	1:A:387:ASN:HA	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	588/611 (96%)	580 (99%)	8 (1%)	0	100	100
1	B	591/611 (97%)	575 (97%)	16 (3%)	0	100	100
All	All	1179/1222 (96%)	1155 (98%)	24 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	510/527 (97%)	495 (97%)	15 (3%)	37	62
1	B	511/527 (97%)	496 (97%)	15 (3%)	37	62
All	All	1021/1054 (97%)	991 (97%)	30 (3%)	37	62

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	39	GLU
1	A	44	VAL
1	A	52	LEU
1	A	91	LYS
1	A	170	GLU
1	A	286	VAL
1	A	294	VAL
1	A	302	VAL
1	A	309	LEU
1	A	340	LEU
1	A	437	GLU
1	A	447	SER
1	A	469	SER
1	A	524	GLU
1	A	549	LYS
1	B	42	ILE
1	B	55	LEU
1	B	85	THR
1	B	133	VAL
1	B	286	VAL
1	B	291	GLU
1	B	295	LEU
1	B	309	LEU
1	B	312	GLU
1	B	351	ILE
1	B	366	SER
1	B	451	SER
1	B	458	LEU
1	B	538	VAL
1	B	549	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	43	ASN
1	A	54	ASN
1	A	144	GLN
1	A	326	GLN
1	A	400	ASN
1	A	548	GLN
1	B	167	GLN
1	B	195	HIS
1	B	208	HIS
1	B	399	HIS
1	B	412	GLN
1	B	448	GLN
1	B	543	ASN
1	B	608	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	592/611 (96%)	-0.37	9 (1%) 72 69	16, 31, 56, 109	0
1	B	593/611 (97%)	-0.30	7 (1%) 76 74	17, 35, 70, 93	0
All	All	1185/1222 (96%)	-0.34	16 (1%) 73 71	16, 33, 63, 109	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	542	PRO	5.2
1	A	36	ALA	3.9
1	A	291	GLU	2.9
1	A	629	GLU	2.9
1	B	590	PRO	2.9
1	B	294	VAL	2.7
1	B	293	THR	2.6
1	A	37	ASP	2.5
1	B	628	LEU	2.4
1	A	293	THR	2.3
1	A	38	THR	2.2
1	A	292	GLY	2.2
1	B	292	GLY	2.2
1	B	534	THR	2.1
1	A	590	PRO	2.1
1	B	629	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.