



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 12:44 PM UTC

PDB ID : 7A53 / pdb_00007a53
Title : Structure of DYRK1A in complex with compound 7
Authors : Dokurno, P.; Surgenor, A.E.; Hubbard, R.E.
Deposited on : 2020-08-20
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	359	
1	B	359	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
1	SEP	B	145	-	-	X	-

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 5672 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dual specificity tyrosine-phosphorylation-regulated kinase 1A.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	335	2737	1759	476	482	2	18	0	1	0
1	B	334	2667	1718	455	474	2	18	0	1	0

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	127	MET	-	initiating methionine	UNP Q13627
A	128	GLY	-	expression tag	UNP Q13627
A	129	SER	-	expression tag	UNP Q13627
A	130	SER	-	expression tag	UNP Q13627
A	131	HIS	-	expression tag	UNP Q13627
A	132	HIS	-	expression tag	UNP Q13627
A	133	HIS	-	expression tag	UNP Q13627
A	134	HIS	-	expression tag	UNP Q13627
A	135	HIS	-	expression tag	UNP Q13627
A	136	HIS	-	expression tag	UNP Q13627
A	137	SER	-	expression tag	UNP Q13627
A	138	SER	-	expression tag	UNP Q13627
A	139	GLY	-	expression tag	UNP Q13627
A	140	LEU	-	expression tag	UNP Q13627
A	141	VAL	-	expression tag	UNP Q13627
A	142	PRO	-	expression tag	UNP Q13627
A	143	ARG	-	expression tag	UNP Q13627
A	144	GLY	-	expression tag	UNP Q13627
A	145	SEP	-	expression tag	UNP Q13627
A	146	HIS	-	expression tag	UNP Q13627
A	147	MET	-	expression tag	UNP Q13627
B	127	MET	-	initiating methionine	UNP Q13627
B	128	GLY	-	expression tag	UNP Q13627
B	129	SER	-	expression tag	UNP Q13627
B	130	SER	-	expression tag	UNP Q13627

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	131	HIS	-	expression tag	UNP Q13627
B	132	HIS	-	expression tag	UNP Q13627
B	133	HIS	-	expression tag	UNP Q13627
B	134	HIS	-	expression tag	UNP Q13627
B	135	HIS	-	expression tag	UNP Q13627
B	136	HIS	-	expression tag	UNP Q13627
B	137	SER	-	expression tag	UNP Q13627
B	138	SER	-	expression tag	UNP Q13627
B	139	GLY	-	expression tag	UNP Q13627
B	140	LEU	-	expression tag	UNP Q13627
B	141	VAL	-	expression tag	UNP Q13627
B	142	PRO	-	expression tag	UNP Q13627
B	143	ARG	-	expression tag	UNP Q13627
B	144	GLY	-	expression tag	UNP Q13627
B	145	SEP	-	expression tag	UNP Q13627
B	146	HIS	-	expression tag	UNP Q13627
B	147	MET	-	expression tag	UNP Q13627

- Molecule 2 is (3 {E})-3-[(5-methylfuran-2-yl)methylidene]-1 {H}-indol-2-one (CCD ID: QYZ) (formula: C₁₄H₁₁NO₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	N			O
2	A	1	17	14	1	2	0	0
2	B	1	17	14	1	2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	143	Total 143	O 143	0	0
3	B	91	Total 91	O 91	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Dual specificity tyrosine-phosphorylation-regulated kinase 1A



- Molecule 1: Dual specificity tyrosine-phosphorylation-regulated kinase 1A



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	57.20Å 84.38Å 84.09Å 90.00° 107.56° 90.00°	Depositor
Resolution (Å)	25.00 – 2.20 25.00 – 2.20	Depositor EDS
% Data completeness (in resolution range)	88.5 (25.00-2.20) 88.5 (25.00-2.20)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.61 (at 2.19Å)	Xtrriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.218 , 0.277 0.223 , 0.279	Depositor DCC
R_{free} test set	1765 reflections (4.55%)	wwPDB-VP
Wilson B-factor (Å ²)	39.6	Xtrriage
Anisotropy	0.087	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 27.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.029 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5672	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: QYZ, SEP, PTR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.02	0/2775	1.38	1/3735 (0.0%)
1	B	1.01	0/2706	1.41	1/3659 (0.0%)
All	All	1.02	0/5481	1.39	2/7394 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	358	GLU	CB-CA-C	6.08	118.32	109.20
1	A	380	PRO	N-CA-CB	5.31	106.16	103.19

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2737	0	2764	22	0
1	B	2667	0	2617	60	0
2	A	17	0	0	0	0
2	B	17	0	0	1	0
3	A	143	0	0	5	1
3	B	91	0	0	11	1
All	All	5672	0	5381	82	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:159:TYR:OH	1:B:228:PHE:CZ	1.69	1.29
1:B:398:LEU:HG	1:B:399:PRO:HD2	1.29	1.08
1:B:339:ASP:HB2	3:B:627:HOH:O	1.62	0.99
1:B:374:VAL:HG11	1:B:405:LEU:HD11	1.51	0.93
1:A:247:ASP:OD2	3:A:601:HOH:O	1.92	0.88
1:B:159:TYR:OH	1:B:228:PHE:CE1	2.30	0.85
1:B:398:LEU:HG	1:B:399:PRO:CD	2.06	0.84
1:B:398:LEU:CG	1:B:399:PRO:HD2	2.08	0.84
1:A:366:GLU:OE2	3:A:602:HOH:O	1.98	0.81
1:B:175:LYS:HG3	1:B:240:MET:HE2	1.63	0.80
1:B:375:GLU:OE1	3:B:602:HOH:O	2.08	0.71
1:A:317[A]:ARG:HG2	3:A:633:HOH:O	1.96	0.65
1:A:214:ASP:HA	1:A:218:LYS:HD2	1.78	0.64
1:A:146:HIS:CE1	1:A:167:LYS:HE3	2.33	0.64
1:B:159:TYR:HD2	1:B:187:ILE:HD11	1.63	0.63
1:B:159:TYR:CD2	1:B:187:ILE:HD11	2.33	0.63
1:B:245:LEU:HD13	1:B:354:MET:HE2	1.80	0.62
1:A:406:LYS:O	1:A:406:LYS:HG2	2.01	0.61
1:B:159:TYR:HD1	1:B:159:TYR:N	2.00	0.60
1:A:217:MET:HB3	1:A:275:PHE:HB2	1.83	0.59
1:B:465:LYS:N	3:B:601:HOH:O	2.36	0.58
1:B:159:TYR:N	1:B:159:TYR:CD1	2.72	0.57
1:A:146:HIS:CE1	1:A:167:LYS:CE	2.89	0.56
1:B:159:TYR:HH	1:B:228:PHE:HZ	0.62	0.56
1:B:145:SEP:O3P	1:B:145:SEP:N	2.38	0.56
1:B:185:VAL:HG11	3:B:650:HOH:O	2.06	0.55
1:A:366:GLU:OE1	1:A:390:LYS:NZ	2.40	0.55
1:B:463:ASP:C	3:B:601:HOH:O	2.50	0.55
1:B:444:HIS:CE1	3:B:624:HOH:O	2.59	0.54
1:A:328:ARG:HD3	1:A:332:VAL:HG12	1.90	0.54
1:B:175:LYS:CG	1:B:240:MET:HE2	2.36	0.53
1:A:214:ASP:HA	1:A:218:LYS:CD	2.40	0.52
1:B:230:PHE:CE2	1:B:231:ARG:HD3	2.44	0.52
1:A:287:ASP:OD2	3:A:603:HOH:O	2.19	0.52
1:B:400:ASP:OD1	1:B:400:ASP:N	2.26	0.51
1:B:145:SEP:P	1:B:145:SEP:H	2.33	0.51
1:B:183:GLU:OE2	1:B:225:LYS:NZ	2.44	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:178:ASP:HB3	3:B:650:HOH:O	2.10	0.50
1:B:185:VAL:HG12	1:B:225:LYS:HE3	1.94	0.50
1:B:161:ILE:HD12	1:B:176:ALA:HB2	1.92	0.50
1:B:374:VAL:CG1	1:B:405:LEU:HD11	2.34	0.49
1:A:315:GLY:HA2	1:A:317[A]:ARG:HH11	1.76	0.49
1:B:277:ALA:HA	1:B:340:LEU:CD2	2.43	0.49
1:A:188:LYS:NZ	1:A:203:GLU:OE2	2.46	0.49
1:B:388:ALA:HB3	1:B:391:ALA:HB2	1.93	0.48
1:B:452:PHE:HB2	1:B:478:PHE:CE1	2.48	0.48
1:B:464:PRO:HG2	3:B:687:HOH:O	2.12	0.48
1:B:151:ASN:HA	1:B:161:ILE:HG22	1.95	0.48
1:A:188:LYS:HE2	3:A:653:HOH:O	2.15	0.47
1:B:221:ILE:HD12	1:B:308:PHE:HZ	1.81	0.46
1:B:318:ILE:N	1:B:318:ILE:HD13	2.31	0.46
1:A:263:ARG:HB2	1:A:478:PHE:CZ	2.50	0.46
1:A:297:ASN:ND2	1:A:300:ARG:HG3	2.31	0.46
1:B:145:SEP:O	1:B:233:HIS:HE1	1.99	0.44
1:B:159:TYR:OH	1:B:228:PHE:HZ	1.42	0.43
1:A:217:MET:CE	1:A:274:LEU:HD23	2.48	0.43
1:B:273:LEU:HB3	1:B:470:PRO:HG3	1.99	0.43
1:B:319[B]:TYR:CE2	1:B:321:PTR:HB2	2.53	0.43
1:A:315:GLY:HA2	1:A:317[A]:ARG:HE	1.83	0.43
1:B:200:ALA:O	1:B:204:VAL:HG23	2.18	0.43
1:B:219:TYR:O	1:B:304:LYS:HE3	2.18	0.43
1:B:164:LEU:HA	1:B:174:VAL:HG12	2.01	0.43
1:B:363:GLY:HA2	1:B:368:ASP:OD2	2.19	0.43
1:A:164:LEU:HD11	1:A:172:GLN:HB2	2.01	0.43
1:B:159:TYR:CZ	1:B:228:PHE:HZ	2.33	0.43
1:B:165:ILE:HG21	2:B:501:QYZ:C1	2.49	0.42
1:A:217:MET:HE1	1:A:274:LEU:HD23	2.00	0.42
1:B:390:LYS:HB3	1:B:390:LYS:HE2	1.84	0.42
1:B:371:ASN:OD1	1:B:405:LEU:HD22	2.21	0.41
1:B:463:ASP:CG	3:B:601:HOH:O	2.63	0.41
1:B:145:SEP:HB3	1:B:191:LYS:HG2	2.03	0.41
1:B:221:ILE:HD12	1:B:308:PHE:CZ	2.55	0.41
1:B:142:PRO:HD2	3:B:673:HOH:O	2.20	0.41
1:B:198:ASN:O	1:B:202:ILE:HG13	2.19	0.41
1:B:243:TYR:HE1	1:B:248:LEU:HD13	1.86	0.41
1:B:319[B]:TYR:HD1	1:B:319[B]:TYR:HA	1.69	0.41
1:B:463:ASP:OD1	3:B:601:HOH:O	2.22	0.40
1:B:319[B]:TYR:CD2	1:B:321:PTR:HB2	2.56	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:318:ILE:HG22	1:A:319:TYR:CD2	2.57	0.40
1:B:398:LEU:CD2	1:B:399:PRO:HD2	2.50	0.40
1:B:288:LEU:HD23	1:B:288:LEU:HA	1.92	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:680:HOH:O	3:B:668:HOH:O[2_555]	1.92	0.28

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	330/359 (92%)	313 (95%)	15 (4%)	2 (1%)	21	23
1	B	329/359 (92%)	305 (93%)	23 (7%)	1 (0%)	36	42
All	All	659/718 (92%)	618 (94%)	38 (6%)	3 (0%)	24	27

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	217	MET
1	B	143	ARG
1	A	143	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	294/315 (93%)	277 (94%)	17 (6%)	18	22
1	B	278/315 (88%)	255 (92%)	23 (8%)	10	11
All	All	572/630 (91%)	532 (93%)	40 (7%)	15	16

All (40) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	142	PRO
1	A	165	ILE
1	A	172	GLN
1	A	201	GLN
1	A	212	LYS
1	A	217	MET
1	A	231	ARG
1	A	303	ILE
1	A	317[A]	ARG
1	A	317[B]	ARG
1	A	318	ILE
1	A	322	ILE
1	A	397	LYS
1	A	400	ASP
1	A	408	THR
1	A	470	PRO
1	A	480	LYS
1	B	143	ARG
1	B	153	GLU
1	B	159	TYR
1	B	161	ILE
1	B	207	LEU
1	B	215	THR
1	B	225	LYS
1	B	239	GLU
1	B	240	MET
1	B	300	ARG
1	B	303	ILE
1	B	310	SER
1	B	318	ILE
1	B	319[A]	TYR
1	B	319[B]	TYR
1	B	350	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	396	GLU
1	B	398	LEU
1	B	400	ASP
1	B	405	LEU
1	B	408	THR
1	B	453	LYS
1	B	480	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	151	ASN
1	A	201	GLN
1	A	227	HIS
1	A	251	ASN
1	A	253	ASN
1	A	425	ASN
1	A	469	GLN
1	B	201	GLN
1	B	211	ASN
1	B	227	HIS
1	B	233	HIS
1	B	251	ASN
1	B	267	GLN
1	B	425	ASN
1	B	469	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	PTR	A	321	1	15,16,17	0.48	0	17,22,24	1.25	3 (17%)
1	SEP	A	145	1	8,9,10	0.71	0	7,12,14	0.88	0
1	SEP	B	145	1	8,9,10	0.55	0	7,12,14	1.08	0
1	PTR	B	321	1	15,16,17	0.98	1 (6%)	17,22,24	0.65	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	A	321	1	-	1/10/11/13	0/1/1/1
1	SEP	A	145	1	-	5/6/8/10	-
1	SEP	B	145	1	-	2/6/8/10	-
1	PTR	B	321	1	-	0/10/11/13	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	321	PTR	P-OH	-3.29	1.53	1.59

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	321	PTR	OH-CZ-CE1	2.75	127.47	119.22
1	A	321	PTR	OH-CZ-CE2	-2.30	112.31	119.22
1	A	321	PTR	P-OH-CZ	2.11	131.40	123.88

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	321	PTR	O-C-CA-CB
1	A	145	SEP	N-CA-CB-OG
1	A	145	SEP	CB-OG-P-O2P
1	A	145	SEP	CB-OG-P-O3P
1	B	145	SEP	N-CA-CB-OG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
1	B	145	SEP	CA-CB-OG-P
1	A	145	SEP	CB-OG-P-O1P
1	A	145	SEP	CA-CB-OG-P

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	145	SEP	4	0
1	B	321	PTR	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	QYZ	B	501	-	19,19,19	0.70	0	26,27,27	0.62	0
2	QYZ	A	501	-	19,19,19	0.84	1 (5%)	26,27,27	0.65	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	QYZ	B	501	-	-	0/4/16/16	0/3/3/3
2	QYZ	A	501	-	-	2/4/16/16	0/3/3/3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	QYZ	C6-C5	-2.09	1.39	1.43

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	501	QYZ	C5-C6-C7-C10
2	A	501	QYZ	C5-C6-C7-C8


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	501	QYZ	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	333/359 (92%)	-0.13	3 (0%) 81 79	18, 41, 79, 116	2 (0%)
1	B	332/359 (92%)	0.52	14 (4%) 40 37	37, 60, 96, 124	2 (0%)
All	All	665/718 (92%)	0.19	17 (2%) 57 54	18, 50, 92, 124	4 (0%)

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	159	TYR	6.4
1	B	319[A]	TYR	4.9
1	A	217	MET	3.6
1	B	338	TYR	3.2
1	B	142	PRO	3.0
1	B	408	THR	2.8
1	B	184	TRP	2.8
1	B	320	GLN	2.8
1	B	308	PHE	2.7
1	B	398	LEU	2.6
1	B	180	VAL	2.4
1	B	243	TYR	2.4
1	B	440	GLY	2.3
1	A	142	PRO	2.2
1	B	399	PRO	2.2
1	B	216	GLU	2.1
1	A	219	TYR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
1	SEP	B	145	10/11	0.80	0.09	65,88,119,121	0
1	SEP	A	145	10/11	0.82	0.11	63,76,91,92	0
1	PTR	B	321	16/17	0.89	0.13	63,83,104,105	0
1	PTR	A	321	16/17	0.92	0.09	41,57,66,68	0

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	QYZ	B	501	17/17	0.81	0.13	45,53,62,63	0
2	QYZ	A	501	17/17	0.86	0.15	48,59,74,76	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.