



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:45 PM UTC

PDB ID : 1A6A / pdb_00001a6a
Title : THE STRUCTURE OF AN INTERMEDIATE IN CLASS II MHC MATURATION: CLIP BOUND TO HLA-DR3
Authors : Ghosh, P.; Amaya, M.; Mellins, E.; Wiley, D.C.
Deposited on : 1998-02-22
Resolution : 2.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1009 (2.76-2.76)
Clashscore	190562	1044 (2.76-2.76)
Ramachandran outliers	187476	1024 (2.76-2.76)
Sidechain outliers	187428	1024 (2.76-2.76)
RSRZ outliers	180081	1009 (2.76-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	176	52% 42% 5%
2	B	187	49% 47%
3	C	15	67% 27% 7%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3158 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class II histocompatibility antigen, DR alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	176	Total	C	N	O	S	6	0	0
			1447	940	236	266	5			


- Molecule 2 is a protein called HLA class II histocompatibility antigen, DR-1 beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	187	Total	C	N	O	S	60	0	0
			1548	973	281	289	5			

- Molecule 3 is a protein called HLA class II histocompatibility antigen, gamma chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	15	Total	C	N	O	S	0	0	0
			113	72	20	18	3			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	8	Total	O	0	0
			8	8		
5	B	13	Total	O	0	0
			13	13		
5	C	1	Total	O	0	0
			1	1		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HLA class II histocompatibility antigen, DR alpha chain



- Molecule 2: HLA class II histocompatibility antigen, DR-1 beta chain



- Molecule 3: HLA class II histocompatibility antigen, gamma chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 1 2	Depositor
Cell constants a, b, c, α , β , γ	78.45Å 78.45Å 159.10Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	6.00 – 2.75 6.00 – 2.75	Depositor EDS
% Data completeness (in resolution range)	97.9 (6.00-2.75) 88.0 (6.00-2.75)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.46 (at 2.61Å)	Xtriage
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.246 , 0.325 0.238 , 0.314	Depositor DCC
R_{free} test set	1696 reflections (10.00%)	wwPDB-VP
Wilson B-factor (Å ²)	51.2	Xtriage
Anisotropy	0.194	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 90.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.057 for -h,-k,l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3158	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:
NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.98	1/1492 (0.1%)	1.16	11/2035 (0.5%)
2	B	0.96	2/1591 (0.1%)	1.00	3/2160 (0.1%)
3	C	0.99	0/114	0.98	1/151 (0.7%)
All	All	0.97	3/3197 (0.1%)	1.07	15/4346 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	31	PHE	C-N	-11.56	1.18	1.33
1	A	85	VAL	CA-CB	-6.07	1.47	1.53
2	B	175	VAL	CA-CB	-5.99	1.47	1.54

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	31	PHE	O-C-N	-9.15	112.66	123.27
1	A	99	LEU	N-CA-C	7.91	121.87	111.75
1	A	86	PRO	CA-C-N	7.49	127.65	120.31
1	A	86	PRO	C-N-CA	7.49	127.65	120.31
1	A	85	VAL	CA-C-N	6.65	124.52	119.66
1	A	85	VAL	C-N-CA	6.65	124.52	119.66
1	A	31	ILE	N-CA-C	-6.44	103.98	111.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	15	ASN	N-CA-C	6.33	122.13	113.77
2	B	13	SER	N-CA-C	-5.64	100.00	109.07
1	A	145	PHE	N-CA-C	5.31	117.90	109.50
1	A	177	HIS	N-CA-C	5.19	116.99	110.24
1	A	79	TYR	N-CA-C	5.13	118.52	111.54
1	A	157	THR	N-CA-C	-5.09	106.20	114.09
3	C	94	ALA	N-CA-C	-5.05	103.05	110.52
2	B	83	TYR	N-CA-C	-5.04	105.48	110.97

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	150	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1447	0	1387	63	0
2	B	1548	0	1463	79	0
3	C	113	0	128	10	0
4	A	28	0	26	3	0
5	A	8	0	0	1	0
5	B	13	0	0	1	0
5	C	1	0	0	0	0
All	All	3158	0	3004	137	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

All (137) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:74:ARG:NH1	3:C:94:ALA:HB1	1.86	0.90

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:176:GLU:HG2	2:B:183:PRO:HB3	1.56	0.85
1:A:16:PRO:HG2	2:B:6:ARG:HA	1.58	0.84
2:B:74:ARG:HH11	3:C:94:ALA:HB1	1.39	0.83
1:A:8:ILE:HG23	1:A:25:ASP:HB3	1.60	0.83
2:B:31:PHE:CE2	2:B:36:GLU:HB2	2.15	0.81
2:B:180:VAL:HG11	2:B:184:LEU:HD11	1.64	0.79
1:A:15:ASN:HB2	1:A:70:LEU:HD21	1.64	0.79
2:B:71:LYS:HE3	3:C:97:LEU:HD13	1.67	0.75
1:A:71:GLU:O	1:A:75:LYS:HD3	1.88	0.73
2:B:27:LEU:HG	2:B:29:ARG:HD3	1.74	0.70
1:A:95:SER:HB2	1:A:96:PRO:HD2	1.74	0.69
4:A:181:NAG:O7	4:A:181:NAG:C3	2.42	0.68
2:B:50:VAL:HG23	2:B:51:THR:HG23	1.76	0.67
1:A:115:PRO:O	1:A:137:PHE:CE1	2.48	0.66
1:A:126:LYS:H	1:A:126:LYS:HD2	1.59	0.66
1:A:8:ILE:HD11	1:A:10:ALA:HB2	1.76	0.66
4:A:181:NAG:O7	4:A:181:NAG:H3	1.96	0.66
2:B:82:ASN:O	2:B:86:VAL:HG22	1.96	0.65
1:A:15:ASN:CB	1:A:70:LEU:HD21	2.27	0.64
1:A:11:GLU:HA	1:A:21:GLU:O	1.97	0.64
1:A:176:LYS:HA	1:A:176:LYS:HE3	1.78	0.64
2:B:16:HIS:HB2	2:B:25:ARG:HB3	1.81	0.62
1:A:119:VAL:HG22	1:A:165:VAL:HG22	1.81	0.62
2:B:97:PRO:HB3	2:B:122:PHE:HB3	1.82	0.61
1:A:23:MET:HE2	1:A:25:ASP:HB2	1.81	0.61
1:A:92:LEU:HD11	2:B:156:GLN:NE2	2.17	0.60
1:A:7:ILE:HG12	1:A:26:PHE:HD1	1.66	0.60
1:A:65:VAL:HG11	3:C:96:PRO:O	2.03	0.59
1:A:87:PRO:HB2	1:A:109:ILE:HG23	1.85	0.58
1:A:109:ILE:HD12	1:A:109:ILE:N	2.18	0.58
2:B:77:ASN:O	3:C:92:ARG:HD2	2.04	0.57
1:A:121:TRP:O	1:A:127:PRO:HA	2.04	0.57
2:B:13:SER:HB3	2:B:26:TYR:HE1	1.67	0.57
2:B:52:GLU:O	2:B:55:ARG:HG2	2.05	0.57
1:A:79:TYR:N	1:A:79:TYR:CD1	2.72	0.57
1:A:94:ASN:HB2	1:A:104:VAL:HB	1.87	0.57
1:A:92:LEU:HD11	2:B:156:GLN:HE22	1.70	0.56
1:A:105:LEU:HB3	1:A:121:TRP:CH2	2.40	0.56
2:B:62:ASN:HA	2:B:68:LEU:HD21	1.86	0.56
2:B:159:VAL:HG23	2:B:159:VAL:O	2.05	0.56
1:A:168:TRP:CZ2	2:B:6:ARG:NH2	2.74	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:163:THR:O	2:B:165:PRO:HD3	2.06	0.56
2:B:12:THR:HG22	2:B:14:GLU:HG3	1.88	0.56
2:B:71:LYS:HZ3	2:B:74:ARG:HH12	1.53	0.55
2:B:87:GLU:HG3	2:B:91:VAL:HB	1.89	0.55
1:A:119:VAL:HB	1:A:149:HIS:CE1	2.42	0.54
2:B:134:ASN:OD1	2:B:169:GLU:HB2	2.08	0.54
1:A:14:LEU:CD2	1:A:116:VAL:HG23	2.38	0.54
2:B:71:LYS:NZ	2:B:74:ARG:NH1	2.55	0.54
1:A:47:GLU:O	1:A:51:PHE:HD1	1.90	0.54
2:B:68:LEU:O	2:B:72:ARG:HG3	2.07	0.54
2:B:24:VAL:HG11	2:B:79:CYS:HB3	1.90	0.53
1:A:61:ALA:O	1:A:64:ALA:HB3	2.08	0.53
2:B:116:VAL:HG22	2:B:160:MET:HG2	1.92	0.52
1:A:103:ASN:HB3	1:A:153:PHE:CE1	2.45	0.52
2:B:177:HIS:O	2:B:180:VAL:HG22	2.09	0.52
1:A:119:VAL:HG13	1:A:163:CYS:SG	2.50	0.51
2:B:81:HIS:CD2	3:C:92:ARG:HG3	2.44	0.51
2:B:122:PHE:HB2	2:B:177:HIS:NE2	2.25	0.51
2:B:28:ASP:HB3	2:B:40:PHE:HB3	1.92	0.51
1:A:115:PRO:O	1:A:137:PHE:CZ	2.63	0.51
2:B:29:ARG:HB2	2:B:31:PHE:HE1	1.76	0.51
1:A:23:MET:CE	1:A:25:ASP:HB2	2.41	0.50
1:A:38:LYS:NZ	1:A:38:LYS:HB2	2.26	0.50
1:A:14:LEU:HD13	2:B:8:LEU:HD13	1.94	0.49
1:A:137:PHE:CE1	1:A:147:LYS:HD2	2.47	0.49
1:A:117:VAL:CG1	1:A:118:ASN:N	2.75	0.49
2:B:17:PHE:HB3	2:B:20:GLY:O	2.12	0.49
2:B:69:GLU:HA	2:B:72:ARG:HB2	1.95	0.48
1:A:92:LEU:HG	1:A:93:THR:N	2.28	0.48
1:A:147:LYS:HE3	1:A:149:HIS:NE2	2.27	0.48
2:B:14:GLU:O	2:B:26:TYR:HA	2.13	0.48
2:B:61:TRP:CG	3:C:97:LEU:HD23	2.49	0.48
2:B:172:THR:HG22	2:B:187:GLU:HG2	1.94	0.48
2:B:71:LYS:HZ3	2:B:74:ARG:NH1	2.12	0.48
2:B:144:SER:HB2	2:B:159:VAL:HG12	1.95	0.48
2:B:161:LEU:HD22	2:B:163:THR:OG1	2.14	0.48
1:A:79:TYR:N	1:A:79:TYR:HD1	2.12	0.47
2:B:129:VAL:CG1	2:B:159:VAL:HG21	2.44	0.47
2:B:131:TRP:NE1	2:B:161:LEU:HB2	2.29	0.47
2:B:41:ASP:HB3	2:B:44:VAL:HG23	1.96	0.47
1:A:67:LYS:O	1:A:70:LEU:HB3	2.14	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:33:ASN:C	2:B:34:GLN:HG3	2.38	0.47
1:A:91:VAL:HG21	1:A:176:LYS:O	2.15	0.47
2:B:29:ARG:HG3	2:B:29:ARG:NH1	2.30	0.47
2:B:55:ARG:N	2:B:56:PRO:CD	2.78	0.46
2:B:144:SER:CB	2:B:159:VAL:HG12	2.46	0.46
2:B:29:ARG:CB	2:B:31:PHE:CE1	2.98	0.46
1:A:16:PRO:CG	2:B:6:ARG:HA	2.39	0.46
2:B:24:VAL:HB	2:B:75:VAL:HG13	1.98	0.46
2:B:49:ALA:HB2	2:B:55:ARG:HA	1.98	0.46
1:A:164:ARG:HH22	1:A:166:GLU:CD	2.24	0.46
2:B:141:GLY:O	2:B:162:GLU:HG2	2.16	0.45
2:B:71:LYS:CE	3:C:97:LEU:HD13	2.41	0.45
2:B:129:VAL:HG13	2:B:175:VAL:HG22	1.99	0.45
2:B:137:GLU:O	2:B:139:LYS:HG3	2.17	0.45
2:B:29:ARG:HB2	2:B:31:PHE:CE1	2.51	0.45
1:A:112:PHE:CE2	1:A:117:VAL:CG2	3.00	0.45
1:A:58:GLY:O	1:A:61:ALA:HB3	2.17	0.45
2:B:177:HIS:CG	2:B:178:PRO:HD2	2.52	0.45
1:A:5:HIS:ND1	1:A:26:PHE:CZ	2.85	0.45
1:A:15:ASN:HB3	1:A:16:PRO:HD3	1.99	0.44
1:A:53:SER:O	3:C:90:LYS:HA	2.17	0.44
1:A:117:VAL:HG12	1:A:118:ASN:N	2.31	0.44
1:A:39:LYS:N	1:A:39:LYS:HD2	2.31	0.44
2:B:27:LEU:CG	2:B:29:ARG:HD3	2.45	0.43
1:A:34:VAL:HG21	1:A:59:ALA:HB2	1.99	0.43
2:B:67:LEU:O	2:B:70:GLN:HB3	2.18	0.43
1:A:118:ASN:OD1	4:A:200:NAG:C7	2.66	0.43
2:B:174:GLN:OE1	2:B:185:THR:HB	2.18	0.43
2:B:97:PRO:HA	2:B:120:SER:O	2.18	0.43
1:A:115:PRO:HG3	1:A:145:PHE:CE2	2.54	0.43
2:B:58:ALA:O	2:B:62:ASN:ND2	2.52	0.43
2:B:94:ARG:NH2	5:B:198:HOH:O	2.51	0.43
1:A:109:ILE:HG22	1:A:112:PHE:CE1	2.54	0.42
2:B:127:ILE:HG12	2:B:128:GLU:N	2.34	0.42
1:A:108:PHE:HE1	1:A:146:ARG:HB3	1.85	0.42
1:A:45:LEU:HD12	1:A:48:PHE:CZ	2.54	0.42
2:B:128:GLU:CD	2:B:130:ARG:HH22	2.28	0.42
1:A:133:SER:HB2	1:A:150:TYR:HB2	2.02	0.42
2:B:39:ARG:O	2:B:39:ARG:HG3	2.20	0.42
1:A:96:PRO:HA	5:A:204:HOH:O	2.20	0.41
1:A:135:THR:O	1:A:147:LYS:NZ	2.48	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:13:SER:HB3	2:B:26:TYR:CE1	2.52	0.41
2:B:28:ASP:O	2:B:39:ARG:HA	2.21	0.41
2:B:29:ARG:HB3	2:B:31:PHE:CE1	2.55	0.41
2:B:31:PHE:CD2	2:B:36:GLU:HB2	2.54	0.41
2:B:66:ASP:O	2:B:70:GLN:HB2	2.20	0.41
2:B:81:HIS:O	2:B:85:VAL:HG23	2.21	0.41
1:A:26:PHE:CD2	2:B:90:THR:HB	2.55	0.41
2:B:127:ILE:HG12	2:B:128:GLU:H	1.85	0.41
1:A:111:LYS:HG2	1:A:140:ARG:CZ	2.51	0.40
1:A:161:TYR:O	1:A:178:TRP:N	2.53	0.40
2:B:166:ARG:O	2:B:169:GLU:HG2	2.22	0.40
1:A:141:GLU:C	1:A:143:HIS:H	2.29	0.40
2:B:81:HIS:HD2	3:C:92:ARG:HG3	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	174/176 (99%)	150 (86%)	23 (13%)	1 (1%)	21	38
2	B	185/187 (99%)	159 (86%)	22 (12%)	4 (2%)	5	10
3	C	13/15 (87%)	13 (100%)	0	0	100	100
All	All	372/378 (98%)	322 (87%)	45 (12%)	5 (1%)	9	18

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	138	GLU
2	B	167	SER
2	B	168	GLY
2	B	112	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	171	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	161/161 (100%)	149 (92%)	12 (8%)	12	24
2	B	172/172 (100%)	162 (94%)	10 (6%)	18	34
3	C	13/13 (100%)	13 (100%)	0	100	100
All	All	346/346 (100%)	324 (94%)	22 (6%)	16	29

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	ILE
1	A	47	GLU
1	A	79	TYR
1	A	83	THR
1	A	92	LEU
1	A	113	THR
1	A	126	LYS
1	A	128	VAL
1	A	171	ASP
1	A	172	GLU
1	A	174	LEU
1	A	176	LYS
2	B	21	THR
2	B	34	GLN
2	B	38	VAL
2	B	88	SER
2	B	96	HIS
2	B	138	GLU
2	B	140	THR
2	B	142	VAL
2	B	161	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	185	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	15	ASN
1	A	84	ASN
2	B	37	ASN
2	B	77	ASN
2	B	81	HIS
2	B	112	HIS
2	B	150	ASN
2	B	156	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	A	200	1	14,14,15	0.82	1 (7%)	17,19,21	0.77	0
4	NAG	A	181	1	14,14,15	0.58	0	17,19,21	0.70	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	200	1	-	0/6/23/26	0/1/1/1
4	NAG	A	181	1	-	3/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	200	NAG	C1-C2	2.50	1.55	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	181	NAG	O7-C7-C8	-2.05	118.40	122.05

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	181	NAG	C3-C2-N2-C7
4	A	181	NAG	O5-C5-C6-O6
4	A	181	NAG	C4-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	200	NAG	1	0
4	A	181	NAG	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	B	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	31:PHE	C	32:HIS	N	1.18

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	176/176 (100%)	-0.56	1 (0%) 85 86	15, 42, 73, 85	1 (0%)
2	B	187/187 (100%)	-0.36	1 (0%) 87 87	22, 43, 77, 85	14 (7%)
3	C	15/15 (100%)	-0.35	0 100 100	22, 41, 69, 71	0
All	All	378/378 (100%)	-0.45	2 (0%) 87 87	15, 43, 73, 85	15 (3%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	166	ARG	2.6
1	A	100	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	A	181	14/15	0.58	0.15	74,77,79,79	0
4	NAG	A	200	14/15	0.59	0.14	85,86,88,88	0

6.5 Other polymers [i](#)

There are no such residues in this entry.