



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 02:17 PM UTC

PDB ID : 3A6E / pdb\_00003a6e  
Title : W174F mutant creatininase, type I  
Authors : Nakajima, Y.; Yamashita, K.; Ito, K.; Yoshimoto, T.  
Deposited on : 2009-08-31  
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	260	
1	B	260	
1	C	260	
1	D	260	
1	E	260	

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	260	 6% 80% 17% ..

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 12873 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Creatinine amidohydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	257	1985	1271	336	367	11	0	0	0
1	B	257	1981	1268	335	367	11	0	0	0
1	C	257	1981	1268	335	367	11	0	0	0
1	D	257	1985	1271	336	367	11	0	0	0
1	E	257	1985	1271	336	367	11	0	0	0
1	F	257	1985	1271	336	367	11	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	174	PHE	TRP	engineered mutation	UNP P83772
B	174	PHE	TRP	engineered mutation	UNP P83772
C	174	PHE	TRP	engineered mutation	UNP P83772
D	174	PHE	TRP	engineered mutation	UNP P83772
E	174	PHE	TRP	engineered mutation	UNP P83772
F	174	PHE	TRP	engineered mutation	UNP P83772

- Molecule 2 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mn	0	0
			1	1		
2	B	1	Total	Mn	0	0
			1	1		
2	C	1	Total	Mn	0	0
			1	1		

*Continued on next page...*


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	D	1	Total Mn 1 1	0	0
2	E	1	Total Mn 1 1	0	0
2	F	1	Total Mn 1 1	0	0

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Zn 1 1	0	0
3	B	1	Total Zn 1 1	0	0
3	C	1	Total Zn 1 1	0	0
3	D	1	Total Zn 1 1	0	0
3	E	1	Total Zn 1 1	0	0
3	F	1	Total Zn 1 1	0	0

- Molecule 4 is CACODYLATE ION (CCD ID: CAC) (formula: C<sub>2</sub>H<sub>6</sub>AsO<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	As	C	O	0	0
			5	1	2	2		
4	B	1	Total	As	C	O	0	0
			5	1	2	2		
4	C	1	Total	As	C	O	0	0
			5	1	2	2		
4	D	1	Total	As	C	O	0	0
			5	1	2	2		
4	E	1	Total	As	C	O	0	0
			5	1	2	2		
4	F	1	Total	As	C	O	0	0
			5	1	2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	167	Total	O	0	0
			167	167		
5	B	151	Total	O	0	0
			151	151		
5	C	151	Total	O	0	0
			151	151		
5	D	173	Total	O	0	0
			173	173		
5	E	150	Total	O	0	0
			150	150		
5	F	137	Total	O	0	0
			137	137		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Creatinine amidohydrolase




- Molecule 1: Creatinine amidohydrolase




- Molecule 1: Creatinine amidohydrolase





- Molecule 1: Creatinine amidohydrolase







● Molecule 1: Creatinine amidohydrolase



● Molecule 1: Creatinine amidohydrolase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	105.90Å 59.70Å 145.10Å 90.00° 99.70° 90.00°	Depositor
Resolution (Å)	20.00 – 2.00 20.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.6 (20.00-2.00) 98.5 (20.00-2.00)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	7.56 (at 1.97Å)	Xtrriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.196 , 0.229 0.194 , 0.227	Depositor DCC
$R_{free}$ test set	6223 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.6	Xtrriage
Anisotropy	0.298	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 57.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	12873	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.29% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CAC, MN, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.37	0/2033	0.91	8/2765 (0.3%)
1	B	0.36	0/2029	0.90	7/2761 (0.3%)
1	C	0.36	0/2029	0.89	7/2761 (0.3%)
1	D	0.36	0/2033	0.90	7/2765 (0.3%)
1	E	0.35	0/2033	0.86	4/2765 (0.1%)
1	F	0.36	0/2033	0.85	2/2765 (0.1%)
All	All	0.36	0/12190	0.89	35/16582 (0.2%)

There are no bond length outliers.

The worst 5 of 35 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	89	SER	N-CA-C	6.17	119.75	109.95
1	D	89	SER	N-CA-C	5.99	119.48	109.95
1	D	39	HIS	N-CA-C	5.91	120.65	113.38
1	C	91	ASP	N-CA-C	-5.86	102.64	110.55
1	B	127	ILE	N-CA-C	-5.79	104.87	110.72

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1985	0	1968	30	0
1	B	1981	0	1957	20	0
1	C	1981	0	1957	28	0
1	D	1985	0	1968	24	0
1	E	1985	0	1968	31	0
1	F	1985	0	1968	31	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
3	F	1	0	0	0	0
4	A	5	0	0	1	0
4	B	5	0	0	1	0
4	C	5	0	0	1	0
4	D	5	0	0	1	0
4	E	5	0	0	1	0
4	F	5	0	0	1	0
5	A	167	0	0	2	0
5	B	151	0	0	0	0
5	C	151	0	0	1	0
5	D	173	0	0	1	0
5	E	150	0	0	3	0
5	F	137	0	0	1	0
All	All	12873	0	11786	156	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 156 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:160:PRO:O	1:A:164:GLN:HG3	1.82	0.80
1:F:232:THR:HG22	1:F:237:LYS:NZ	2.02	0.74
1:E:120:HIS:HB3	1:E:123:ASN:ND2	2.04	0.72
1:D:28:LEU:HD23	1:D:116:LEU:HD21	1.73	0.70

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:120:HIS:HB3	1:F:123:ASN:ND2	2.06	0.69

There are no symmetry-related clashes.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	255/260 (98%)	252 (99%)	3 (1%)	0	100	100
1	B	255/260 (98%)	249 (98%)	6 (2%)	0	100	100
1	C	255/260 (98%)	250 (98%)	5 (2%)	0	100	100
1	D	255/260 (98%)	252 (99%)	3 (1%)	0	100	100
1	E	255/260 (98%)	240 (94%)	15 (6%)	0	100	100
1	F	255/260 (98%)	246 (96%)	9 (4%)	0	100	100
All	All	1530/1560 (98%)	1489 (97%)	41 (3%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	212/215 (99%)	212 (100%)	0	100	100
1	B	211/215 (98%)	210 (100%)	1 (0%)	81	87

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	211/215 (98%)	210 (100%)	1 (0%)	81	87
1	D	212/215 (99%)	212 (100%)	0	100	100
1	E	212/215 (99%)	210 (99%)	2 (1%)	70	78
1	F	212/215 (99%)	209 (99%)	3 (1%)	59	66
All	All	1270/1290 (98%)	1263 (99%)	7 (1%)	78	85

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	144	GLN
1	F	158	LYS
1	F	232	THR
1	F	212	ASP
1	E	116	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	164	GLN
1	E	76	GLN
1	F	165	GLN
1	E	165	GLN
1	F	75	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry

Of 18 ligands modelled in this entry, 12 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	CAC	A	302	3,2	2,4,4	2.56	2 (100%)	4,6,6	0.42	0
4	CAC	F	307	3,2	2,4,4	2.54	2 (100%)	4,6,6	0.32	0
4	CAC	B	303	3,2	2,4,4	2.53	2 (100%)	4,6,6	0.28	0
4	CAC	C	304	3,2	2,4,4	2.59	2 (100%)	4,6,6	0.33	0
4	CAC	D	305	3,2	2,4,4	2.57	2 (100%)	4,6,6	0.55	0
4	CAC	E	306	3,2	2,4,4	2.49	2 (100%)	4,6,6	0.38	0

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	304	CAC	AS-C2	2.71	1.96	1.90
4	D	305	CAC	AS-C2	2.62	1.96	1.90
4	A	302	CAC	AS-C2	2.60	1.96	1.90
4	B	303	CAC	AS-C2	2.59	1.96	1.90
4	F	307	CAC	AS-C1	2.57	1.96	1.90

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	302	CAC	1	0
4	F	307	CAC	1	0
4	B	303	CAC	1	0
4	C	304	CAC	1	0
4	D	305	CAC	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	E	306	CAC	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	257/260 (98%)	-0.22	3 (1%) 76 76	8, 17, 32, 47	0
1	B	257/260 (98%)	-0.22	4 (1%) 70 70	8, 18, 35, 51	0
1	C	257/260 (98%)	-0.02	9 (3%) 47 46	7, 21, 43, 56	0
1	D	257/260 (98%)	-0.09	3 (1%) 76 76	10, 20, 37, 47	0
1	E	257/260 (98%)	0.44	16 (6%) 26 25	10, 27, 55, 67	0
1	F	257/260 (98%)	0.39	16 (6%) 26 25	10, 26, 52, 62	0
All	All	1542/1560 (98%)	0.05	51 (3%) 49 48	7, 20, 43, 67	0

The worst 5 of 51 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	172	LEU	3.9
1	E	173	GLY	3.3
1	C	171	PHE	3.3
1	E	200	VAL	3.2
1	E	176	ILE	3.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MN	F	300	1/1	0.95	0.12	45,45,45,45	0
4	CAC	E	306	5/5	0.95	0.16	35,38,40,40	0
2	MN	E	300	1/1	0.96	0.14	41,41,41,41	0
4	CAC	F	307	5/5	0.96	0.14	35,37,39,39	0
4	CAC	D	305	5/5	0.97	0.13	25,27,29,30	0
4	CAC	A	302	5/5	0.97	0.12	21,22,23,27	0
4	CAC	C	304	5/5	0.97	0.13	26,29,31,31	0
2	MN	C	300	1/1	0.98	0.13	36,36,36,36	0
4	CAC	B	303	5/5	0.98	0.11	23,25,26,27	0
2	MN	D	300	1/1	0.99	0.10	32,32,32,32	0
2	MN	B	300	1/1	0.99	0.14	30,30,30,30	0
2	MN	A	300	1/1	0.99	0.11	29,29,29,29	0
3	ZN	C	301	1/1	0.99	0.02	23,23,23,23	0
3	ZN	E	301	1/1	0.99	0.03	27,27,27,27	0
3	ZN	F	301	1/1	0.99	0.03	26,26,26,26	0
3	ZN	D	301	1/1	1.00	0.01	22,22,22,22	0
3	ZN	B	301	1/1	1.00	0.01	19,19,19,19	0
3	ZN	A	301	1/1	1.00	0.01	18,18,18,18	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.