



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 15, 2026 – 10:17 AM UTC

PDB ID : 5A64 / pdb_00005a64
Title : Crystal structure of mouse thiamine triphosphatase in complex with thiamine triphosphate.
Authors : Martinez, J.; Truffault, V.; Hothorn, M.
Deposited on : 2015-06-24
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	226	17% 82% 8% 10%
1	B	226	34% 81% • 15%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3268 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called THIAMINE TRIPHOSPHATASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	203	Total	C	N	O	S	0	8	0
			1588	1009	262	311	6			
1	B	192	Total	C	N	O	S	0	0	0
			1448	923	243	277	5			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP Q8JZL3
A	225	SER	-	expression tag	UNP Q8JZL3
B	0	GLY	-	expression tag	UNP Q8JZL3
B	225	SER	-	expression tag	UNP Q8JZL3

- Molecule 2 is Thiamine Triphosphate (CCD ID: V4E) (formula: $C_{12}H_{20}N_4O_{10}P_3S$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	S	
			30	12	4	10	3	1	0
2	B	1	Total	C	N	O	P	S	
			30	12	4	10	3	1	0

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O		
			4	2	2	0	0
3	A	1	Total	C	O		
			4	2	2	0	0
3	A	1	Total	C	O		
			4	2	2	0	0
3	A	1	Total	C	O		
			4	2	2	0	0

- Molecule 4 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: $C_6H_{14}O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			10	6	4		
4	A	1	Total	C	O	0	0
			10	6	4		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	90	Total	O	0	0
			90	90		
5	B	46	Total	O	0	0
			46	46		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: THIAMINE TRIPHOSPHATASE



• Molecule 1: THIAMINE TRIPHOSPHATASE



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	103.18Å 93.58Å 70.78Å 90.00° 93.04° 90.00°	Depositor
Resolution (Å)	70.68 – 2.10 70.67 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.5 (70.68-2.10) 99.6 (70.67-2.10)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.89 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.8.0123	Depositor
R, R_{free}	0.211 , 0.232 0.216 , 0.233	Depositor DCC
R_{free} test set	1963 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	30.2	Xtriage
Anisotropy	0.253	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 50.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.005 for -1/2*h+1/2*k+1,1/2*h-1/2*k+1,1/2*h+1/2*k 0.015 for -1/2*h-1/2*k+1,-1/2*h-1/2*k-1,1/2*h-1/2*k	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3268	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: V4E, EDO, PGE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.79	0/1641	0.89	0/2224
1	B	0.72	0/1474	0.84	0/2001
All	All	0.76	0/3115	0.87	0/4225

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1588	0	1576	13	0
1	B	1448	0	1401	4	0
2	A	30	0	16	1	0
2	B	30	0	16	0	0
3	A	16	0	24	1	0
4	A	20	0	28	2	0
5	A	90	0	0	0	0
5	B	46	0	0	0	0
All	All	3268	0	3061	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:10:ARG:NH1	1:A:11:LYS:O	2.31	0.64
1:B:206:TYR:O	1:B:210:LEU:HD13	2.02	0.60
1:A:62:TRP:C	1:A:63[B]:GLU:HG3	2.29	0.57
1:A:36:ARG:HE	4:A:1221:PGE:C5	2.17	0.57
1:A:55:ARG:NH2	1:A:63[A]:GLU:OE1	2.31	0.56
1:A:72:VAL:HG11	1:A:78:GLU:HB2	1.88	0.54
1:A:72:VAL:O	1:A:72:VAL:HG22	2.11	0.50
1:A:45:LEU:HD22	1:A:48[B]:MET:HE2	1.93	0.49
1:A:36:ARG:HB3	4:A:1221:PGE:H32	1.98	0.45
1:A:75:PRO:HG2	1:A:76:HIS:CE1	2.54	0.43
1:A:199[B]:GLN:NE2	1:B:201:PHE:CE1	2.87	0.42
1:A:39:TYR:OH	2:A:1000:V4E:H052	2.20	0.42
1:B:198:LEU:O	1:B:202:ARG:C	2.63	0.42
1:A:83:THR:O	3:A:1217:EDO:H12	2.19	0.42
1:B:9:GLU:HB3	1:B:159:GLU:HB3	2.01	0.41
1:A:41:ASP:OD2	1:A:48[B]:MET:HG3	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	207/226 (92%)	203 (98%)	4 (2%)	0	100	100
1	B	186/226 (82%)	182 (98%)	4 (2%)	0	100	100
All	All	393/452 (87%)	385 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	170/181 (94%)	168 (99%)	2 (1%)	63	72
1	B	145/181 (80%)	142 (98%)	3 (2%)	47	54
All	All	315/362 (87%)	310 (98%)	5 (2%)	55	64

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	GLU
1	A	81	GLU
1	B	9	GLU
1	B	98	SER
1	B	127	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	PGE	A	1220	-	9,9,9	0.63	0	8,8,8	0.48	0
3	EDO	A	1219	-	3,3,3	0.50	0	2,2,2	0.19	0
3	EDO	A	1218	-	3,3,3	0.46	0	2,2,2	0.21	0
2	V4E	A	1000	-	30,31,31	2.88	8 (26%)	43,47,47	3.96	15 (34%)
2	V4E	B	1000	-	30,31,31	2.94	8 (26%)	43,47,47	4.20	13 (30%)
3	EDO	A	1216	-	3,3,3	0.43	0	2,2,2	0.20	0
4	PGE	A	1221	-	9,9,9	0.81	0	8,8,8	1.27	1 (12%)
3	EDO	A	1217	-	3,3,3	0.44	0	2,2,2	0.06	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PGE	A	1220	-	-	4/7/7/7	-
3	EDO	A	1219	-	-	1/1/1/1	-
3	EDO	A	1218	-	-	1/1/1/1	-
2	V4E	A	1000	-	-	4/23/23/23	0/2/2/2
2	V4E	B	1000	-	-	1/23/23/23	0/2/2/2
3	EDO	A	1216	-	-	0/1/1/1	-
4	PGE	A	1221	-	-	5/7/7/7	-
3	EDO	A	1217	-	-	1/1/1/1	-

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1000	V4E	C02-N21	9.31	1.59	1.39
2	A	1000	V4E	C02-N21	8.94	1.58	1.39
2	B	1000	V4E	C03-S19	-7.62	1.52	1.72
2	A	1000	V4E	C03-S19	-7.40	1.53	1.72
2	A	1000	V4E	C20-N21	4.77	1.44	1.32
2	B	1000	V4E	C20-S19	-4.60	1.55	1.69

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1000	V4E	C20-N21	4.58	1.43	1.32
2	B	1000	V4E	C03-C02	4.56	1.44	1.35
2	A	1000	V4E	C03-C02	4.52	1.44	1.35
2	A	1000	V4E	C20-S19	-4.33	1.55	1.69
2	B	1000	V4E	C24-N25	3.64	1.43	1.34
2	A	1000	V4E	C24-N25	3.44	1.43	1.34
2	B	1000	V4E	C22-C23	2.65	1.56	1.51
2	A	1000	V4E	C22-C23	2.38	1.55	1.51
2	A	1000	V4E	P15-O18	-2.37	1.46	1.54
2	B	1000	V4E	C27-N29	2.18	1.37	1.34

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1000	V4E	C20-S19-C03	24.87	107.69	91.22
2	A	1000	V4E	C20-S19-C03	22.57	106.16	91.22
2	A	1000	V4E	C20-N21-C02	-4.84	107.52	114.06
2	B	1000	V4E	C20-N21-C02	-4.56	107.91	114.06
2	A	1000	V4E	N29-C27-N26	-3.97	118.92	125.53
2	B	1000	V4E	N29-C27-N26	-3.66	119.44	125.53
2	A	1000	V4E	O17-P15-O14	3.57	116.61	104.64
2	B	1000	V4E	C28-C27-N29	3.50	120.93	117.20
2	A	1000	V4E	C28-C27-N29	3.42	120.84	117.20
2	A	1000	V4E	C30-N29-C27	3.32	121.53	116.07
2	B	1000	V4E	S19-C20-N21	-3.27	108.15	112.30
2	A	1000	V4E	O12-P11-O14	-2.86	99.53	107.27
2	B	1000	V4E	C30-N29-C27	2.80	120.67	116.07
4	A	1221	PGE	O3-C4-C3	2.80	123.12	110.35
2	A	1000	V4E	C30-C23-C24	2.76	118.94	115.55
2	B	1000	V4E	C30-C23-C24	2.75	118.93	115.55
2	B	1000	V4E	C01-C02-N21	2.67	126.72	120.57
2	A	1000	V4E	C23-C30-N29	-2.66	119.50	123.83
2	A	1000	V4E	S19-C20-N21	-2.56	109.05	112.30
2	B	1000	V4E	C23-C30-N29	-2.43	119.89	123.83
2	A	1000	V4E	O18-P15-O16	-2.35	101.67	110.83
2	A	1000	V4E	O12-P11-O10	2.30	113.49	107.27
2	A	1000	V4E	C01-C02-N21	2.28	125.83	120.57
2	B	1000	V4E	O09-P07-O10	2.28	113.43	107.27
2	B	1000	V4E	C27-N26-C24	2.18	121.45	118.10
2	B	1000	V4E	O09-P07-O08	-2.13	102.52	112.44
2	B	1000	V4E	O17-P15-O14	2.12	111.76	104.64
2	A	1000	V4E	C28-C27-N26	2.08	120.24	117.13

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1000	V4E	C22-N21-C20	2.00	127.66	123.48

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1000	V4E	P11-O14-P15-O17
2	B	1000	V4E	C05-O06-P07-O08
4	A	1220	PGE	O2-C3-C4-O3
4	A	1221	PGE	O1-C1-C2-O2
4	A	1221	PGE	O3-C5-C6-O4
3	A	1217	EDO	O1-C1-C2-O2
2	A	1000	V4E	N21-C22-C23-C30
4	A	1220	PGE	O1-C1-C2-O2
2	A	1000	V4E	P11-O14-P15-O18
3	A	1218	EDO	O1-C1-C2-O2
4	A	1221	PGE	C1-C2-O2-C3
4	A	1220	PGE	O3-C5-C6-O4
4	A	1220	PGE	C1-C2-O2-C3
4	A	1221	PGE	O2-C3-C4-O3
3	A	1219	EDO	O1-C1-C2-O2
4	A	1221	PGE	C6-C5-O3-C4
2	A	1000	V4E	N21-C22-C23-C24


There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1000	V4E	1	0
4	A	1221	PGE	2	0
3	A	1217	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	203/226 (89%)	0.84	39 (19%) 3 3	13, 37, 84, 109	9 (4%)
1	B	192/226 (84%)	1.75	77 (40%) 0 1	27, 67, 106, 122	0
All	All	395/452 (87%)	1.28	116 (29%) 1 1	13, 53, 100, 122	9 (2%)

All (116) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	162	VAL	6.6
1	A	188	GLU	5.6
1	A	69	VAL	5.2
1	B	164	GLU	5.0
1	B	5	LEU	4.9
1	A	72	VAL	4.9
1	B	166	ALA	4.8
1	B	175	ILE	4.7
1	A	75	PRO	4.7
1	A	73	SER	4.5
1	A	76	HIS	4.5
1	A	71	GLY	4.5
1	B	25	LEU	4.4
1	B	130	LEU	4.3
1	B	180	SER	4.3
1	B	76	HIS	4.2
1	B	168	VAL	4.2
1	B	163	HIS	4.1
1	A	74	GLY	4.1
1	A	183	GLY	4.1
1	B	6	ILE	4.1
1	B	141	GLN	4.0
1	B	172	LEU	4.0
1	B	204	LEU	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	182	LEU	4.0
1	A	185	PRO	4.0
1	B	207	GLN	3.9
1	A	180	SER	3.9
1	A	161	MET	3.8
1	B	8	VAL	3.7
1	B	178	VAL	3.7
1	B	203	PRO	3.7
1	A	184	VAL	3.7
1	A	70	THR	3.7
1	A	186	ALA	3.6
1	B	171	ALA	3.6
1	B	206	TYR	3.6
1	B	186	ALA	3.5
1	B	18	THR	3.4
1	B	182	LEU	3.4
1	A	4	GLY	3.3
1	B	212	ALA	3.3
1	B	131	ALA	3.3
1	B	199	GLN	3.2
1	B	209	LEU	3.2
1	B	210	LEU	3.1
1	B	12	PHE	3.1
1	B	165	LYS	3.1
1	B	161	MET	3.1
1	A	77	ASN	3.1
1	A	215	SER	3.1
1	B	160	ALA	3.1
1	B	190	ALA	3.1
1	B	158	VAL	3.1
1	A	5	LEU	3.1
1	A	10	ARG	3.0
1	B	21	ARG	3.0
1	B	142	LEU	3.0
1	B	208	ARG	2.9
1	B	24	GLU	2.9
1	B	184	VAL	2.9
1	A	129	LYS	2.9
1	B	17	ASP	2.9
1	B	185	PRO	2.9
1	B	198	LEU	2.9
1	B	169	PRO	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	189	GLU	2.9
1	A	174	LYS	2.9
1	A	214	SER	2.9
1	B	176	ILE	2.8
1	B	16	PRO	2.8
1	A	164	GLU	2.8
1	A	6	ILE	2.8
1	B	7	GLU	2.8
1	B	128	TRP	2.8
1	B	187	GLN	2.8
1	B	129	LYS	2.8
1	B	27	ALA	2.7
1	B	77	ASN	2.7
1	A	25	LEU	2.7
1	B	14	PRO	2.6
1	B	4	GLY	2.6
1	A	187	GLN	2.6
1	B	44	GLU	2.6
1	A	176	ILE	2.5
1	B	22	LEU	2.5
1	A	166	ALA	2.4
1	B	49	LEU	2.4
1	B	196	VAL	2.4
1	B	31	HIS	2.4
1	B	15	GLY	2.4
1	A	15	GLY	2.3
1	A	163	HIS	2.3
1	B	201	PHE	2.2
1	B	174	LYS	2.2
1	B	191	PRO	2.2
1	B	173	GLU	2.2
1	B	29	LEU	2.2
1	B	45	LEU	2.2
1	B	197	TYR	2.2
1	A	181	MET	2.2
1	B	126	SER	2.2
1	B	78	GLU	2.1
1	B	159	GLU	2.1
1	B	189	GLU	2.1
1	A	179	SER	2.1
1	B	170	ALA	2.1
1	A	162	VAL	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	13	ALA	2.1
1	A	178	VAL	2.1
1	A	131	ALA	2.1
1	A	160	ALA	2.1
1	B	192	ALA	2.1
1	B	143	THR	2.0
1	B	188	GLU	2.0
1	B	195	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	PGE	A	1220	10/10	0.78	0.24	54,59,60,61	0
4	PGE	A	1221	10/10	0.81	0.18	36,39,40,41	0
3	EDO	A	1219	4/4	0.83	0.21	56,57,58,59	0
3	EDO	A	1218	4/4	0.90	0.18	47,50,51,52	0
2	V4E	B	1000	30/30	0.90	0.16	43,63,86,88	0
3	EDO	A	1216	4/4	0.92	0.12	40,41,41,42	0
3	EDO	A	1217	4/4	0.92	0.14	36,41,41,45	0
2	V4E	A	1000	30/30	0.96	0.11	30,42,62,65	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around V4E B 1000:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around V4E A 1000:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.