



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 13, 2026 – 05:15 AM UTC

PDB ID : 5A6T / pdb_00005a6t
Title : 1.65 Å resolution Sulphite inhibited *Sporosarcina pasteurii* urease
Authors : Mazzei, L.; Cianci, M.; Benini, S.; Bertini, L.; Musiani, F.; Ciurli, S.
Deposited on : 2015-07-01
Resolution : 1.65 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.66-1.66)
Clashscore	190562	2662 (1.66-1.66)
Ramachandran outliers	187476	2621 (1.66-1.66)
Sidechain outliers	187428	2621 (1.66-1.66)
RSRZ outliers	180081	2564 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	100	 96%
2	B	126	 2% 92% 5%
3	C	570	 0% 93% 6%

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 6863 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UREASE SUBUNIT GAMMA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	100	Total	C	N	O	S	0	2	0
			791	499	133	152	7			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	20	ALA	LEU	conflict	UNP P41022
A	22	LYS	ARG	conflict	UNP P41022

- Molecule 2 is a protein called UREASE SUBUNIT BETA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	122	Total	C	N	O	S	0	1	0
			960	594	172	193	1			

- Molecule 3 is a protein called UREASE SUBUNIT ALPHA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	570	Total	C	N	O	S	0	17	0
			4438	2786	762	864	26			

There are 10 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	19	GLN	ARG	conflict	UNP P41020
C	28	TRP	GLY	conflict	UNP P41020
C	29	ILE	-	insertion	UNP P41020
C	36	THR	TYR	conflict	UNP P41020
C	37	THR	TYR	conflict	UNP P41020
C	38	TYR	LEU	conflict	UNP P41020
C	42	ALA	VAL	conflict	UNP P41020
C	263	LEU	VAL	conflict	UNP P41020

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	403	ALA	LEU	conflict	UNP P41020
C	420	ILE	MET	conflict	UNP P41020

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is NICKEL (II) ION (CCD ID: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	C	2	Total	Ni	0	0
			2	2		

- Molecule 7 is SULFITE ION (CCD ID: SO3) (formula: O₃S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	C	1	Total	O	S	0	0
			4	3	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	97	Total	O	0	0
			97	97		
8	B	139	Total	O	0	0
			139	139		
8	C	399	Total	O	0	0
			399	399		

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: UREASE SUBUNIT GAMMA

Chain A:  96% .




• Molecule 2: UREASE SUBUNIT BETA

Chain B:  2% 92% 5% .



• Molecule 3: UREASE SUBUNIT ALPHA

Chain C:  % 93% 6% .



4 Data and refinement statistics

Property	Value	Source
Space group	P 63 2 2	Depositor
Cell constants a, b, c, α , β , γ	131.26Å 131.26Å 189.07Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	113.68 – 1.65 113.68 – 1.65	Depositor EDS
% Data completeness (in resolution range)	99.7 (113.68-1.65) 99.7 (113.68-1.65)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.87 (at 1.65Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.129 , 0.152 (Not available) , 0.157	Depositor DCC
R_{free} test set	5779 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	17.2	Xtriage
Anisotropy	0.279	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 37.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6863	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CXM, SO4, KCX, SO3, NI, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.34	2/794 (0.3%)	1.16	1/1068 (0.1%)
2	B	1.33	1/972 (0.1%)	1.29	1/1308 (0.1%)
3	C	1.27	8/4531 (0.2%)	1.15	7/6136 (0.1%)
All	All	1.29	11/6297 (0.2%)	1.17	9/8512 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	C	0	1

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	520	CYS	CB-SG	-9.34	1.50	1.81
3	C	520	CYS	CA-CB	8.63	1.66	1.54
1	A	9	GLU	CA-C	-5.85	1.45	1.52
3	C	107	ASP	N-CA	-5.57	1.38	1.45
3	C	531	ASN	CA-CB	5.56	1.57	1.53
3	C	5	ARG	NE-CZ	-5.51	1.26	1.33
3	C	328	ASN	CA-C	5.28	1.59	1.52
3	C	453	GLY	C-O	-5.24	1.18	1.24
2	B	33	SER	CB-OG	-5.17	1.31	1.42
1	A	78	ASP	CG-OD1	-5.11	1.15	1.25
3	C	460	GLN	N-CA	5.08	1.52	1.46

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	302	ASN	CA-C-N	-7.09	112.40	119.56
3	C	302	ASN	C-N-CA	-7.09	112.40	119.56
3	C	5	ARG	NE-CZ-NH2	-6.81	113.07	119.20
1	A	97	ASN	CA-C-O	6.27	125.28	120.26
3	C	322	CYS	CB-CA-C	-5.34	101.60	110.68
3	C	8	TYR	N-CA-C	-5.33	105.37	111.07
3	C	1	MET	CG-SD-CE	-5.16	89.54	100.90
3	C	329	ILE	N-CA-C	-5.15	97.76	108.88
2	B	125	VAL	O-C-N	5.03	128.46	122.83

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	C	234	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	791	0	811	0	0
2	B	960	0	942	3	0
3	C	4438	0	4419	19	0
4	A	8	0	12	0	0
4	B	4	0	6	0	0
4	C	16	0	24	0	0
5	B	5	0	0	0	0
6	C	2	0	0	0	0
7	C	4	0	0	0	0
8	A	97	0	0	0	1
8	B	139	0	0	0	1
8	C	399	0	0	3	0
All	All	6863	0	6214	19	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:272:HIS:HE1	3:C:299:SER:OG	1.75	0.70
3:C:507[A]:LYS:HD2	8:C:2170:HOH:O	1.91	0.69
3:C:270[B]:VAL:HG12	8:C:2359:HOH:O	2.00	0.61
3:C:270[B]:VAL:HG11	3:C:523:ILE:O	2.01	0.61
3:C:272:HIS:HD2	3:C:418[B]:GLN:NE2	1.98	0.60
3:C:293:HIS:HE1	8:C:2245:HOH:O	1.86	0.59
3:C:522[B]:ASN:OD1	3:C:522[B]:ASN:N	2.41	0.53
2:B:95:GLU:O	3:C:104:ASP:HB3	2.11	0.51
3:C:132:GLY:HA3	3:C:155:THR:OG1	2.14	0.48
3:C:272:HIS:CE1	3:C:299:SER:OG	2.63	0.47
3:C:272:HIS:HD2	3:C:418[B]:GLN:HE21	1.62	0.47
3:C:316:LEU:O	3:C:320:MET:HG2	2.16	0.46
3:C:70:LEU:HD11	3:C:86:ASP:HB3	1.98	0.46
3:C:274:PHE:O	3:C:275:HIS:C	2.61	0.43
2:B:71:SER:OG	3:C:49:VAL:HG21	2.19	0.43
3:C:507[A]:LYS:HE2	3:C:507[A]:LYS:HB3	1.80	0.43
3:C:137:HIS:CE1	3:C:274:PHE:CD2	3.08	0.42
3:C:362:THR:O	3:C:368:GLY:HA3	2.21	0.40
2:B:15:ALA:O	3:C:5:ARG:HD2	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:A:2015:HOH:O	8:B:2049:HOH:O[8_676]	1.96	0.24

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	100/100 (100%)	100 (100%)	0	0	100	100
2	B	121/126 (96%)	118 (98%)	2 (2%)	1 (1%)	16	4

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	C	584/570 (102%)	561 (96%)	23 (4%)	0	100	100
All	All	805/796 (101%)	779 (97%)	25 (3%)	1 (0%)	48	30

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	99	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	85/83 (102%)	85 (100%)	0	100	100
2	B	102/105 (97%)	102 (100%)	0	100	100
3	C	475/458 (104%)	468 (98%)	7 (2%)	57	37
All	All	662/646 (102%)	655 (99%)	7 (1%)	65	48

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	1	MET
3	C	158	PHE
3	C	272	HIS
3	C	395	LYS
3	C	402	ARG
3	C	452	LYS
3	C	551	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	96	HIS
2	B	5	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	C	65	ASN
3	C	179	ASN
3	C	254	ASN
3	C	272	HIS
3	C	293	HIS
3	C	323	HIS
3	C	324	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	KCX	C	220	3,6	10,11,12	1.74	1 (10%)	6,12,14	3.53	1 (16%)
1	CXM	A	1	1	9,10,11	2.57	1 (11%)	9,11,13	1.20	1 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	KCX	C	220	3,6	-	0/9/10/12	-
1	CXM	A	1	1	-	3/9/10/12	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1	CXM	ON1-CN	7.32	1.35	1.21
3	C	220	KCX	OQ1-CX	4.94	1.30	1.21

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	220	KCX	OQ1-CX-NZ	-8.56	111.92	124.92
1	A	1	CXM	ON1-CN-N	-2.18	121.28	124.86

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	1	CXM	O-C-CA-CB
1	A	1	CXM	CB-CA-N-CN
1	A	1	CXM	C-CA-N-CN

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 2 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	EDO	C	1577	-	3,3,3	0.36	0	2,2,2	0.29	0
7	SO3	C	1573	6	1,3,3	5.45	1 (100%)	0,3,3	-	-
4	EDO	A	1102	-	3,3,3	1.58	0	2,2,2	0.52	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EDO	B	1127	-	3,3,3	0.21	0	2,2,2	0.91	0
4	EDO	C	1575	-	3,3,3	0.79	0	2,2,2	0.75	0
4	EDO	C	1574	-	3,3,3	0.50	0	2,2,2	0.36	0
4	EDO	A	1101	-	3,3,3	0.71	0	2,2,2	0.94	0
4	EDO	C	1576	-	3,3,3	0.51	0	2,2,2	0.25	0
5	SO4	B	1128	-	4,4,4	0.56	0	6,6,6	0.94	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	C	1577	-	-	0/1/1/1	-
4	EDO	A	1102	-	-	0/1/1/1	-
4	EDO	B	1127	-	-	1/1/1/1	-
4	EDO	C	1575	-	-	0/1/1/1	-
4	EDO	C	1574	-	-	0/1/1/1	-
4	EDO	A	1101	-	-	0/1/1/1	-
4	EDO	C	1576	-	-	0/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	C	1573	SO3	O1-S	5.45	1.67	1.44

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	1127	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	99/100 (99%)	-0.60	0 100 100	9, 16, 23, 26	2 (2%)
2	B	122/126 (96%)	-0.26	2 (1%) 70 75	11, 18, 33, 52	1 (0%)
3	C	569/570 (99%)	-0.50	8 (1%) 73 78	7, 15, 28, 80	17 (2%)
All	All	790/796 (99%)	-0.48	10 (1%) 75 80	7, 16, 28, 80	20 (2%)

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	390	PRO	5.2
3	C	391	LEU	4.9
3	C	396	ASN	4.1
3	C	329	ILE	4.1
3	C	392	ALA	4.0
3	C	328	ASN	2.8
3	C	397	GLY	2.8
3	C	395	LYS	2.4
2	B	5	ASN	2.3
2	B	126	GLU	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	KCX	C	220	12/13	0.98	0.04	11,13,15,15	0
1	CXM	A	1	11/12	0.99	0.04	14,16,19,22	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	SO4	B	1128	5/5	0.87	0.10	42,48,57,66	0
4	EDO	C	1576	4/4	0.90	0.12	41,41,41,43	0
4	EDO	C	1577	4/4	0.93	0.10	32,35,36,46	0
4	EDO	B	1127	4/4	0.93	0.12	33,40,43,52	0
4	EDO	A	1102	4/4	0.94	0.11	23,24,26,28	0
4	EDO	C	1574	4/4	0.96	0.07	27,30,30,31	0
4	EDO	C	1575	4/4	0.97	0.07	26,29,32,33	0
4	EDO	A	1101	4/4	0.97	0.06	20,24,24,25	0
7	SO3	C	1573	4/4	0.97	0.10	22,23,25,28	0
6	NI	C	1572	1/1	1.00	0.04	16,16,16,16	0
6	NI	C	1571	1/1	1.00	0.05	18,18,18,18	0

6.5 Other polymers [i](#)

There are no such residues in this entry.