



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 12:53 AM UTC

PDB ID : 6A60 / pdb_00006a60
Title : Crystal structure of human FXR/RXR-LBD heterodimer bound to GW4064 and 9cRA and SRC1
Authors : Wang, N.; Liu, J.
Deposited on : 2018-06-25
Resolution : 3.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2469 (3.10-3.02)
Clashscore	190562	2569 (3.10-3.02)
Ramachandran outliers	187476	2424 (3.10-3.02)
Sidechain outliers	187428	2423 (3.10-3.02)
RSRZ outliers	180081	2469 (3.10-3.02)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	229	
2	B	16	
2	F	16	
3	D	238	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3697 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bile acid receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	219	Total	C	N	O	S	0	0	0
			1808	1157	302	340	9			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	432	GLU	CYS	engineered mutation	UNP Q96RI1
A	466	GLU	CYS	engineered mutation	UNP Q96RI1


- Molecule 2 is a protein called Nuclear receptor coactivator 1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	8	Total	C	N	O	0	0	0
			71	47	15	9			
2	F	11	Total	C	N	O	0	0	0
			101	64	23	14			

- Molecule 3 is a protein called Retinoic acid receptor RXR-alpha.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	210	Total	C	N	O	S	0	0	0
			1658	1065	284	299	10			

- Molecule 4 is 3-[(E)-2-(2-chloro-4-{[3-(2,6-dichlorophenyl)-5-(1-methylethyl)isoxazol-4-yl]methoxy}phenyl)ethenyl]benzoic acid (CCD ID: 064) (formula: C₂₈H₂₂Cl₃NO₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	Cl	N	O	
			36	28	3	1	4	

- Molecule 5 is (9cis)-retinoic acid (CCD ID: 9CR) (formula: $C_{20}H_{28}O_2$).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	D	1	Total	C	O	
			22	20	2	


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	O	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Bile acid receptor




- Molecule 2: Nuclear receptor coactivator 1



- Molecule 2: Nuclear receptor coactivator 1



- Molecule 3: Retinoic acid receptor RXR-alpha



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	102.85Å 102.85Å 109.46Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	74.95 – 3.05 74.95 – 3.05	Depositor EDS
% Data completeness (in resolution range)	98.5 (74.95-3.05) 98.4 (74.95-3.05)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.33 (at 3.07Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.225 , 0.288 0.225 , 0.293	Depositor DCC
R_{free} test set	528 reflections (4.51%)	wwPDB-VP
Wilson B-factor (Å ²)	106.2	Xtriage
Anisotropy	0.100	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 74.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3697	wwPDB-VP
Average B, all atoms (Å ²)	128.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 9CR, 064

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.99	4/1844 (0.2%)	1.19	3/2488 (0.1%)
2	B	0.69	0/71	0.97	0/93
2	F	1.18	2/102 (2.0%)	1.35	1/134 (0.7%)
3	D	1.03	2/1690 (0.1%)	1.15	3/2284 (0.1%)
All	All	1.01	8/3707 (0.2%)	1.18	7/4999 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
2	F	0	1
3	D	0	1
All	All	0	3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	443	PHE	C-O	-5.96	1.16	1.24
1	A	373	GLY	C-O	-5.76	1.17	1.23
1	A	257	MET	C-O	-5.67	1.16	1.24
2	F	638	GLN	N-CA	5.25	1.52	1.46
3	D	232	VAL	C-O	-5.22	1.18	1.24
3	D	342	VAL	N-CA	5.15	1.52	1.46
1	A	320	LEU	C-O	-5.14	1.18	1.24
2	F	634	HIS	C-O	5.11	1.29	1.24

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	444	ASP	CA-CB-CG	6.04	118.64	112.60
1	A	377	MET	N-CA-C	5.86	118.55	111.40
1	A	249	GLN	CB-CA-C	-5.77	108.28	115.89
1	A	393	PRO	N-CA-C	5.30	120.54	114.03
3	D	289	PHE	N-CA-C	-5.25	105.47	111.14
3	D	436	LEU	CA-C-O	-5.18	114.93	120.42
2	F	638	GLN	N-CA-C	5.05	119.69	111.37

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	436	ARG	Sidechain
3	D	414	ARG	Sidechain
2	F	635	ARG	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1808	0	1799	38	0
2	B	71	0	84	2	0
2	F	101	0	110	1	0
3	D	1658	0	1695	19	0
4	A	36	0	21	2	0
5	D	22	0	27	4	0
6	A	1	0	0	0	0
All	All	3697	0	3736	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:377:MET:O	1:A:378:THR:HG22	1.29	1.25

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:377:MET:O	1:A:378:THR:CG2	1.96	1.14
1:A:365:MET:HE1	1:A:450:MET:HE1	1.30	1.07
1:A:377:MET:O	1:A:381:GLU:HB2	1.85	0.76
1:A:246:THR:O	1:A:249:GLN:HG3	1.88	0.74
1:A:365:MET:HE1	1:A:450:MET:CE	2.18	0.65
1:A:248:ASP:OD1	1:A:251:THR:OG1	2.15	0.65
1:A:377:MET:C	1:A:378:THR:HG22	2.20	0.65
2:F:631:LYS:O	2:F:635:ARG:HD3	1.98	0.62
3:D:347:ASP:O	3:D:351:THR:HG23	2.01	0.60
1:A:365:MET:CE	1:A:450:MET:HE1	2.20	0.59
1:A:465:LEU:HA	1:A:468:ILE:CG2	2.33	0.58
1:A:464:LEU:O	1:A:468:ILE:HG22	2.04	0.57
1:A:465:LEU:HA	1:A:468:ILE:HG22	1.88	0.56
2:B:748:ILE:HG22	2:B:748:ILE:O	2.06	0.55
1:A:442:THR:OG1	3:D:426:ARG:NH1	2.39	0.55
1:A:379:GLN:O	1:A:380:GLU:C	2.51	0.53
1:A:269:ILE:HG23	1:A:269:ILE:O	2.08	0.53
1:A:363:THR:N	1:A:364:PRO:HD2	2.24	0.53
1:A:286:ILE:HD13	1:A:351:ARG:NH1	2.23	0.52
1:A:246:THR:OG1	1:A:249:GLN:HA	2.09	0.51
1:A:295:VAL:O	1:A:299:VAL:HG23	2.10	0.51
3:D:327:ALA:HB2	5:D:501:9CR:O2	2.10	0.51
4:A:501:064:H27	4:A:501:064:CL25	2.46	0.51
1:A:434:LEU:O	1:A:437:LEU:HB2	2.11	0.51
3:D:414:ARG:HG3	3:D:414:ARG:HH11	1.76	0.51
1:A:258:ASP:HA	1:A:261:ASN:HB2	1.93	0.50
1:A:264:ARG:CG	1:A:266:PRO:HD3	2.43	0.49
3:D:442:ILE:HD11	3:D:444:ASP:OD1	2.13	0.49
3:D:442:ILE:HD11	3:D:444:ASP:CG	2.38	0.48
1:A:250:GLN:HE21	1:A:250:GLN:N	2.10	0.48
3:D:268:ILE:HG23	5:D:501:9CR:C20	2.44	0.48
1:A:260:TYR:CD2	1:A:383:ALA:HB2	2.51	0.46
1:A:264:ARG:HG2	1:A:266:PRO:HD3	1.97	0.46
1:A:246:THR:O	1:A:249:GLN:CG	2.61	0.46
1:A:377:MET:HB2	1:A:382:TYR:CE2	2.50	0.46
3:D:307:GLU:OE2	3:D:426:ARG:HD3	2.15	0.46
3:D:365:THR:HG21	3:D:403:TYR:CE1	2.51	0.45
3:D:286:ILE:O	3:D:287:PRO:C	2.59	0.45
1:A:291:ALA:O	1:A:295:VAL:HG23	2.18	0.45
3:D:265:VAL:HG12	3:D:445:THR:HG21	2.00	0.44
1:A:436:ARG:HG2	1:A:436:ARG:HH11	1.83	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:245:LEU:HD22	1:A:250:GLN:NE2	2.33	0.43
3:D:360:MET:HE3	3:D:417:LYS:CE	2.48	0.43
3:D:447:ILE:HG23	3:D:451:LEU:HD23	1.99	0.43
5:D:501:9CR:H27	5:D:501:9CR:H9	2.01	0.43
1:A:267:GLN:HG2	1:A:270:THR:HG23	2.00	0.43
1:A:328:MET:SD	4:A:501:064:CL15	3.13	0.43
1:A:303:LYS:HG2	1:A:309:GLN:HE21	1.83	0.43
1:A:347:LEU:HD13	1:A:350:GLU:CG	2.49	0.43
3:D:330:LEU:HD12	3:D:330:LEU:N	2.33	0.42
3:D:450:PHE:CE1	3:D:454:MET:HE2	2.53	0.42
1:A:449:GLU:O	1:A:450:MET:C	2.63	0.42
3:D:350:LEU:HA	3:D:350:LEU:HD23	1.84	0.42
3:D:306:ASN:ND2	3:D:433:LEU:HG	2.35	0.42
3:D:306:ASN:HD22	3:D:433:LEU:HG	1.84	0.41
1:A:340:LEU:HD13	1:A:340:LEU:HA	1.96	0.41
3:D:432:CYS:HB3	5:D:501:9CR:H7	2.03	0.41
1:A:253:LEU:C	1:A:253:LEU:HD12	2.46	0.40
2:B:748:ILE:O	2:B:748:ILE:CG2	2.68	0.40
1:A:248:ASP:OD2	1:A:252:LEU:CB	2.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	213/229 (93%)	192 (90%)	20 (9%)	1 (0%)	24	52
2	B	6/16 (38%)	5 (83%)	1 (17%)	0	100	100
2	F	9/16 (56%)	8 (89%)	1 (11%)	0	100	100
3	D	206/238 (87%)	198 (96%)	7 (3%)	1 (0%)	24	52
All	All	434/499 (87%)	403 (93%)	29 (7%)	2 (0%)	24	52

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	D	322	ASP
1	A	269	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	203/212 (96%)	178 (88%)	25 (12%)	4	16
2	B	8/15 (53%)	3 (38%)	5 (62%)	0	0
2	F	11/15 (73%)	9 (82%)	2 (18%)	2	7
3	D	180/205 (88%)	167 (93%)	13 (7%)	13	37
All	All	402/447 (90%)	357 (89%)	45 (11%)	6	20

All (45) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	246	THR
1	A	248	ASP
1	A	249	GLN
1	A	250	GLN
1	A	253	LEU
1	A	263	GLN
1	A	267	GLN
1	A	269	ILE
1	A	293	ASN
1	A	325	VAL
1	A	331	ARG
1	A	340	LEU
1	A	347	LEU
1	A	360	GLU
1	A	367	SER
1	A	370	LYS
1	A	379	GLN
1	A	381	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	399	LYS
1	A	418	LEU
1	A	425	GLU
1	A	456	VAL
1	A	459	HIS
1	A	463	PRO
1	A	468	ILE
2	B	747	LYS
2	B	749	LEU
2	B	750	HIS
2	B	751	ARG
2	B	753	LEU
3	D	278	THR
3	D	288	HIS
3	D	290	SER
3	D	320	VAL
3	D	324	ILE
3	D	336	SER
3	D	345	ILE
3	D	351	THR
3	D	360	MET
3	D	407	LYS
3	D	431	LYS
3	D	442	ILE
3	D	445	THR
2	F	629	ARG
2	F	635	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	250	GLN
1	A	296	GLN
1	A	309	GLN
1	A	337	ASN
1	A	396	GLN
1	A	444	ASN
3	D	270	GLN
3	D	275	GLN
3	D	331	HIS
3	D	333	HIS
3	D	377	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	D	435	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	064	A	501	-	39,39,39	3.03	10 (25%)	51,55,55	2.49	17 (33%)
5	9CR	D	501	-	22,22,22	1.72	6 (27%)	30,30,30	1.41	6 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	064	A	501	-	-	7/22/22/22	0/4/4/4
5	9CR	D	501	-	-	10/15/32/32	0/1/1/1

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	501	064	C2-C5	11.92	1.48	1.34
4	A	501	064	C24-C22	9.48	1.50	1.39
4	A	501	064	C3-C8	4.89	1.47	1.39
4	A	501	064	C3-C7	4.85	1.46	1.39
4	A	501	064	O9-N4	-4.27	1.34	1.42
5	D	501	9CR	C14-C13	3.79	1.40	1.35
4	A	501	064	C8-CL15	3.34	1.81	1.73
5	D	501	9CR	C11-C12	2.92	1.42	1.34
5	D	501	9CR	C10-C9	2.88	1.42	1.35
4	A	501	064	C7-CL13	2.82	1.80	1.73
5	D	501	9CR	C8-C9	-2.74	1.40	1.46
5	D	501	9CR	O2-C15	-2.58	1.23	1.30
4	A	501	064	O9-C5	2.43	1.37	1.34
4	A	501	064	C1-N4	2.39	1.35	1.31
4	A	501	064	C22-CL25	2.38	1.79	1.73
5	D	501	9CR	C12-C13	-2.17	1.41	1.46

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	501	064	C22-C24-C26	7.43	128.57	121.87
4	A	501	064	C24-C22-CL25	6.38	125.69	120.17
4	A	501	064	C5-O9-N4	5.56	114.01	109.45
4	A	501	064	C3-C1-N4	4.82	126.53	119.67
4	A	501	064	O9-C5-C2	-4.49	105.85	110.15
4	A	501	064	C20-C22-C24	-3.75	119.63	122.45
4	A	501	064	C23-C24-C26	-3.32	114.50	121.13
5	D	501	9CR	C10-C11-C12	3.30	132.76	123.20
4	A	501	064	C6-O11-C18	3.28	123.51	117.63
4	A	501	064	C12-C7-C3	-3.23	117.54	121.86
5	D	501	9CR	C12-C13-C14	3.19	127.99	119.15
4	A	501	064	C12-C7-CL13	3.15	124.62	118.42
4	A	501	064	C8-C3-C1	-3.09	118.27	121.58
4	A	501	064	C14-C8-CL15	3.06	124.45	118.42
4	A	501	064	C3-C1-C2	-3.03	122.80	129.54
4	A	501	064	C14-C8-C3	-3.01	117.83	121.86
4	A	501	064	C8-C3-C7	2.91	120.77	116.70
5	D	501	9CR	C8-C9-C10	2.60	123.10	119.01
5	D	501	9CR	C20-C13-C14	-2.38	115.70	122.84
4	A	501	064	C20-C22-CL25	-2.31	114.67	118.45
5	D	501	9CR	C11-C10-C9	2.31	130.51	127.28
5	D	501	9CR	C19-C9-C10	-2.21	119.23	122.82

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	501	064	O9-C5-C10	2.21	119.62	116.87

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	501	064	C22-C24-C26-C27
5	D	501	9CR	C5-C6-C7-C8
5	D	501	9CR	C6-C7-C8-C9
5	D	501	9CR	C10-C11-C12-C13
5	D	501	9CR	C12-C13-C14-C15
5	D	501	9CR	C20-C13-C14-C15
4	A	501	064	C29-C31-C34-O35
4	A	501	064	C33-C31-C34-O35
4	A	501	064	C33-C31-C34-O36
4	A	501	064	C29-C31-C34-O36
5	D	501	9CR	C7-C8-C9-C19
5	D	501	9CR	C7-C8-C9-C10
5	D	501	9CR	C1-C6-C7-C8
5	D	501	9CR	C11-C10-C9-C8
4	A	501	064	C23-C24-C26-C27
5	D	501	9CR	C11-C10-C9-C19
4	A	501	064	C1-C2-C6-O11


There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	501	064	2	0
5	D	501	9CR	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	219/229 (95%)	-0.33	2 (0%) 81 61	91, 152, 202, 266	0
2	B	8/16 (50%)	0.04	1 (12%) 8 4	147, 179, 193, 210	0
2	F	11/16 (68%)	-0.49	0 100 100	75, 85, 145, 161	0
3	D	210/238 (88%)	-0.64	0 100 100	69, 97, 134, 164	0
All	All	448/499 (89%)	-0.47	3 (0%) 84 66	69, 122, 190, 266	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	320	LEU	2.9
1	A	349	GLU	2.2
2	B	752	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	064	A	501	36/36	0.88	0.10	109,133,174,182	0
5	9CR	D	501	22/22	0.96	0.08	72,85,91,94	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 064 A 501:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 9CR D 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.