



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:14 PM UTC

PDB ID : 7A6T / pdb_00007a6t
Title : Crystal Structure of Asn173Ser variant of Human Deoxyhypusine Synthase in complex with NAD and spermidine
Authors : Wator, E.; Wilk, P.; Grudnik, P.
Deposited on : 2020-08-26
Resolution : 1.66 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.66 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.66-1.66)
Clashscore	190562	2662 (1.66-1.66)
Ramachandran outliers	187476	2621 (1.66-1.66)
Sidechain outliers	187428	2621 (1.66-1.66)
RSRZ outliers	180081	2564 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	369	4% 84% 7% 9%
1	B	369	3% 85% • 11%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SPD	B	1602	-	-	X	-
5	BME	A	406	-	-	-	X

2 Entry composition ⓘ

There are 7 unique types of molecules in this entry. The entry contains 11380 atoms, of which 5500 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Deoxyhypusine synthase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	336	5381	1720	2685	456	503	17	0	17	0
1	B	328	5343	1705	2681	446	494	17	0	20	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	173	SER	ASN	engineered mutation	UNP P49366
B	173	SER	ASN	engineered mutation	UNP P49366


- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (CCD ID: NAD) (formula: C₂₁H₂₇N₇O₁₄P₂).



Continued from previous page...

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	B	1	Total	C	H	N	O	P	0	0
			70	21	26	7	14	2		

- Molecule 3 is SPERMIDINE (CCD ID: SPD) (formula: $C_7H_{19}N_3$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	N	0	0
			29	7	19	3		
3	B	1	Total	C	H	N	0	0
			29	7	19	3		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			10	2	6	2		
4	A	1	Total	C	H	O	0	0
			10	2	6	2		
4	A	1	Total	C	H	O	0	0
			10	2	6	2		
4	A	1	Total	C	H	O	0	0
			10	2	6	2		
4	B	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 5 is BETA-MERCAPTOETHANOL (CCD ID: BME) (formula: C_2H_6OS).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	H	O	S	0	0
			10	2	6	1	1		

- Molecule 6 is OXAMIC ACID (CCD ID: OXM) (formula: $C_2H_3NO_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	B	1	Total	C	H	N	O	0	0
			8	2	2	1	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	191	Total 191	O 191	0	0
7	B	189	Total 189	O 189	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Deoxyhypusine synthase



- Molecule 1: Deoxyhypusine synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	105.03Å 105.03Å 160.00Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.01 – 1.66 46.01 – 1.66	Depositor EDS
% Data completeness (in resolution range)	99.7 (46.01-1.66) 99.9 (46.01-1.66)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.02 (at 1.66Å)	Xtriage
Refinement program	PHENIX 1.17.1	Depositor
R, R_{free}	0.171 , 0.193 0.171 , 0.193	Depositor DCC
R_{free} test set	2099 reflections (1.74%)	wwPDB-VP
Wilson B-factor (Å ²)	25.7	Xtriage
Anisotropy	0.538	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 53.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.017 for -h,-k,l	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	11380	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BME, NAD, SPD, EDO, CSS, OXM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.61	4/2798 (0.1%)	0.75	8/3792 (0.2%)
1	B	0.52	2/2770 (0.1%)	0.69	2/3754 (0.1%)
All	All	0.57	6/5568 (0.1%)	0.72	10/7546 (0.1%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	284	GLY	C-N	-15.15	1.15	1.33
1	A	285	VAL	CA-C	12.95	1.68	1.52
1	B	40	VAL	CB-CG1	-7.12	1.29	1.52
1	A	173	SER	C-O	-6.08	1.16	1.24
1	B	114	GLU	C-O	-5.69	1.17	1.24
1	A	43	ARG	C-O	-5.41	1.17	1.24

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	284	GLY	CA-C-N	12.49	136.30	120.56
1	A	284	GLY	C-N-CA	12.49	136.30	120.56
1	B	40	VAL	CA-CB-CG1	-9.92	93.54	110.40
1	A	285	VAL	N-CA-C	-8.36	102.40	110.42
1	A	285	VAL	N-CA-CB	7.81	119.68	110.55
1	A	285	VAL	O-C-N	7.19	128.96	121.91
1	A	284	GLY	O-C-N	-6.00	116.07	122.67
1	B	42	TYR	N-CA-C	5.45	116.90	111.07
1	A	285	VAL	CA-C-N	-5.00	113.61	120.46
1	A	285	VAL	C-N-CA	-5.00	113.61	120.46

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2696	2685	2645	22	0
1	B	2662	2681	2641	16	0
2	A	44	26	26	2	0
2	B	44	26	26	0	0
3	A	10	19	19	4	0
3	B	10	19	19	8	0
4	A	20	30	30	5	0
4	B	4	6	6	0	0
5	A	4	6	6	3	0
6	B	6	2	2	0	0
7	A	191	0	0	3	0
7	B	189	0	0	5	0
All	All	5880	5500	5420	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:288:HIS:NE2	3:A:402:SPD:H51	1.84	0.93
1:A:320:ARG:HD2	4:A:405:EDO:H21	1.55	0.89
1:B:329:LYS:HZ1	3:B:1602:SPD:H51	1.47	0.79
1:A:329:LYS:NZ	3:A:402:SPD:H52	2.05	0.72
1:A:261:GLU:OE2	5:A:406:BME:S2	2.44	0.71
1:A:43:ARG:NH1	1:A:357:GLN:CD	2.53	0.67
1:B:137[B]:GLU:OE1	7:B:1701:HOH:O	2.12	0.66
1:A:329:LYS:HZ3	3:A:402:SPD:H52	1.61	0.65
1:B:288:HIS:NE2	3:B:1602:SPD:H52	2.16	0.60
1:A:320:ARG:HH11	4:A:405:EDO:H21	1.67	0.60
1:B:329:LYS:NZ	3:B:1602:SPD:H51	2.19	0.55
1:B:358:LYS:HG3	7:B:1863:HOH:O	2.06	0.55
1:A:77:LEU:HD13	1:A:78:SER:N	2.24	0.53
1:A:164:ASN:HB2	1:A:171[B]:VAL:HG23	1.90	0.52
4:A:407:EDO:H22	7:A:549:HOH:O	2.10	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:595:HOH:O	1:B:296:MET:HE2	2.12	0.50
4:A:408:EDO:H21	7:B:1758:HOH:O	2.11	0.50
1:A:307:ASN:HA	2:A:401:NAD:H2A	1.96	0.48
4:A:408:EDO:O2	3:B:1602:SPD:N10	2.47	0.48
1:B:92:ARG:NH1	1:B:123:ASN:OD1	2.30	0.48
1:B:327:TRP:CZ3	3:B:1602:SPD:H72	2.49	0.47
1:B:327:TRP:CZ2	3:B:1602:SPD:H31	2.50	0.47
1:A:43:ARG:HH11	1:A:357:GLN:NE2	2.14	0.45
1:B:329:LYS:NZ	3:B:1602:SPD:C5	2.79	0.45
1:B:283:GLY:O	1:B:286[B]:VAL:HG22	2.16	0.45
1:B:137[B]:GLU:OE2	7:B:1702:HOH:O	2.21	0.44
1:A:83:GLN:OE1	1:A:83:GLN:HA	2.16	0.44
1:A:268:THR:HG21	5:A:406:BME:H12	2.00	0.44
1:A:137:GLU:OE2	2:A:401:NAD:H4N	2.17	0.44
1:B:92:ARG:HD3	7:B:1821:HOH:O	2.18	0.43
1:A:43:ARG:NH1	1:A:357:GLN:NE2	2.66	0.42
1:A:123:ASN:ND2	1:A:227:ASN:HB3	2.33	0.42
1:A:283:GLY:O	1:A:286[B]:VAL:HG22	2.19	0.42
1:B:268[B]:THR:HG23	1:B:272:PHE:CD2	2.54	0.41
1:A:32:ARG:NH1	7:A:510:HOH:O	2.53	0.41
1:A:123:ASN:HD21	1:A:227:ASN:HB3	1.85	0.41
1:A:327:TRP:CE2	1:B:164:ASN:HB3	2.55	0.41
1:B:329:LYS:HZ1	3:B:1602:SPD:C5	2.23	0.41
1:A:265:LEU:HD23	5:A:406:BME:H22	2.03	0.40
1:A:323:GLU:OE1	3:A:402:SPD:H91	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	350/369 (95%)	345 (99%)	5 (1%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	343/369 (93%)	337 (98%)	6 (2%)	0	100	100
All	All	693/738 (94%)	682 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	293/309 (95%)	293 (100%)	0	100	100
1	B	293/309 (95%)	293 (100%)	0	100	100
All	All	586/618 (95%)	586 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	89	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CSS	A	177	1	4,6,7	0.64	0	2,6,8	0.34	0
1	CSS	B	177	1	4,6,7	1.15	0	2,6,8	0.86	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSS	A	177	1	-	0/1/5/7	-
1	CSS	B	177	1	-	0/1/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EDO	A	405	-	3,3,3	0.36	0	2,2,2	0.22	0
4	EDO	A	404	-	3,3,3	0.35	0	2,2,2	0.52	0
5	BME	A	406	-	3,3,3	0.35	0	2,2,2	0.96	0
3	SPD	A	402	-	9,9,9	0.81	0	8,8,8	0.98	0
3	SPD	B	1602	-	9,9,9	0.19	0	8,8,8	0.34	0
4	EDO	A	407	-	3,3,3	0.31	0	2,2,2	0.40	0
4	EDO	B	1604	-	3,3,3	0.42	0	2,2,2	0.53	0
2	NAD	A	401	-	46,48,48	0.63	0	64,73,73	1.11	5 (7%)
6	OXM	B	1603	-	5,5,5	2.72	1 (20%)	2,6,6	0.95	0
2	NAD	B	1601	-	46,48,48	2.07	17 (36%)	64,73,73	1.94	18 (28%)
4	EDO	A	403	-	3,3,3	0.41	0	2,2,2	0.34	0
4	EDO	A	408	-	3,3,3	0.30	0	2,2,2	0.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	405	-	-	1/1/1/1	-
4	EDO	A	404	-	-	0/1/1/1	-
5	BME	A	406	-	-	0/1/1/1	-
3	SPD	A	402	-	-	1/7/7/7	-
3	SPD	B	1602	-	-	4/7/7/7	-
4	EDO	A	407	-	-	1/1/1/1	-
4	EDO	B	1604	-	-	0/1/1/1	-
2	NAD	A	401	-	-	5/30/62/62	0/5/5/5
6	OXM	B	1603	-	-	0/4/4/4	-
2	NAD	B	1601	-	-	6/30/62/62	0/5/5/5
4	EDO	A	403	-	-	0/1/1/1	-
4	EDO	A	408	-	-	0/1/1/1	-

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	1603	OXM	C1-C2	-5.75	1.48	1.55
2	B	1601	NAD	C5A-N7A	-4.47	1.30	1.39
2	B	1601	NAD	PN-O3	-4.12	1.55	1.59
2	B	1601	NAD	C8A-N9A	-3.88	1.30	1.37
2	B	1601	NAD	PN-O2N	-3.57	1.38	1.55
2	B	1601	NAD	O7N-C7N	-3.42	1.17	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1601	NAD	C4A-N9A	-3.39	1.30	1.37
2	B	1601	NAD	PA-O2A	-3.32	1.40	1.55
2	B	1601	NAD	C5A-C4A	3.23	1.44	1.39
2	B	1601	NAD	C2N-N1N	-2.94	1.31	1.35
2	B	1601	NAD	PN-O1N	-2.60	1.41	1.50
2	B	1601	NAD	PA-O3	-2.59	1.56	1.59
2	B	1601	NAD	PA-O1A	-2.41	1.42	1.50
2	B	1601	NAD	O4B-C4B	-2.13	1.40	1.45
2	B	1601	NAD	O5D-C5D	-2.08	1.36	1.44
2	B	1601	NAD	O4D-C4D	-2.07	1.40	1.45
2	B	1601	NAD	C4A-N3A	-2.02	1.30	1.34
2	B	1601	NAD	C2N-C3N	-2.01	1.35	1.39

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1601	NAD	C5A-C4A-N3A	-6.56	117.68	126.72
2	B	1601	NAD	C4A-C5A-N7A	-4.57	105.36	110.58
2	B	1601	NAD	N3A-C4A-N9A	4.52	134.85	127.17
2	B	1601	NAD	C2A-N3A-C4A	4.15	121.97	111.83
2	B	1601	NAD	O3D-C3D-C4D	3.87	122.20	111.08
2	B	1601	NAD	N3A-C2A-N1A	-3.34	123.53	128.58
2	B	1601	NAD	O4B-C1B-N9A	3.32	114.46	108.09
2	B	1601	NAD	O2A-PA-O3	3.18	115.86	107.27
2	A	401	NAD	O4B-C1B-N9A	3.12	114.08	108.09
2	A	401	NAD	C4D-O4D-C1D	-3.08	107.11	109.92
2	B	1601	NAD	C2D-C3D-C4D	3.00	108.40	102.61
2	B	1601	NAD	C5A-N7A-C8A	2.78	107.83	103.45
2	B	1601	NAD	O3D-C3D-C2D	2.71	120.49	111.82
2	B	1601	NAD	O5D-PN-O1N	2.71	119.66	108.94
2	A	401	NAD	O2B-C2B-C1B	2.63	119.15	110.10
2	B	1601	NAD	C6A-C5A-N7A	2.58	137.07	132.09
2	B	1601	NAD	O4D-C4D-C5D	2.53	117.43	109.33
2	B	1601	NAD	C5N-C4N-C3N	-2.26	118.14	120.36
2	B	1601	NAD	O5D-C5D-C4D	2.25	116.64	108.99
2	A	401	NAD	O2A-PA-O1A	2.23	122.83	112.44
2	A	401	NAD	O3-PA-O1A	-2.21	104.04	110.70
2	B	1601	NAD	C2A-N1A-C6A	2.19	122.33	118.73
2	B	1601	NAD	C2B-C1B-N9A	-2.06	108.20	113.30

There are no chirality outliers.

All (18) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	401	NAD	PA-O3-PN-O5D
2	A	401	NAD	C5D-O5D-PN-O3
2	A	401	NAD	C5D-O5D-PN-O1N
2	A	401	NAD	C5D-O5D-PN-O2N
2	B	1601	NAD	PA-O3-PN-O5D
2	B	1601	NAD	C5D-O5D-PN-O3
2	B	1601	NAD	C5D-O5D-PN-O1N
2	B	1601	NAD	C5D-O5D-PN-O2N
3	B	1602	SPD	C3-C4-C5-N6
3	B	1602	SPD	N6-C7-C8-C9
3	A	402	SPD	N6-C7-C8-C9
3	B	1602	SPD	N1-C2-C3-C4
3	B	1602	SPD	C7-C8-C9-N10
4	A	405	EDO	O1-C1-C2-O2
4	A	407	EDO	O1-C1-C2-O2
2	A	401	NAD	PN-O3-PA-O1A
2	B	1601	NAD	PN-O3-PA-O1A
2	B	1601	NAD	PN-O3-PA-O2A


There are no ring outliers.

7 monomers are involved in 21 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	405	EDO	2	0
5	A	406	BME	3	0
3	A	402	SPD	4	0
3	B	1602	SPD	8	0
4	A	407	EDO	1	0
2	A	401	NAD	2	0
4	A	408	EDO	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	284:GLY	C	285:VAL	N	1.14

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	335/369 (90%)	-0.15	14 (4%)	40 44	17, 36, 76, 135	13 (3%)
1	B	327/369 (88%)	-0.23	10 (3%)	51 55	15, 35, 66, 137	17 (5%)
All	All	662/738 (89%)	-0.19	24 (3%)	46 50	15, 36, 72, 137	30 (4%)

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	93	PRO	6.2
1	A	84	HIS	5.6
1	B	364	HIS	5.4
1	A	78	SER	5.3
1	A	87	LEU	4.8
1	A	81	GLU	4.0
1	B	94	LEU	3.7
1	A	77	LEU	3.4
1	A	363	MET	3.4
1	A	82	ASP	2.9
1	A	85	ALA	2.7
1	A	88	THR	2.7
1	B	28	SER	2.7
1	B	91	ARG	2.6
1	B	80	ASP	2.6
1	B	81	GLU	2.4
1	A	80	ASP	2.4
1	A	94	LEU	2.4
1	A	28	SER	2.3
1	A	86[A]	ASP	2.3
1	A	359	MET	2.3
1	B	79	GLN	2.1
1	B	78	SER	2.1
1	B	77	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CSS	B	177	7/8	0.96	0.07	40,50,68,82	0
1	CSS	A	177	7/8	0.97	0.06	28,36,51,61	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	EDO	A	407	4/4	0.77	0.37	114,136,137,137	0
5	BME	A	406	4/4	0.77	0.45	180,217,219,219	0
4	EDO	A	405	4/4	0.82	0.27	63,76,87,87	0
4	EDO	A	404	4/4	0.89	0.25	74,89,100,102	0
6	OXM	B	1603	6/6	0.89	0.13	47,53,67,67	0
4	EDO	A	408	4/4	0.91	0.14	32,44,51,53	10
3	SPD	A	402	10/10	0.93	0.11	36,47,55,57	0
3	SPD	B	1602	10/10	0.95	0.09	29,39,52,53	0
4	EDO	B	1604	4/4	0.97	0.06	22,27,29,33	10
2	NAD	B	1601	44/44	0.97	0.06	22,32,42,45	0
4	EDO	A	403	4/4	0.97	0.07	32,38,40,44	0
2	NAD	A	401	44/44	0.98	0.05	21,30,38,44	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around NAD B 1601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAD A 401:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.