



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 05:37 AM UTC

PDB ID : 1A7N / pdb_00001a7n
Title : FV FRAGMENT OF MOUSE MONOCLONAL ANTIBODY D1.3 (BALB/C, IGG1, K) VARIANT FOR CHAIN L GLU81->ASP AND CHAIN H LEU312->VAL
Authors : Marks, C.; Henrick, K.; Winter, G.
Deposited on : 1998-03-16
Resolution : 2.01 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.01 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	107	
2	H	116	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1909 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IGG1-KAPPA D1.3 FV (LIGHT CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	107	Total	C	N	O	S	0	0	0
			816	518	135	161	2			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
L	3	VAL	GLU	conflict	UNP P01635
L	50	TYR	LYS	conflict	UNP P01635
L	51	THR	ALA	conflict	UNP P01635
L	52	THR	GLN	conflict	UNP P01635
L	81	ASP	GLU	variant	UNP P01635
L	?	-	PRO	deletion	UNP P01635
L	96	ARG	TRP	conflict	UNP P01635

- Molecule 2 is a protein called IGG1-KAPPA D1.3 FV (HEAVY CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	116	Total	C	N	O	S	0	0	0
			901	563	157	177	4			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	L	90	Total	O	0	0
			90	90		
3	H	102	Total	O	0	0
			102	102		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: IGG1-KAPPA D1.3 FV (LIGHT CHAIN)

Chain L: 



- Molecule 2: IGG1-KAPPA D1.3 FV (HEAVY CHAIN)

Chain H: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	90.18Å 90.18Å 56.63Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.00 – 2.01 6.00 – 2.02	Depositor EDS
% Data completeness (in resolution range)	(Not available) (6.00-2.01) 95.2 (6.00-2.02)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.64 (at 2.02Å)	Xtriage
Refinement program	CCP4	Depositor
R, R_{free}	0.155 , (Not available) 0.159 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	21.0	Xtriage
Anisotropy	0.094	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 84.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	1909	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	L	1.14	2/837 (0.2%)	2.19	34/1138 (3.0%)
2	H	1.12	0/921	2.42	55/1250 (4.4%)
All	All	1.13	2/1758 (0.1%)	2.31	89/2388 (3.7%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	L	52	THR	CA-CB	5.23	1.61	1.54
1	L	27	GLY	N-CA	5.15	1.50	1.45

All (89) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	308	GLN	CB-CG-CD	26.64	157.90	112.60
1	L	45	GLN	CB-CG-CD	14.43	137.14	112.60
1	L	81	ASP	CA-CB-CG	10.63	123.23	112.60
1	L	70	GLN	OE1-CD-NE2	10.01	132.61	122.60
2	H	201	GLN	CA-C-O	9.58	137.09	120.80
2	H	308	GLN	OE1-CD-NE2	-9.55	113.05	122.60
1	L	62	PHE	CA-C-O	-8.77	110.88	120.43
2	H	215	SER	O-C-N	8.41	129.55	121.56
1	L	72	SER	N-CA-CB	-8.18	96.32	111.53
2	H	308	GLN	CA-CB-CG	-8.01	98.08	114.10
1	L	37	GLN	OE1-CD-NE2	-7.98	114.62	122.60
2	H	283	ASN	OD1-CG-ND2	7.79	130.40	122.60
2	H	266	ARG	NE-CZ-NH2	-7.76	112.22	119.20
1	L	67	SER	O-C-N	7.75	131.13	123.46
2	H	300	ASP	CA-CB-CG	-7.72	104.88	112.60
2	H	301	TYR	N-CA-C	7.53	121.79	111.54
2	H	209	PRO	CA-C-O	-7.53	109.42	118.90
2	H	240	PRO	N-CA-CB	7.50	107.39	103.19
2	H	282	MET	CA-CB-CG	7.48	129.07	114.10
2	H	286	HIS	CA-CB-CG	-7.43	106.36	113.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	256	ASN	CA-CB-CG	-7.34	105.26	112.60
1	L	2	ILE	CA-C-N	-7.09	112.24	122.45
1	L	2	ILE	C-N-CA	-7.09	112.24	122.45
1	L	70	GLN	N-CA-CB	-7.01	99.77	110.77
2	H	291	ALA	O-C-N	6.97	129.81	123.26
2	H	267	LEU	N-CA-C	6.91	120.94	109.95
1	L	21	ILE	CB-CA-C	6.83	121.34	110.82
2	H	201	GLN	CA-CB-CG	6.75	127.59	114.10
1	L	105	GLU	CA-C-O	6.62	128.59	121.11
2	H	300	ASP	O-C-N	-6.61	113.80	122.59
2	H	204	LEU	O-C-N	6.51	130.93	123.31
2	H	202	VAL	N-CA-C	-6.49	99.13	108.53
1	L	7	SER	CB-CA-C	6.42	118.77	109.45
2	H	240	PRO	O-C-N	6.40	124.25	121.31
2	H	255	GLY	CA-C-O	-6.34	112.19	119.10
1	L	4	LEU	O-C-N	6.30	130.73	123.29
2	H	237	VAL	CA-C-N	-6.26	112.76	122.59
2	H	237	VAL	C-N-CA	-6.26	112.76	122.59
1	L	99	GLY	N-CA-C	-6.23	102.82	112.85
1	L	94	THR	N-CA-CB	6.15	119.92	111.25
2	H	206	GLU	CA-C-O	-6.15	114.03	121.28
2	H	312	VAL	CA-C-O	-6.14	113.94	120.39
1	L	22	THR	CA-CB-OG1	-6.12	100.42	109.60
2	H	283	ASN	O-C-N	6.12	129.95	123.03
2	H	307	GLY	N-CA-C	-6.02	102.92	112.66
1	L	70	GLN	CG-CD-NE2	-5.92	107.52	116.40
2	H	218	LEU	N-CA-CB	-5.89	100.35	111.00
2	H	295	CYS	N-CA-C	-5.88	100.61	109.95
2	H	239	GLN	CA-C-N	5.86	123.94	119.66
2	H	239	GLN	C-N-CA	5.86	123.94	119.66
2	H	203	GLN	CB-CA-C	5.80	121.16	109.38
2	H	302	ARG	O-C-N	5.75	129.66	123.56
2	H	291	ALA	CA-C-N	-5.71	114.84	122.72
2	H	291	ALA	C-N-CA	-5.71	114.84	122.72
2	H	284	SER	O-C-N	5.67	130.13	122.59
2	H	265	SER	N-CA-CB	-5.63	101.54	110.22
2	H	204	LEU	CA-C-O	-5.63	114.07	120.32
2	H	274	SER	N-CA-C	5.62	117.08	111.07
1	L	67	SER	N-CA-CB	5.60	120.08	111.56
1	L	27	GLY	CA-C-O	-5.60	115.27	121.37
2	H	208	GLY	O-C-N	5.59	127.36	121.77
1	L	10	SER	N-CA-CB	5.57	121.89	111.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	76	ASN	CB-CG-ND2	5.53	124.69	116.40
2	H	306	TRP	CA-C-O	-5.47	114.93	121.11
1	L	86	TYR	O-C-N	5.46	129.80	123.30
1	L	106	ILE	CB-CA-C	5.44	119.44	111.33
2	H	302	ARG	CB-CG-CD	-5.40	98.88	111.30
2	H	283	ASN	CB-CA-C	-5.39	100.91	109.80
2	H	262	ALA	CA-C-O	-5.37	114.86	120.55
2	H	240	PRO	N-CA-C	-5.36	105.33	110.47
1	L	29	ILE	CB-CA-C	5.34	117.82	111.20
2	H	248	LEU	O-C-N	5.31	129.95	122.20
2	H	312	VAL	O-C-N	5.25	128.93	123.26
1	L	73	LEU	CA-C-O	5.21	125.88	120.30
1	L	14	SER	O-C-N	5.20	128.90	123.03
1	L	101	GLY	CA-C-O	5.19	126.75	122.24
1	L	22	THR	O-C-N	5.14	129.63	123.16
2	H	304	ASP	CA-C-O	-5.13	113.73	119.79
2	H	223	THR	OG1-CB-CG2	5.13	119.56	109.30
2	H	223	THR	CA-CB-OG1	-5.11	101.94	109.60
1	L	31	ASN	CA-CB-CG	-5.10	107.50	112.60
1	L	71	TYR	CA-C-N	-5.07	113.92	122.29
1	L	71	TYR	C-N-CA	-5.07	113.92	122.29
2	H	313	THR	CA-CB-OG1	-5.07	102.00	109.60
2	H	299	ARG	NE-CZ-NH2	5.05	123.75	119.20
2	H	300	ASP	CB-CA-C	5.05	120.47	110.42
1	L	94	THR	CA-CB-OG1	-5.04	102.05	109.60
2	H	289	ASP	N-CA-C	-5.02	106.94	113.16
1	L	76	ASN	OD1-CG-ND2	-5.01	117.59	122.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	816	0	783	7	0
2	H	901	0	864	14	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	H	102	0	0	2	0
3	L	90	0	0	4	0
All	All	1909	0	1647	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:3:VAL:HG12	3:L:448:HOH:O	1.37	1.23
1:L:30:HIS:HE1	3:L:427:HOH:O	1.60	0.84
1:L:30:HIS:HD2	3:L:472:HOH:O	1.70	0.74
2:H:300:ASP:O	2:H:300:ASP:OD1	2.19	0.60
2:H:218:LEU:HA	3:H:420:HOH:O	2.01	0.60
1:L:37:GLN:O	1:L:45:GLN:HG2	2.06	0.55
2:H:276:SER:OG	3:H:507:HOH:O	2.19	0.53
1:L:94:THR:HB	1:L:95:PRO:HA	1.91	0.53
2:H:218:LEU:HB2	2:H:285:LEU:HD11	1.92	0.51
2:H:220:ILE:HD12	2:H:280:LEU:HD23	1.94	0.48
1:L:37:GLN:OE1	1:L:45:GLN:NE2	2.46	0.48
2:H:214:PRO:O	2:H:215:SER:HB2	2.15	0.46
2:H:267:LEU:HD22	2:H:280:LEU:HD11	1.97	0.46
2:H:254:ASP:CG	2:H:256:ASN:HD22	2.25	0.45
2:H:203:GLN:HB3	2:H:225:SER:HB2	2.01	0.43
2:H:275:LYS:O	2:H:276:SER:HB2	2.19	0.43
2:H:297:ARG:O	2:H:303:LEU:HA	2.19	0.42
1:L:30:HIS:CE1	3:L:427:HOH:O	2.49	0.42
2:H:240:PRO:HA	2:H:241:PRO:HD3	1.94	0.41
2:H:300:ASP:OD1	2:H:300:ASP:C	2.62	0.41
2:H:302:ARG:HH11	2:H:302:ARG:HD3	1.67	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	105/107 (98%)	102 (97%)	2 (2%)	1 (1%)	12	8
2	H	114/116 (98%)	110 (96%)	4 (4%)	0	100	100
All	All	219/223 (98%)	212 (97%)	6 (3%)	1 (0%)	24	21

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	77	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	89/91 (98%)	86 (97%)	3 (3%)	32	33
2	H	99/100 (99%)	93 (94%)	6 (6%)	17	14
All	All	188/191 (98%)	179 (95%)	9 (5%)	23	21

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	39	LYS
1	L	81	ASP
1	L	104	LEU
2	H	201	GLN
2	H	224	VAL
2	H	228	SER
2	H	260	ASN
2	H	275	LYS
2	H	299	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	30	HIS
1	L	45	GLN
1	L	76	ASN
1	L	89	GLN
2	H	216	GLN
2	H	256	ASN
2	H	286	HIS
2	H	308	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	L	107/107 (100%)	-1.02	0 100 100	11, 18, 31, 44	6 (5%)
2	H	116/116 (100%)	-1.04	1 (0%) 81 80	9, 18, 32, 36	4 (3%)
All	All	223/223 (100%)	-1.03	1 (0%) 88 88	9, 18, 32, 44	10 (4%)

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	300	ASP	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.