



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2026 – 10:28 PM UTC

PDB ID : 2A7P / pdb_00002a7p
Title : Crystal Structure of the G81A mutant of the Active Chimera of (S)-Mandelate Dehydrogenase in complex with its substrate 3-indolelactate
Authors : Sukumar, N.; Xu, Y.; Mitra, B.; Mathews, F.S.
Deposited on : 2005-07-05
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	380	

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called (S)-Mandelate Dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	353	Total 2748	C 1740	N 488	O 507	S 13	10	0	0

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	81	ALA	GLY	engineered mutation	UNP P20932
A	375	HIS	-	expression tag	UNP P20932
A	376	HIS	-	expression tag	UNP P20932
A	377	HIS	-	expression tag	UNP P20932
A	378	HIS	-	expression tag	UNP P20932
A	379	HIS	-	expression tag	UNP P20932
A	380	HIS	-	expression tag	UNP P20932

- Molecule 2 is FLAVIN MONONUCLEOTIDE (CCD ID: FMN) (formula: $\text{C}_{17}\text{H}_{21}\text{N}_4\text{O}_9\text{P}$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			31	17	4	9	1		

- Molecule 3 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: $C_6H_{13}NO_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 4 is 3-(INDOL-3-YL) LACTATE (CCD ID: 3IL) (formula: $C_{11}H_{11}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			15	11	1	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	224	Total	O	0	0
			224	224		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: (S)-Mandelate Dehydrogenase



4 Data and refinement statistics

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, α , β , γ	99.40 Å 99.40 Å 87.60 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 2.20 40.00 – 2.20	Depositor EDS
% Data completeness (in resolution range)	(Not available) (40.00-2.20) 88.5 (40.00-2.20)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.06 (at 2.05 Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.198 , 0.243 0.198 , 0.243	Depositor DCC
R_{free} test set	1360 reflections (5.84%)	wwPDB-VP
Wilson B-factor (Å ²)	17.8	Xtriage
Anisotropy	0.321	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 47.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.039 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3030	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.90% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 3IL, MES, FMN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.63	5/2797 (0.2%)	1.01	8/3784 (0.2%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	181	LEU	C-N	-11.55	1.17	1.33
1	A	195	ALA	C-N	-9.52	1.20	1.33
1	A	203	LEU	CA-C	6.52	1.61	1.52
1	A	195	ALA	CA-C	-5.66	1.45	1.52
1	A	197	LEU	C-N	5.22	1.41	1.33

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	196	ASN	O-C-N	-14.97	102.68	122.59
1	A	133	ILE	N-CA-C	-7.29	104.67	111.45
1	A	198	GLU	N-CA-C	-6.54	104.78	112.89
1	A	176	PRO	CB-CA-C	6.53	118.89	110.92
1	A	35	TYR	N-CA-C	-6.20	104.42	112.23
1	A	29	GLY	N-CA-C	5.87	118.78	112.33
1	A	265	SER	N-CA-C	-5.85	102.67	109.93
1	A	20	PRO	N-CA-C	-5.62	102.37	111.19

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2748	0	2782	85	0
2	A	31	0	19	2	0
3	A	12	0	13	0	0
4	A	15	0	10	0	0
5	A	224	0	0	6	0
All	All	3030	0	2824	85	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (85) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:191:LYS:HB2	1:A:196:ASN:HB3	1.35	1.07
1:A:195:ALA:O	1:A:196:ASN:HB2	1.73	0.86
1:A:84:GLY:H	1:A:112:ASN:HD22	1.24	0.86
1:A:40:ASN:HD21	1:A:254:ASN:HD21	1.23	0.84
1:A:180:THR:HG23	1:A:182:LYS:HG2	1.60	0.83
1:A:355:ASN:HB3	5:A:1067:HOH:O	1.78	0.83
1:A:189:LEU:HD12	1:A:200:GLN:HG2	1.63	0.81
1:A:191:LYS:HB2	1:A:196:ASN:CB	2.12	0.80
1:A:137:ILE:CD1	1:A:203:LEU:HD21	2.14	0.77
1:A:355:ASN:CB	5:A:1067:HOH:O	2.35	0.74
1:A:137:ILE:HD11	1:A:203:LEU:HD21	1.68	0.74
1:A:112:ASN:O	1:A:182:LYS:HG3	1.90	0.71
1:A:200:GLN:HA	1:A:203:LEU:HD12	1.73	0.70
1:A:133:ILE:HG22	1:A:202:ALA:HB1	1.74	0.70
1:A:199:MET:O	1:A:203:LEU:HG	1.95	0.65
1:A:40:ASN:ND2	1:A:254:ASN:HD21	1.94	0.64
1:A:137:ILE:CD1	1:A:203:LEU:CD2	2.74	0.64
1:A:116:GLU:H	1:A:116:GLU:CD	2.06	0.63
1:A:284:ASP:O	1:A:285:SER:HB2	1.97	0.62
1:A:92:LEU:HB2	1:A:113:MET:HE1	1.81	0.62
1:A:84:GLY:N	1:A:112:ASN:HD22	1.95	0.62
1:A:180:THR:CG2	1:A:182:LYS:HG2	2.30	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:134:HIS:CD2	1:A:136:GLU:HB3	2.38	0.59
1:A:176:PRO:HG2	1:A:179:LEU:HD12	1.85	0.58
1:A:191:LYS:HG3	1:A:196:ASN:CG	2.28	0.58
1:A:189:LEU:CD1	1:A:200:GLN:HG2	2.30	0.58
1:A:84:GLY:H	1:A:112:ASN:ND2	1.97	0.58
1:A:194:LYS:C	1:A:195:ALA:O	2.44	0.58
1:A:205:SER:C	1:A:207:GLN:H	2.12	0.58
1:A:75:LEU:HD21	1:A:293:ILE:HD13	1.87	0.56
1:A:137:ILE:HD13	1:A:203:LEU:CD2	2.34	0.56
1:A:45:GLN:NE2	5:A:1111:HOH:O	2.39	0.55
1:A:253:SER:HB2	1:A:284:ASP:OD1	2.05	0.55
1:A:284:ASP:OD1	1:A:284:ASP:C	2.50	0.55
1:A:133:ILE:HD12	1:A:133:ILE:N	2.22	0.54
1:A:116:GLU:HG2	1:A:144:LYS:HE3	1.90	0.53
1:A:5:LEU:HB3	1:A:11:TYR:CE1	2.44	0.53
1:A:40:ASN:HD21	1:A:254:ASN:ND2	2.01	0.53
1:A:129:GLN:NE2	2:A:390:FMN:HN3	2.07	0.53
1:A:191:LYS:CB	1:A:196:ASN:HB3	2.24	0.52
1:A:199:MET:HE3	1:A:203:LEU:HD11	1.91	0.52
1:A:189:LEU:HD12	1:A:200:GLN:CG	2.38	0.51
1:A:108:SER:HA	1:A:129:GLN:HE21	1.76	0.51
1:A:137:ILE:HD13	1:A:203:LEU:HD21	1.89	0.51
1:A:191:LYS:CB	1:A:196:ASN:CB	2.87	0.51
1:A:139:GLN:HG3	1:A:223:LEU:CD2	2.41	0.50
1:A:32:GLU:HG3	5:A:933:HOH:O	2.10	0.50
1:A:198:GLU:O	1:A:201:ALA:HB3	2.12	0.49
1:A:109:THR:OG1	1:A:130:LEU:HA	2.12	0.49
1:A:284:ASP:O	1:A:285:SER:CB	2.61	0.49
1:A:74:PRO:HG2	1:A:345:ILE:HD13	1.93	0.48
1:A:317:ARG:HG2	1:A:317:ARG:HH11	1.79	0.48
1:A:116:GLU:HG2	1:A:144:LYS:CE	2.44	0.48
1:A:194:LYS:O	1:A:197:LEU:HB2	2.13	0.48
1:A:200:GLN:HB3	1:A:204:MET:HE3	1.95	0.48
1:A:194:LYS:O	1:A:195:ALA:O	2.32	0.47
1:A:191:LYS:CB	1:A:196:ASN:CG	2.88	0.47
1:A:195:ALA:O	1:A:196:ASN:CB	2.39	0.47
1:A:139:GLN:HG3	1:A:223:LEU:HD21	1.96	0.47
1:A:187:ILE:HG21	1:A:203:LEU:HD21	1.98	0.46
1:A:229:LEU:HD22	1:A:249:GLY:C	2.41	0.46
1:A:115:ILE:HG23	1:A:150:TYR:CE1	2.51	0.46
1:A:195:ALA:C	1:A:197:LEU:N	2.74	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:75:LEU:C	1:A:76:LEU:HD12	2.41	0.45
1:A:85:ALA:HA	1:A:179:LEU:HD11	1.99	0.45
1:A:308:ARG:HG3	2:A:390:FMN:O2P	2.16	0.45
1:A:195:ALA:C	1:A:197:LEU:H	2.24	0.45
1:A:253:SER:CB	1:A:284:ASP:OD1	2.65	0.45
1:A:134:HIS:HD2	1:A:137:ILE:HG13	1.81	0.45
1:A:355:ASN:CG	5:A:1067:HOH:O	2.58	0.45
1:A:62:LEU:HD12	1:A:62:LEU:N	2.32	0.44
1:A:116:GLU:CG	1:A:144:LYS:HE3	2.48	0.43
1:A:115:ILE:HD11	1:A:128:PHE:CZ	2.54	0.43
1:A:77:ILE:HD12	1:A:98:ALA:HB2	2.01	0.42
1:A:153:LEU:HD21	1:A:220:LEU:HD11	2.01	0.42
1:A:200:GLN:HB3	1:A:204:MET:CE	2.50	0.42
1:A:159:VAL:HG12	1:A:159:VAL:O	2.20	0.42
1:A:20:PRO:O	1:A:21:LYS:C	2.63	0.41
1:A:284:ASP:HB3	1:A:305:LEU:HB2	2.03	0.41
1:A:59:ARG:HG3	1:A:59:ARG:HH11	1.84	0.41
1:A:233:LEU:HD13	1:A:239:ALA:HA	2.03	0.41
1:A:29:GLY:HA3	5:A:977:HOH:O	2.20	0.41
1:A:94:LEU:HD23	1:A:322:VAL:HG21	2.03	0.41
1:A:182:LYS:HE2	1:A:182:LYS:HB3	1.33	0.41
1:A:131:TYR:HB3	1:A:208:MET:SD	2.60	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	232/380 (61%)	215 (93%)	14 (6%)	3 (1%)	9 8

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	194	LYS
1	A	195	ALA
1	A	285	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	288/311 (93%)	282 (98%)	6 (2%)	47 63

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	69	LYS
1	A	182	LYS
1	A	197	LEU
1	A	302	GLU
1	A	310	THR
1	A	339	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	16	GLN
1	A	40	ASN
1	A	46	GLN
1	A	112	ASN
1	A	129	GLN
1	A	134	HIS
1	A	162	ASN
1	A	200	GLN
1	A	207	GLN
1	A	259	GLN
1	A	272	GLN
1	A	339	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FMN	A	390	-	33,33,33	2.99	15 (45%)	48,50,50	1.61	11 (22%)
3	MES	A	890	-	12,12,12	2.94	4 (33%)	15,16,16	1.03	0
4	3IL	A	410	-	16,16,16	2.34	10 (62%)	21,22,22	2.00	5 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FMN	A	390	-	-	1/18/18/18	0/3/3/3
3	MES	A	890	-	-	0/6/14/14	0/1/1/1
4	3IL	A	410	-	-	0/8/8/8	0/2/2/2

All (29) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	890	MES	O1S-S	7.89	1.67	1.45
2	A	390	FMN	C9A-C5A	7.61	1.53	1.41
2	A	390	FMN	C8-C7	5.77	1.54	1.40
2	A	390	FMN	C9-C9A	5.76	1.48	1.39
2	A	390	FMN	C6-C7	5.59	1.47	1.39
2	A	390	FMN	C9A-N10	4.67	1.49	1.41
3	A	890	MES	O2S-S	4.25	1.57	1.45
2	A	390	FMN	C6-C5A	4.01	1.46	1.40
2	A	390	FMN	C2-N3	3.89	1.47	1.39
4	A	410	3IL	C4-C9	3.56	1.45	1.39
2	A	390	FMN	O2-C2	-3.56	1.17	1.24
2	A	390	FMN	C4A-C10	3.45	1.54	1.44
4	A	410	3IL	C2-C3	3.42	1.41	1.36
2	A	390	FMN	C10-N10	3.27	1.44	1.37
2	A	390	FMN	C9-C8	3.04	1.43	1.39
2	A	390	FMN	O2'-C2'	-2.98	1.37	1.43
4	A	410	3IL	C7-C8	2.96	1.44	1.39
4	A	410	3IL	C2-N1	2.84	1.41	1.37
4	A	410	3IL	C9-C8	2.67	1.44	1.41
3	A	890	MES	C5-N4	2.59	1.53	1.46
3	A	890	MES	O3S-S	2.50	1.56	1.47
2	A	390	FMN	C4-N3	2.48	1.43	1.38
4	A	410	3IL	C9-C3	-2.47	1.39	1.44
4	A	410	3IL	C5-C4	2.36	1.42	1.38
4	A	410	3IL	C6-C7	2.36	1.42	1.38
4	A	410	3IL	C6-C5	2.32	1.43	1.38
4	A	410	3IL	C15-C3	2.30	1.54	1.50
2	A	390	FMN	O3'-C3'	2.14	1.48	1.43
2	A	390	FMN	C4A-N5	2.11	1.35	1.30

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	410	3IL	C8-N1-C2	-5.84	103.88	109.08
2	A	390	FMN	C9-C9A-N10	4.43	127.81	121.85
4	A	410	3IL	C9-C8-N1	3.87	110.75	107.52
2	A	390	FMN	C5A-C9A-N10	-3.35	114.94	117.97
4	A	410	3IL	O11-C10-C13	2.90	118.88	112.74
2	A	390	FMN	O3'-C3'-C4'	-2.63	102.96	108.93
2	A	390	FMN	C9A-C9-C8	2.62	124.50	119.22
2	A	390	FMN	C4-C4A-N5	2.59	121.79	118.21
2	A	390	FMN	C5A-N5-C4A	2.56	122.23	118.09
2	A	390	FMN	C9A-N10-C10	2.32	124.29	120.75

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	390	FMN	C4'-C3'-C2'	2.20	117.23	113.57
2	A	390	FMN	O4-C4-C4A	-2.15	120.86	126.53
2	A	390	FMN	O4'-C4'-C3'	2.11	114.19	109.25
2	A	390	FMN	C10-N1-C2	2.10	121.39	116.85
4	A	410	3IL	C15-C3-C9	2.09	130.72	126.70
4	A	410	3IL	C3-C2-N1	2.09	112.60	110.31

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	390	FMN	C4'-C5'-O5'-P

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	390	FMN	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	195:ALA	C	196:ASN	N	1.20
1	A	181:LEU	C	182:LYS	N	1.17

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	353/380 (92%)	0.22	28 (7%) 18 16	10, 24, 82, 91	2 (0%)

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	195	ALA	4.9
1	A	189	LEU	4.8
1	A	187	ILE	4.7
1	A	203	LEU	4.5
1	A	197	LEU	4.1
1	A	201	ALA	4.1
1	A	178	PHE	3.9
1	A	192	MET	3.7
1	A	194	LYS	3.6
1	A	190	GLY	3.6
1	A	196	ASN	3.5
1	A	188	ASP	3.4
1	A	199	MET	3.4
1	A	204	MET	3.1
1	A	180	THR	2.9
1	A	202	ALA	2.8
1	A	207	GLN	2.7
1	A	175	ILE	2.7
1	A	191	LYS	2.6
1	A	186	GLY	2.5
1	A	193	ASP	2.5
1	A	177	PRO	2.5
1	A	208	MET	2.4
1	A	198	GLU	2.4
1	A	200	GLN	2.3
1	A	205	SER	2.3
1	A	206	ARG	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	176	PRO	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	3IL	A	410	15/15	0.89	0.11	35,40,42,42	0
3	MES	A	890	12/12	0.95	0.09	21,23,25,26	0
2	FMN	A	390	31/31	0.96	0.06	10,15,18,20	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FMN A 390:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.