



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 04:26 AM UTC

PDB ID : 3A7Q / pdb_00003a7q
Title : Structural basis for specific recognition of reelin by its receptors
Authors : Yasui, N.; Nogi, T.; Takagi, J.
Deposited on : 2009-10-01
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	725	 3% 75% 18% 6%
2	B	44	 2% 64% 16% 20%
3	C	2	 50% 50%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 5803 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Reelin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	685	Total	C	N	O	S	0	2	0
			5419	3441	925	1017	36			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1946	GLY	-	expression tag	UNP Q60841
A	1947	ARG	-	expression tag	UNP Q60841
A	2101	ALA	CYS	engineered mutation	UNP Q60841
A	2662	SER	-	expression tag	UNP Q60841
A	2663	ARG	-	expression tag	UNP Q60841
A	2664	LEU	-	expression tag	UNP Q60841
A	2665	GLU	-	expression tag	UNP Q60841
A	2666	ASN	-	expression tag	UNP Q60841
A	2667	LEU	-	expression tag	UNP Q60841
A	2668	TYR	-	expression tag	UNP Q60841
A	2669	PHE	-	expression tag	UNP Q60841
A	2670	GLN	-	expression tag	UNP Q60841

- Molecule 2 is a protein called Low-density lipoprotein receptor-related protein 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	35	Total	C	N	O	S	0	0	0
			289	166	51	66	6			

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	40	GLY	-	expression tag	UNP Q14114
B	41	SER	-	expression tag	UNP Q14114
B	46	GLU	ASP	SEE REMARK 999	UNP Q14114

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	4	Total	Ca	0	0
			4	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Ca	0	0
			1	1		

- Molecule 6 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Zn	0	0
			1	1		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	18	Total	O	0	0
			18	18		
7	B	1	Total	O	0	0
			1	1		



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	56.98Å 93.84Å 73.46Å 90.00° 107.41° 90.00°	Depositor
Resolution (Å)	46.93 – 2.60 46.93 – 2.60	Depositor EDS
% Data completeness (in resolution range)	96.5 (46.93-2.60) 96.4 (46.93-2.60)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.46 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.223 , 0.296 0.225 , 0.296	Depositor DCC
R_{free} test set	1135 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	50.1	Xtriage
Anisotropy	0.150	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 51.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	5803	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, ZN, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.57	0/5569	0.83	2/7578 (0.0%)
2	B	0.56	0/293	0.85	0/394
All	All	0.57	0/5862	0.83	2/7972 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	2547	VAL	N-CA-C	-5.58	106.26	111.45
1	A	2275	ARG	N-CA-C	5.39	117.27	109.07

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5419	0	5134	79	0
2	B	289	0	231	5	0
3	C	28	0	25	0	0
4	A	42	0	39	0	0
5	A	4	0	0	0	0
5	B	1	0	0	0	0
6	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	18	0	0	1	0
7	B	1	0	0	0	0
All	All	5803	0	5429	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2221:LEU:HD12	1:A:2290:THR:HB	1.63	0.81
2:B:49:LYS:HD2	2:B:49:LYS:H	1.50	0.76
1:A:1985:TYR:O	1:A:2088:TRP:HD1	1.70	0.74
1:A:2055:LEU:HD11	1:A:2095:PHE:HD1	1.54	0.72
1:A:2416:LEU:HB3	1:A:2417:PRO:HD2	1.73	0.68
1:A:1976:GLY:HA2	1:A:2003:GLU:HG2	1.77	0.66
1:A:2195:CYS:HB3	1:A:2308:VAL:HG21	1.76	0.66
2:B:49:LYS:HD2	2:B:49:LYS:N	2.13	0.60
1:A:2611:MET:SD	1:A:2626:ILE:HD11	2.42	0.60
1:A:2022:ILE:HD13	1:A:2055:LEU:HD21	1.83	0.60
1:A:2231:MET:HB3	1:A:2277:ILE:HD13	1.85	0.58
1:A:2217:ARG:HD3	1:A:2218:ASP:O	2.04	0.57
1:A:2042:LEU:HD13	1:A:2108:TRP:CE2	2.39	0.57
1:A:2412:VAL:O	1:A:2428:GLN:HA	2.06	0.56
1:A:2630:PRO:HA	1:A:2633:LYS:HD2	1.89	0.54
1:A:2046:ARG:NH2	1:A:2100:LEU:O	2.41	0.54
1:A:2059:CYS:SG	1:A:2075:HIS:CE1	3.02	0.53
1:A:2318:ILE:C	1:A:2320:GLY:H	2.16	0.53
1:A:2055:LEU:HD11	1:A:2095:PHE:CD1	2.40	0.53
1:A:2137:CYS:O	1:A:2160:CYS:HB2	2.09	0.53
1:A:2562:LEU:CD1	1:A:2642:TRP:HB3	2.39	0.52
1:A:2074:HIS:HB2	1:A:2250:GLN:HE22	1.75	0.51
1:A:2230:PHE:HD2	1:A:2311:GLN:HB2	1.75	0.51
1:A:2563:LEU:O	1:A:2640:ARG:HA	2.10	0.51
1:A:2036:SER:HB3	1:A:2209:ASP:HB3	1.92	0.51
1:A:1983:CYS:O	1:A:1985:TYR:N	2.37	0.51
1:A:2047:ASP:CG	1:A:2050:ALA:HB3	2.36	0.50
1:A:2318:ILE:C	1:A:2320:GLY:N	2.68	0.50
1:A:2029:GLY:HA2	1:A:2120:VAL:HG22	1.94	0.49
1:A:2212:ARG:HH12	1:A:2306:PRO:HA	1.78	0.48
2:B:49:LYS:C	2:B:51:GLN:H	2.21	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2270:SER:O	1:A:2275:ARG:NH2	2.46	0.48
1:A:2206:PHE:HB3	1:A:2212:ARG:HD3	1.96	0.48
2:B:56:ASN:O	2:B:57:GLU:HB2	2.14	0.47
1:A:2268:SER:H	1:A:2271:SER:HB3	1.78	0.47
1:A:2319:SER:O	1:A:2320:GLY:C	2.57	0.47
1:A:1964:PHE:O	1:A:1964:PHE:CD2	2.68	0.46
1:A:2056:LEU:HB3	1:A:2078:SER:HB3	1.97	0.46
1:A:2229:PHE:HA	1:A:2311:GLN:O	2.16	0.46
1:A:1963:ASP:HA	1:A:1995:SER:HB2	1.98	0.45
1:A:2058:LEU:HB2	1:A:2077:SER:HB3	1.98	0.45
1:A:2325:GLU:HG3	1:A:2477:TYR:HD1	1.80	0.45
1:A:2320:GLY:O	1:A:2372:ALA:CB	2.64	0.45
1:A:2420:VAL:C	1:A:2422:CYS:N	2.72	0.45
1:A:2320:GLY:O	1:A:2372:ALA:HB2	2.17	0.45
1:A:2416:LEU:O	1:A:2418:THR:N	2.49	0.45
1:A:2595:VAL:HG21	1:A:2654:TRP:CD2	2.52	0.44
1:A:2562:LEU:HD13	1:A:2642:TRP:HB3	1.98	0.44
2:B:72:CYS:O	2:B:73:LEU:HB2	2.17	0.44
1:A:2212:ARG:NH1	1:A:2306:PRO:HA	2.33	0.44
1:A:2522:ASN:OD1	1:A:2522:ASN:N	2.42	0.44
1:A:2434:ASP:OD1	1:A:2559:CYS:HB3	2.17	0.43
1:A:2052:TRP:CZ3	1:A:2107:ARG:HG2	2.52	0.43
1:A:2074:HIS:HB2	1:A:2250:GLN:NE2	2.33	0.43
1:A:2156:SER:HB2	1:A:2162:ILE:HD11	2.00	0.43
1:A:2170:LEU:O	1:A:2313:LEU:HA	2.18	0.43
1:A:2488:GLY:HA2	1:A:2605:ILE:HD11	2.00	0.43
1:A:2022:ILE:HD12	1:A:2022:ILE:N	2.34	0.43
1:A:2292:LEU:HD12	1:A:2314:ILE:CD1	2.49	0.43
1:A:2173:ASP:HB2	1:A:2175:GLU:HG2	2.01	0.42
1:A:2230:PHE:CD2	1:A:2311:GLN:HB2	2.54	0.42
1:A:2038:ASP:N	1:A:2039:PRO:CD	2.82	0.42
1:A:2328:PHE:O	1:A:2353:ASP:HB2	2.18	0.42
1:A:2271:SER:C	1:A:2273:VAL:H	2.28	0.42
1:A:2001:SER:OG	1:A:2119:PRO:O	2.28	0.42
1:A:2044:PHE:HA	1:A:2105:ARG:O	2.20	0.42
1:A:2538:GLY:HA2	1:A:2554:HIS:O	2.20	0.42
1:A:2273:VAL:HB	1:A:2275:ARG:NH2	2.35	0.41
1:A:2601:VAL:HG22	7:A:6004:HOH:O	2.19	0.41
1:A:2400:TYR:CE2	1:A:2409:HIS:HB2	2.56	0.41
1:A:2300:ASN:N	1:A:2300:ASN:OD1	2.53	0.41
1:A:2357:PHE:HB3	1:A:2364:ARG:HD3	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2029:GLY:HA2	1:A:2120:VAL:CG2	2.51	0.41
1:A:2213:MET:HA	1:A:2294:TRP:O	2.21	0.41
1:A:2249:LEU:HD13	1:A:2294:TRP:CE2	2.56	0.41
1:A:2000:VAL:HA	1:A:2121:THR:HG23	2.02	0.41
1:A:2591:ARG:HA	1:A:2615:TYR:CD2	2.55	0.41
1:A:2616:ASP:N	1:A:2616:ASP:OD1	2.51	0.40
1:A:2174:PHE:CZ	1:A:2312:ILE:HD12	2.56	0.40
1:A:2245:GLN:HA	1:A:2246:PRO:HD3	1.97	0.40
1:A:2381:ILE:HG22	1:A:2476:VAL:HG22	2.04	0.40
1:A:2424:ARG:HD3	1:A:2424:ARG:HA	1.76	0.40
1:A:2287:SER:C	1:A:2289:SER:H	2.30	0.40
1:A:2397:GLU:OE2	1:A:2397:GLU:HA	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	679/725 (94%)	631 (93%)	44 (6%)	4 (1%)	21	42
2	B	33/44 (75%)	28 (85%)	4 (12%)	1 (3%)	3	6
All	All	712/769 (93%)	659 (93%)	48 (7%)	5 (1%)	18	38

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	2272	ASN
1	A	2320	GLY
1	A	2421	GLU
1	A	2417	PRO
2	B	50	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	599/634 (94%)	581 (97%)	18 (3%)	36	64
2	B	35/41 (85%)	35 (100%)	0	100	100
All	All	634/675 (94%)	616 (97%)	18 (3%)	38	66

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1963	ASP
1	A	2032	THR
1	A	2181	ASP
1	A	2195	CYS
1	A	2198	LEU
1	A	2253	LEU
1	A	2262	LEU
1	A	2305	SER
1	A	2317	ASN
1	A	2366	VAL
1	A	2414	ASP
1	A	2501	GLU
1	A	2510	GLU
1	A	2513	THR
1	A	2522	ASN
1	A	2559	CYS
1	A	2587	THR
1	A	2627	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2245	GLN
1	A	2590	ASN
1	A	2625	ASN
1	A	2651	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	2658	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	C	1	1,3	14,14,15	0.54	0	17,19,21	0.99	2 (11%)
3	NAG	C	2	3	14,14,15	0.54	0	17,19,21	0.75	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	C	2	3	-	4/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1	NAG	C4-C3-C2	2.28	114.36	111.02
3	C	1	NAG	O5-C1-C2	-2.24	107.82	111.29

There are no chirality outliers.


All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	2	NAG	C8-C7-N2-C2
3	C	2	NAG	O7-C7-N2-C2
3	C	2	NAG	C4-C5-C6-O6
3	C	2	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 6 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	A	3001	1	14,14,15	0.55	0	17,19,21	1.19	1 (5%)
4	NAG	A	3005	1	14,14,15	0.54	0	17,19,21	1.12	1 (5%)
4	NAG	A	3004	1	14,14,15	0.53	0	17,19,21	1.51	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	3001	1	-	2/6/23/26	0/1/1/1
4	NAG	A	3005	1	-	4/6/23/26	0/1/1/1
4	NAG	A	3004	1	-	1/6/23/26	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	3004	NAG	C1-O5-C5	5.55	119.62	112.19
4	A	3001	NAG	C1-O5-C5	3.63	117.05	112.19
4	A	3005	NAG	C1-O5-C5	2.58	115.65	112.19

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	3005	NAG	O5-C5-C6-O6
4	A	3001	NAG	O5-C5-C6-O6
4	A	3005	NAG	C4-C5-C6-O6
4	A	3005	NAG	C8-C7-N2-C2
4	A	3005	NAG	O7-C7-N2-C2
4	A	3001	NAG	C4-C5-C6-O6
4	A	3004	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	685/725 (94%)	0.54	23 (3%) 48 42	29, 57, 64, 71	2 (0%)
2	B	35/44 (79%)	0.45	1 (2%) 53 48	54, 57, 60, 60	0
All	All	720/769 (93%)	0.54	24 (3%) 49 43	29, 57, 64, 71	2 (0%)

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	81	CYS	4.8
1	A	2299	GLU	4.2
1	A	2319	SER	4.1
1	A	1964	PHE	3.5
1	A	2059	CYS	3.1
1	A	2141	GLY	3.0
1	A	2316	GLY	3.0
1	A	2186	MET	2.8
1	A	2209	ASP	2.8
1	A	2526	ALA	2.8
1	A	2468	GLN	2.7
1	A	2460	HIS	2.7
1	A	2184	LEU	2.4
1	A	2300	ASN	2.3
1	A	2157	GLY	2.3
1	A	2237	LYS	2.2
1	A	2055	LEU	2.2
1	A	2085	THR	2.1
1	A	1953	ASN	2.1
1	A	2154	GLY	2.1
1	A	2198	LEU	2.1
1	A	2527	PRO	2.1
1	A	2661	ILE	2.1
1	A	2172	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	C	2	14/15	0.61	0.18	39,43,44,44	0
3	NAG	C	1	14/15	0.67	0.18	44,46,47,47	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	A	3004	14/15	0.74	0.15	63,65,67,67	0
4	NAG	A	3001	14/15	0.83	0.12	50,52,53,53	0
4	NAG	A	3005	14/15	0.84	0.11	57,59,63,63	0
5	CA	A	4001	1/1	0.92	0.07	52,52,52,52	0
6	ZN	A	4005	1/1	0.95	0.11	118,118,118,118	0
5	CA	A	4004	1/1	0.96	0.05	53,53,53,53	0
5	CA	A	4003	1/1	0.97	0.05	55,55,55,55	0
5	CA	B	5001	1/1	0.98	0.04	55,55,55,55	0
5	CA	A	4002	1/1	0.98	0.07	56,56,56,56	0

6.5 Other polymers [i](#)

There are no such residues in this entry.