



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:35 PM UTC

PDB ID : 5A78 / pdb_00005a78
Title : Crystal structure of the homing endonuclease I-CvuI in complex with I- CreI target (C1221) in the presence of 2 mM Mg revealing DNA not cleaved
Authors : Molina, R.; Redondo, P.; LopezMendez, B.; Villate, M.; Merino, N.; Blanco, F.J.; Valton, J.; Grizot, S.; Duchateau, P.; Prieto, J.; Montoya, G.
Deposited on : 2015-07-03
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	172	
1	B	172	
2	C	24	
2	D	24	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3474 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA ENDONUCLEASE I-CVUI.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	157	Total	C	N	O	S	0	0	0
			1233	791	215	221	6			
1	B	158	Total	C	N	O	S	0	0	0
			1239	794	216	223	6			

There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	ALA	-	expression tag	UNP P56347
A	163	SER	-	expression tag	UNP P56347
A	164	ALA	-	expression tag	UNP P56347
A	165	ALA	-	expression tag	UNP P56347
A	166	LEU	-	expression tag	UNP P56347
A	167	GLU	-	expression tag	UNP P56347
A	168	HIS	-	expression tag	UNP P56347
A	169	HIS	-	expression tag	UNP P56347
A	170	HIS	-	expression tag	UNP P56347
A	171	HIS	-	expression tag	UNP P56347
A	172	HIS	-	expression tag	UNP P56347
A	173	HIS	-	expression tag	UNP P56347
A	54	GLN	ARG	conflict	UNP P56347
B	2	ALA	-	expression tag	UNP P56347
B	163	SER	-	expression tag	UNP P56347
B	164	ALA	-	expression tag	UNP P56347
B	165	ALA	-	expression tag	UNP P56347
B	166	LEU	-	expression tag	UNP P56347
B	167	GLU	-	expression tag	UNP P56347
B	168	HIS	-	expression tag	UNP P56347
B	169	HIS	-	expression tag	UNP P56347
B	170	HIS	-	expression tag	UNP P56347
B	171	HIS	-	expression tag	UNP P56347
B	172	HIS	-	expression tag	UNP P56347
B	173	HIS	-	expression tag	UNP P56347

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	54	GLN	ARG	conflict	UNP P56347

- Molecule 2 is a DNA chain called 24MER DNA, 5'-D(*TP*CP*AP*AP*AP*AP*CP*GP*TP*CP*GP*TP*AP *CP*GP*AP*CP*GP*TP*TP*TP*TP*GP*A)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	24	Total 489	C 235	N 89	O 142	P 23	0	0	0
2	D	24	Total 489	C 235	N 89	O 142	P 23	0	0	0

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total 1	Mg 1	0	0
3	D	1	Total 1	Mg 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	12	Total 12	O 12	0	0
4	B	6	Total 6	O 6	0	0
4	C	4	Total 4	O 4	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: DNA ENDONUCLEASE I-CVUI



• Molecule 1: DNA ENDONUCLEASE I-CVUI



• Molecule 2: 24MER DNA, 5'-D(*TP*CP*AP*AP*AP*AP*CP*GP*TP*CP*GP*TP*AP *CP *GP*AP*CP*GP*TP*TP*TP*TP*GP*A)-3'



• Molecule 2: 24MER DNA, 5'-D(*TP*CP*AP*AP*AP*AP*CP*GP*TP*CP*GP*TP*AP *CP *GP*AP*CP*GP*TP*TP*TP*TP*GP*A)-3'



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	48.57Å 95.08Å 52.15Å 90.00° 103.71° 90.00°	Depositor
Resolution (Å)	47.54 – 2.50 47.54 – 2.50	Depositor EDS
% Data completeness (in resolution range)	97.9 (47.54-2.50) 98.7 (47.54-2.50)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.86 (at 2.51Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.215 , 0.280 0.226 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	36.9	Xtriage
Anisotropy	0.970	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 37.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3474	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.81	0/1249	1.03	4/1683 (0.2%)
1	B	0.73	0/1255	1.31	14/1691 (0.8%)
2	C	0.37	0/548	0.65	0/844
2	D	0.36	0/548	0.63	0/844
All	All	0.67	0/3600	1.03	18/5062 (0.4%)

There are no bond length outliers.

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	142	LYS	N-CA-C	-23.43	81.33	110.61
1	B	141	GLY	N-CA-C	-13.95	80.11	113.18
1	B	140	ASP	CB-CA-C	-12.35	92.07	111.83
1	B	57	ILE	CA-C-N	-8.54	106.19	121.14
1	B	57	ILE	C-N-CA	-8.54	106.19	121.14
1	A	142	LYS	N-CA-CB	-7.14	98.93	111.42
1	B	142	LYS	N-CA-CB	-6.94	101.93	111.65
1	B	68	GLY	N-CA-C	6.81	118.42	111.95
1	B	140	ASP	N-CA-C	-6.70	96.86	108.52
1	B	146	ILE	N-CA-C	6.55	117.28	107.51
1	B	144	ARG	N-CA-C	5.72	118.06	108.96
1	A	68	GLY	N-CA-C	5.60	119.68	112.18
1	A	144	ARG	N-CA-C	5.53	118.55	109.76
1	A	142	LYS	O-C-N	5.29	127.75	122.30
1	B	61	ILE	CA-C-N	5.11	127.08	120.44
1	B	61	ILE	C-N-CA	5.11	127.08	120.44
1	B	121	ASP	CA-C-N	5.00	124.67	119.56
1	B	121	ASP	C-N-CA	5.00	124.67	119.56

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1233	0	1306	44	1
1	B	1239	0	1309	27	0
2	C	489	0	272	11	0
2	D	489	0	272	7	1
3	B	1	0	0	0	0
3	D	1	0	0	0	0
4	A	12	0	0	5	0
4	B	6	0	0	1	0
4	C	4	0	0	1	0
All	All	3474	0	3159	76	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:143:LYS:NZ	2:C:515:DG:N2	1.64	1.40
4:A:2009:HOH:O	1:B:22:ALA:HA	1.68	0.94
4:A:2003:HOH:O	2:C:515:DG:H5'	1.68	0.93
1:A:143:LYS:HZ2	2:C:515:DG:N2	1.34	0.92
4:B:2001:HOH:O	2:D:615:DG:H5'	1.76	0.86
1:A:6:ASN:ND2	1:A:9:ASP:OD1	2.12	0.83
1:A:143:LYS:NZ	2:C:515:DG:C2	2.47	0.81
1:B:126:MET:HG2	1:B:162:VAL:HG11	1.63	0.80
2:C:502:DC:O2	4:C:2001:HOH:O	2.00	0.79
1:A:21:ASP:O	4:A:2002:HOH:O	2.01	0.78
1:A:75:ASP:OD2	4:A:2011:HOH:O	2.04	0.75
1:B:146:ILE:HG22	2:D:618:DG:OP1	1.86	0.75
1:A:139:THR:C	1:A:141:GLY:H	1.93	0.74
1:A:126:MET:HG2	1:A:162:VAL:HG11	1.71	0.72
1:B:40:TYR:HE1	1:B:146:ILE:HD11	1.58	0.69

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:139:THR:C	1:A:141:GLY:N	2.53	0.66
1:A:147:LEU:O	1:A:150:ASN:N	2.30	0.65
1:B:40:TYR:CE1	1:B:146:ILE:HD11	2.32	0.65
1:A:6:ASN:ND2	1:A:9:ASP:H	1.96	0.64
2:C:507:DC:H2''	2:C:508:DG:H5'	1.80	0.63
1:A:139:THR:O	1:A:141:GLY:N	2.33	0.62
1:A:149:GLU:O	1:A:152:ARG:N	2.32	0.61
1:B:146:ILE:HG23	1:B:146:ILE:O	2.00	0.60
1:A:143:LYS:NZ	2:C:515:DG:H22	1.87	0.60
1:B:152:ARG:HH12	1:B:156:LYS:NZ	2.00	0.59
1:B:152:ARG:HH12	1:B:156:LYS:HZ2	1.50	0.59
1:A:6:ASN:HD21	1:A:9:ASP:H	1.49	0.59
1:A:38:LEU:HD11	1:A:120:LYS:O	2.05	0.57
1:A:27:ASN:OD1	1:A:28:ALA:N	2.37	0.55
1:A:30:ILE:HB	1:A:146:ILE:HG22	1.89	0.55
1:A:96:LEU:HD21	1:A:106:ALA:HB3	1.88	0.54
4:A:2006:HOH:O	2:D:603:DA:N7	2.33	0.54
1:A:30:ILE:HB	1:A:146:ILE:CG2	2.38	0.53
1:A:86:THR:HG22	1:A:86:THR:O	2.09	0.52
1:B:89:GLN:O	1:B:93:GLU:HG3	2.09	0.52
1:B:90:THR:HG22	1:B:94:LYS:NZ	2.25	0.52
1:A:59:LEU:HD23	1:A:70:VAL:HG21	1.91	0.51
1:A:145:THR:HG22	1:A:145:THR:O	2.11	0.51
1:B:135:VAL:O	1:B:139:THR:HG23	2.11	0.50
1:B:50:GLN:HG2	1:B:57:ILE:CG2	2.43	0.49
1:B:49:PHE:CE2	2:D:615:DG:H2'	2.48	0.48
1:A:133:ASP:OD2	1:A:148:ALA:CB	2.61	0.48
1:A:144:ARG:NH1	2:C:517:DC:H5'	2.29	0.48
2:D:607:DC:H2''	2:D:608:DG:H5'	1.96	0.47
1:A:101:LEU:HD21	1:B:57:ILE:HD13	1.96	0.47
1:A:147:LEU:HB2	1:A:149:GLU:HG2	1.95	0.47
1:A:6:ASN:ND2	1:A:6:ASN:C	2.73	0.46
1:A:6:ASN:C	1:A:6:ASN:HD22	2.24	0.46
1:B:32:SER:O	1:B:33:ARG:HD2	2.15	0.45
1:A:134:LYS:O	1:A:138:LEU:HG	2.15	0.45
1:A:73:ARG:CD	1:A:79:GLU:HG3	2.47	0.45
1:A:101:LEU:CD2	1:B:57:ILE:HD13	2.47	0.45
1:B:59:LEU:HD23	1:B:70:VAL:HG21	1.98	0.45
2:C:506:DA:C4	2:C:507:DC:C6	3.05	0.45
1:A:141:GLY:C	1:A:142:LYS:HG3	2.43	0.44
1:A:23:ASP:CG	1:B:22:ALA:HB1	2.43	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:504:DA:H2''	2:C:505:DA:C8	2.53	0.44
1:B:40:TYR:CD2	1:B:151:VAL:HG22	2.53	0.43
1:A:109:VAL:O	1:A:113:ILE:HG13	2.18	0.43
1:B:90:THR:HG22	1:B:94:LYS:HZ1	1.83	0.43
1:B:38:LEU:HD11	1:B:120:LYS:O	2.18	0.43
1:B:152:ARG:HH22	1:B:156:LYS:HZ3	1.67	0.43
1:A:95:LEU:O	1:A:96:LEU:C	2.63	0.42
1:A:12:LYS:HG2	1:A:64:ILE:HD13	2.02	0.42
1:A:57:ILE:HB	1:B:101:LEU:HD21	2.02	0.42
2:C:506:DA:C5	2:C:507:DC:C5	3.07	0.42
2:D:604:DA:H2''	2:D:605:DA:C8	2.55	0.42
1:A:147:LEU:CD2	1:A:147:LEU:N	2.83	0.42
1:A:101:LEU:HD22	1:B:54:GLN:OE1	2.20	0.42
1:B:91:THR:HA	1:B:94:LYS:HE2	2.01	0.41
1:A:105:GLN:HG3	1:A:138:LEU:HD12	2.03	0.41
1:A:12:LYS:HG2	1:A:64:ILE:CD1	2.51	0.41
1:A:22:ALA:HB3	1:B:22:ALA:HB3	2.01	0.41
1:A:91:THR:O	1:A:94:LYS:HB2	2.21	0.41
2:D:606:DA:C4	2:D:607:DC:C6	3.09	0.41
1:B:20:VAL:O	1:B:24:GLY:N	2.42	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:157:LYS:NZ	2:D:620:DT:OP1[2_445]	2.15	0.05

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	155/172 (90%)	146 (94%)	9 (6%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	156/172 (91%)	153 (98%)	3 (2%)	0	100	100
All	All	311/344 (90%)	299 (96%)	12 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	137/149 (92%)	124 (90%)	13 (10%)	8	17
1	B	138/149 (93%)	122 (88%)	16 (12%)	5	11
All	All	275/298 (92%)	246 (90%)	29 (10%)	6	14

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	ASN
1	A	12	LYS
1	A	33	ARG
1	A	34	GLU
1	A	37	LEU
1	A	51	SER
1	A	75	ASP
1	A	83	VAL
1	A	144	ARG
1	A	146	ILE
1	A	147	LEU
1	A	152	ARG
1	A	156	LYS
1	B	29	GLN
1	B	35	ASP
1	B	57	ILE
1	B	69	THR
1	B	83	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	100	GLN
1	B	110	LEU
1	B	114	LYS
1	B	134	LYS
1	B	139	THR
1	B	144	ARG
1	B	146	ILE
1	B	147	LEU
1	B	149	GLU
1	B	152	ARG
1	B	153	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	6	ASN
1	A	29	GLN
1	A	62	GLN
1	A	105	GLN
1	B	62	GLN
1	B	118	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	157/172 (91%)	1.14	21 (13%) 7 5	24, 41, 59, 78	0
1	B	158/172 (91%)	1.36	29 (18%) 3 2	24, 43, 62, 75	0
2	C	24/24 (100%)	0.20	0 100 100	34, 44, 54, 56	0
2	D	24/24 (100%)	0.32	0 100 100	36, 43, 51, 53	0
All	All	363/392 (92%)	1.12	50 (13%) 6 5	24, 42, 60, 78	0

All (50) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	142	LYS	7.5
1	B	141	GLY	6.3
1	A	142	LYS	5.8
1	B	153	GLU	5.2
1	B	7	PHE	5.1
1	B	140	ASP	4.6
1	A	7	PHE	4.5
1	A	141	GLY	4.4
1	A	150	ASN	4.3
1	B	6	ASN	3.8
1	B	163	SER	3.7
1	A	86	THR	3.7
1	A	153	GLU	3.5
1	A	162	VAL	3.4
1	A	143	LYS	3.3
1	B	110	LEU	3.3
1	B	143	LYS	3.3
1	A	149	GLU	3.2
1	A	145	THR	3.2
1	B	60	ASP	3.2
1	B	162	VAL	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	10	GLN	2.9
1	A	8	HIS	2.9
1	A	148	ALA	2.8
1	B	34	GLU	2.8
1	A	84	GLY	2.7
1	A	60	ASP	2.7
1	A	85	GLY	2.6
1	A	34	GLU	2.6
1	B	74	ASN	2.6
1	A	53	THR	2.5
1	B	152	ARG	2.5
1	A	146	ILE	2.5
1	B	159	GLY	2.5
1	A	140	ASP	2.5
1	A	83	VAL	2.4
1	B	84	GLY	2.3
1	B	23	ASP	2.3
1	B	8	HIS	2.3
1	B	149	GLU	2.3
1	A	139	THR	2.3
1	B	139	THR	2.3
1	B	147	LEU	2.3
1	B	32	SER	2.2
1	B	93	GLU	2.1
1	B	53	THR	2.1
1	B	133	ASP	2.1
1	B	50	GLN	2.0
1	B	161	VAL	2.0
1	B	90	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MG	B	1164	1/1	0.83	0.11	51,51,51,51	0
3	MG	D	1625	1/1	0.83	0.15	53,53,53,53	0

6.5 Other polymers [i](#)

There are no such residues in this entry.