



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 22, 2026 – 01:47 AM UTC

PDB ID : 5A7D / pdb_00005a7d
Title : Tetrameric assembly of LGN with Inscuteable
Authors : Culurgioni, S.; Mari, S.; Bonetto, G.; Gallini, S.; Brennich, M.; Round, A.;
Mapelli, M.
Deposited on : 2015-07-07
Resolution : 3.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1001 (3.44-3.36)
Clashscore	190562	1022 (3.44-3.36)
Ramachandran outliers	187476	1012 (3.44-3.36)
Sidechain outliers	187428	1012 (3.44-3.36)
RSRZ outliers	180081	1001 (3.44-3.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	382	 2% 76% 14% 7%
1	C	382	 2% 80% 10% 9%
1	D	382	 79% 11% 10%
1	E	382	 2% 77% 15% 8%
1	F	382	 81% 9% 9%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	382	<p>% 78% 12% • 9%</p>
1	H	382	<p>78% 11% • 9%</p>
1	I	382	<p>% 79% 12% • 9%</p>
2	L	341	<p>3% 69% 14% 17%</p>
2	M	341	<p>2% 63% 12% • 24%</p>
2	N	341	<p>71% 12% 17%</p>
2	O	341	<p>2% 68% 15% • 16%</p>
2	P	341	<p>% 71% 13% • 16%</p>
2	Q	341	<p>% 65% 17% • 18%</p>
2	R	341	<p>% 77% 11% • 9%</p>
2	S	341	<p>% 69% 14% • 16%</p>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 38980 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PINS.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	354	Total 2742	C 1695	N 513	O 521	S 13	0	0	0
1	C	348	Total 2685	C 1655	N 504	O 513	S 13	0	0	0
1	D	345	Total 2643	C 1635	N 489	O 506	S 13	0	0	0
1	E	352	Total 2616	C 1618	N 489	O 497	S 12	0	0	0
1	F	346	Total 2676	C 1652	N 502	O 509	S 13	0	0	0
1	G	346	Total 2679	C 1655	N 504	O 507	S 13	0	0	0
1	H	347	Total 2668	C 1650	N 499	O 506	S 13	0	0	0
1	I	346	Total 2608	C 1615	N 481	O 499	S 13	0	0	0

- Molecule 2 is a protein called INSCUTEABLE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	L	284	Total 2127	C 1329	N 386	O 402	S 10	0	0	0
2	M	259	Total 1977	C 1237	N 361	O 369	S 10	0	0	0
2	N	283	Total 2228	C 1395	N 407	O 416	S 10	0	0	0
2	O	288	Total 2250	C 1410	N 409	O 421	S 10	0	0	0
2	P	288	Total 2233	C 1396	N 406	O 421	S 10	0	0	0
2	Q	281	Total 2215	C 1388	N 401	O 416	S 10	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	R	311	Total	C	N	O	S	0	0	0
			2386	1490	439	447	10			
2	S	287	Total	C	N	O	S	0	0	0
			2243	1405	408	420	10			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total	O	0	0
			2	2		
3	H	2	Total	O	0	0
			2	2		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: PINS




- Molecule 1: PINS




- Molecule 1: PINS





- Molecule 1: PINS









• Molecule 1: PINS




• Molecule 1: PINS




• Molecule 1: PINS




• Molecule 1: PINS







• Molecule 2: INSCUTEABLE



• Molecule 2: INSCUTEABLE




• Molecule 2: INSCUTEABLE





- Molecule 2: INSCUTEABLE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	128.19Å 212.58Å 280.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	75.58 – 3.40 75.58 – 3.40	Depositor EDS
% Data completeness (in resolution range)	99.8 (75.58-3.40) 99.9 (75.58-3.40)	Depositor EDS
R_{merge}	0.23	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.17 (at 3.41Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.209 , 0.250 0.212 , 0.252	Depositor DCC
R_{free} test set	5219 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	84.1	Xtrriage
Anisotropy	0.393	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 70.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	38980	wwPDB-VP
Average B, all atoms (Å ²)	77.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	B	0.26	0/2787	0.65	0/3748
1	C	0.28	0/2729	0.66	0/3673
1	D	0.26	0/2688	0.67	0/3623
1	E	0.26	0/2657	0.67	0/3583
1	F	0.26	0/2721	0.67	0/3662
1	G	0.26	0/2724	0.67	0/3665
1	H	0.27	0/2713	0.68	2/3654 (0.1%)
1	I	0.26	0/2650	0.68	0/3576
2	L	0.28	0/2149	0.72	1/2914 (0.0%)
2	M	0.29	0/1999	0.71	1/2710 (0.0%)
2	N	0.28	0/2254	0.70	0/3049
2	O	0.27	0/2278	0.69	0/3085
2	P	0.28	0/2260	0.70	0/3063
2	Q	0.27	0/2242	0.68	1/3034 (0.0%)
2	R	0.28	0/2416	0.71	1/3273 (0.0%)
2	S	0.28	0/2270	0.69	0/3074
All	All	0.27	0/39537	0.68	6/53386 (0.0%)

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	L	603	PRO	N-CA-CB	7.06	110.67	103.25
2	R	603	PRO	N-CA-CB	6.92	110.52	103.25
1	H	74	THR	CA-C-N	5.91	132.33	121.70
1	H	74	THR	C-N-CA	5.91	132.33	121.70
2	M	581	ARG	CB-CA-C	-5.26	110.53	116.63

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2742	0	2647	34	0
1	C	2685	0	2569	25	0
1	D	2643	0	2503	20	0
1	E	2616	0	2438	31	0
1	F	2676	0	2560	20	0
1	G	2679	0	2570	26	0
1	H	2668	0	2545	28	0
1	I	2608	0	2457	22	0
2	L	2127	0	2088	19	0
2	M	1977	0	1958	20	0
2	N	2228	0	2268	20	0
2	O	2250	0	2285	30	0
2	P	2233	0	2253	26	0
2	Q	2215	0	2251	25	0
2	R	2386	0	2372	22	0
2	S	2243	0	2277	20	0
3	B	2	0	0	0	0
3	H	2	0	0	0	0
All	All	38980	0	38041	342	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 342 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:M:567:ILE:HG23	2:M:587:LEU:HD21	1.63	0.80
1:B:74:THR:OG1	1:B:75:GLU:N	2.16	0.76
2:O:523:LEU:HD22	2:O:563:ILE:HD11	1.70	0.74
2:P:342:TYR:N	2:P:343:THR:HA	2.04	0.73
1:E:47:ALA:HB1	2:O:334:ILE:HD13	1.71	0.72

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	352/382 (92%)	329 (94%)	17 (5%)	6 (2%)	7	28
1	C	346/382 (91%)	334 (96%)	11 (3%)	1 (0%)	36	65
1	D	343/382 (90%)	332 (97%)	10 (3%)	1 (0%)	36	65
1	E	348/382 (91%)	330 (95%)	16 (5%)	2 (1%)	21	50
1	F	344/382 (90%)	334 (97%)	9 (3%)	1 (0%)	36	65
1	G	344/382 (90%)	331 (96%)	13 (4%)	0	100	100
1	H	345/382 (90%)	337 (98%)	7 (2%)	1 (0%)	36	65
1	I	344/382 (90%)	330 (96%)	9 (3%)	5 (2%)	8	30
2	L	278/341 (82%)	255 (92%)	20 (7%)	3 (1%)	11	38
2	M	251/341 (74%)	236 (94%)	13 (5%)	2 (1%)	16	44
2	N	279/341 (82%)	263 (94%)	16 (6%)	0	100	100
2	O	286/341 (84%)	270 (94%)	12 (4%)	4 (1%)	9	31
2	P	286/341 (84%)	273 (96%)	12 (4%)	1 (0%)	36	65
2	Q	277/341 (81%)	269 (97%)	6 (2%)	2 (1%)	18	47
2	R	307/341 (90%)	283 (92%)	17 (6%)	7 (2%)	5	23
2	S	285/341 (84%)	266 (93%)	10 (4%)	9 (3%)	3	17
All	All	5015/5784 (87%)	4772 (95%)	198 (4%)	45 (1%)	14	42

5 of 45 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	183	PRO
1	E	351	ILE
2	M	408	VAL
2	S	343	THR
1	B	75	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	271/298 (91%)	250 (92%)	21 (8%)	12	37
1	C	263/298 (88%)	255 (97%)	8 (3%)	36	59
1	D	256/298 (86%)	244 (95%)	12 (5%)	23	50
1	E	241/298 (81%)	228 (95%)	13 (5%)	20	47
1	F	262/298 (88%)	256 (98%)	6 (2%)	44	63
1	G	262/298 (88%)	248 (95%)	14 (5%)	20	47
1	H	259/298 (87%)	249 (96%)	10 (4%)	28	53
1	I	247/298 (83%)	236 (96%)	11 (4%)	24	51
2	L	216/292 (74%)	200 (93%)	16 (7%)	13	38
2	M	206/292 (70%)	194 (94%)	12 (6%)	18	45
2	N	242/292 (83%)	230 (95%)	12 (5%)	22	49
2	O	243/292 (83%)	224 (92%)	19 (8%)	11	37
2	P	241/292 (82%)	226 (94%)	15 (6%)	16	43
2	Q	242/292 (83%)	225 (93%)	17 (7%)	14	40
2	R	248/292 (85%)	227 (92%)	21 (8%)	10	33
2	S	242/292 (83%)	224 (93%)	18 (7%)	13	38
All	All	3941/4720 (84%)	3716 (94%)	225 (6%)	18	45

5 of 225 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	M	456	LEU
2	S	542	LEU
2	O	444	ILE
2	S	515	GLN
2	R	495	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 53 such sidechains are listed below:

Mol	Chain	Res	Type
1	I	346	ASN
2	N	442	HIS
2	S	516	ASN
2	L	317	GLN
2	L	553	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	B	354/382 (92%)	-0.14	3 (0%) 82 71	35, 56, 99, 127	0
1	C	348/382 (91%)	0.07	2 (0%) 85 75	46, 79, 126, 152	0
1	D	345/382 (90%)	-0.04	1 (0%) 90 84	51, 89, 121, 166	0
1	E	352/382 (92%)	0.33	9 (2%) 57 42	46, 104, 147, 191	0
1	F	346/382 (90%)	-0.11	0 100 100	31, 70, 132, 150	0
1	G	346/382 (90%)	-0.07	2 (0%) 85 75	50, 84, 113, 154	0
1	H	347/382 (90%)	-0.14	1 (0%) 90 84	31, 64, 119, 132	0
1	I	346/382 (90%)	0.31	5 (1%) 73 59	50, 103, 147, 161	0
2	L	284/341 (83%)	0.13	9 (3%) 50 37	43, 75, 125, 150	0
2	M	259/341 (75%)	0.19	8 (3%) 51 38	45, 82, 115, 132	0
2	N	283/341 (82%)	-0.12	1 (0%) 88 81	42, 69, 109, 135	0
2	O	288/341 (84%)	-0.09	7 (2%) 59 45	33, 62, 124, 161	0
2	P	288/341 (84%)	-0.13	3 (1%) 79 66	28, 55, 115, 176	0
2	Q	281/341 (82%)	-0.04	2 (0%) 84 73	43, 68, 99, 117	0
2	R	311/341 (91%)	-0.13	4 (1%) 75 61	28, 52, 134, 153	0
2	S	287/341 (84%)	-0.08	2 (0%) 84 73	36, 67, 128, 153	0
All	All	5065/5784 (87%)	-0.00	59 (1%) 76 63	28, 74, 130, 191	0

The worst 5 of 59 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	L	345	THR	4.0
1	E	379	THR	3.6
2	N	570	VAL	3.6
1	E	344	LEU	3.5
1	G	40	SER	3.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.