



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 02:25 PM UTC

PDB ID : 6A7W / pdb_00006a7w
Title : Structure of a catalytic domain of the colistin resistance enzyme
Authors : Wang, X.D.; Chai, Y.; Qi, J.X.; Gao, G.F.
Deposited on : 2018-07-04
Resolution : 2.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3580 (3.00-2.96)
Clashscore	190562	3904 (3.00-2.96)
Ramachandran outliers	187476	3761 (3.00-2.96)
Sidechain outliers	187428	3764 (3.00-2.96)
RSRZ outliers	180081	3579 (3.00-2.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	324	 83% 14% .
1	B	324	 84% 13% ..

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5084 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative integral membrane protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	324	Total	C	N	O	S	0	0	0
			2541	1600	427	500	14			
1	B	324	Total	C	N	O	S	0	0	0
			2541	1600	427	500	14			


- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative integral membrane protein



- Molecule 1: Putative integral membrane protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	46.44Å 82.30Å 79.91Å 90.00° 97.39° 90.00°	Depositor
Resolution (Å)	41.15 – 2.99 41.15 – 2.99	Depositor EDS
% Data completeness (in resolution range)	96.3 (41.15-2.99) 87.9 (41.15-2.99)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.19	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.54 (at 3.01Å)	Xtriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, R_{free}	0.180 , 0.220 (Not available) , 0.211	Depositor DCC
R_{free} test set	585 reflections (4.78%)	wwPDB-VP
Wilson B-factor (Å ²)	54.7	Xtriage
Anisotropy	0.102	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 24.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5084	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.51	3/2600 (0.1%)	0.73	13/3533 (0.4%)
1	B	0.32	1/2600 (0.0%)	0.53	6/3533 (0.2%)
All	All	0.43	4/5200 (0.1%)	0.64	19/7066 (0.3%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	475	LEU	C-O	-5.71	1.16	1.24
1	A	476	HIS	C-O	-5.66	1.16	1.24
1	A	476	HIS	CA-C	-5.27	1.45	1.52
1	B	476	HIS	C-O	-5.06	1.17	1.24

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	474	TYR	N-CA-C	15.43	134.69	112.94
1	A	475	LEU	N-CA-CB	12.65	122.41	109.51
1	A	408	PHE	O-C-N	-9.36	112.55	123.22
1	A	331	LYS	CA-C-N	-8.31	109.46	122.51
1	A	331	LYS	C-N-CA	-8.31	109.46	122.51
1	B	331	LYS	CA-C-N	-7.93	111.45	122.63
1	B	331	LYS	C-N-CA	-7.93	111.45	122.63
1	A	475	LEU	N-CA-C	-7.88	99.43	108.49
1	B	409	THR	C-N-CD	7.35	136.78	120.60
1	A	409	THR	C-N-CD	6.44	134.76	120.60
1	A	329	ASP	O-C-N	-6.33	116.46	123.04
1	B	331	LYS	N-CA-C	-6.19	106.14	113.38
1	A	393	HIS	N-CA-C	6.04	117.68	110.19
1	A	474	TYR	CA-C-N	5.69	130.09	120.81
1	A	474	TYR	C-N-CA	5.69	130.09	120.81

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	394	GLY	CA-C-N	-5.63	113.81	119.56
1	B	394	GLY	C-N-CA	-5.63	113.81	119.56
1	A	474	TYR	CB-CA-C	-5.33	101.48	110.64
1	A	331	LYS	N-CA-C	-5.11	106.99	113.23

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2541	0	2416	33	1
1	B	2541	0	2416	28	1
2	A	1	0	0	0	0
2	B	1	0	0	0	0
All	All	5084	0	4832	61	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:474:TYR:O	1:A:475:LEU:HD13	1.63	0.99
1:B:283:THR:O	1:B:331:LYS:HE3	1.80	0.80
1:A:458:MET:HE1	1:A:460:TYR:HB2	1.70	0.73
1:B:458:MET:HE1	1:B:460:TYR:HB2	1.72	0.72
1:A:393:HIS:CG	1:A:394:GLY:H	2.11	0.69
1:B:467:SER:O	1:B:474:TYR:HA	1.95	0.66
1:A:244:GLU:OE1	1:A:393:HIS:HD2	1.79	0.65
1:B:310:VAL:HB	1:B:516:THR:HG23	1.84	0.60
1:A:244:GLU:OE1	1:A:393:HIS:CD2	2.57	0.58
1:A:378:ASN:HB3	1:A:381:LYS:HE2	1.86	0.57
1:A:393:HIS:CG	1:A:394:GLY:N	2.71	0.57
1:B:268:ASP:OD2	1:B:269:GLY:N	2.38	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:283:THR:O	1:B:331:LYS:CE	2.53	0.56
1:A:466:GLU:OE1	1:A:476:HIS:ND1	2.30	0.53
1:A:270:LEU:HD23	1:A:495:TRP:HB2	1.89	0.53
1:A:284:ALA:HA	1:A:331:LYS:HD2	1.90	0.52
1:A:293:TYR:O	1:A:533:ARG:NH2	2.43	0.52
1:B:270:LEU:HD23	1:B:495:TRP:HB2	1.93	0.50
1:B:474:TYR:O	1:B:475:LEU:HB2	2.11	0.50
1:B:409:THR:HB	1:B:410:PRO:HA	1.92	0.50
1:A:239:VAL:HG23	1:A:447:LEU:HD21	1.93	0.49
1:B:467:SER:HB3	1:B:473:VAL:O	2.12	0.49
1:A:279:CYS:SG	1:A:296:GLN:HB3	2.52	0.49
1:A:393:HIS:ND1	1:A:394:GLY:N	2.60	0.49
1:B:378:ASN:HB3	1:B:381:LYS:HE2	1.93	0.49
1:A:409:THR:HB	1:A:410:PRO:HA	1.94	0.48
1:B:460:TYR:O	1:B:492:ALA:HA	2.13	0.48
1:B:467:SER:O	1:B:475:LEU:N	2.46	0.48
1:A:408:PHE:C	1:A:409:THR:HG23	2.39	0.48
1:B:261:PHE:CD1	1:B:437:ASP:HB2	2.48	0.48
1:A:416:GLU:H	1:A:416:GLU:HG2	1.60	0.47
1:B:239:VAL:HG23	1:B:447:LEU:HD21	1.95	0.47
1:B:330:SER:C	1:B:332:GLY:N	2.72	0.47
1:A:310:VAL:HB	1:A:516:THR:HG23	1.96	0.47
1:B:325:ASP:OD1	1:B:388:HIS:ND1	2.39	0.47
1:B:256:TYR:CE1	1:B:258:ARG:HB2	2.49	0.47
1:A:408:PHE:O	1:A:409:THR:CG2	2.64	0.46
1:A:261:PHE:CD1	1:A:437:ASP:HB2	2.51	0.46
1:A:531:LYS:HD3	1:A:531:LYS:HA	1.70	0.45
1:A:330:SER:O	1:A:331:LYS:HB2	2.17	0.45
1:A:503:PRO:HA	1:A:537:ILE:HA	1.97	0.45
1:A:367:MET:HB3	1:A:387:LEU:HD22	1.97	0.45
1:B:412:CYS:N	1:B:424:SER:HB3	2.32	0.45
1:A:244:GLU:HG3	1:A:245:THR:OG1	2.17	0.45
1:A:251:VAL:HG21	1:A:491:PRO:HB3	1.99	0.44
1:B:327:ASN:HB2	1:B:389:GLN:O	2.18	0.43
1:B:279:CYS:SG	1:B:296:GLN:HB3	2.58	0.43
1:A:517:PRO:HB2	1:A:535:ALA:HB2	2.00	0.43
1:A:507:ASP:OD2	1:A:507:ASP:N	2.47	0.42
1:A:286:SER:O	1:A:290:MET:HG3	2.19	0.42
1:B:245:THR:HG22	1:B:245:THR:O	2.20	0.41
1:B:393:HIS:CG	1:B:394:GLY:H	2.33	0.41
1:B:467:SER:HB2	1:B:475:LEU:O	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:296:GLN:H	1:B:296:GLN:HG3	1.44	0.41
1:B:346:LYS:HE3	1:B:346:LYS:HB2	1.92	0.41
1:B:330:SER:C	1:B:332:GLY:H	2.29	0.41
1:A:247:ARG:HB3	1:A:249:ASP:OD1	2.21	0.41
1:B:233:ARG:NH1	1:B:234:LYS:O	2.53	0.41
1:A:384:LEU:HG	1:A:386:MET:SD	2.61	0.40
1:A:245:THR:HG23	1:A:466:GLU:HB2	2.03	0.40
1:A:327:ASN:HB2	1:A:389:GLN:O	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:471:ASN:ND2	1:B:331:LYS:O[2_658]	1.88	0.32

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	322/324 (99%)	307 (95%)	14 (4%)	1 (0%)	36	67
1	B	322/324 (99%)	307 (95%)	14 (4%)	1 (0%)	36	67
All	All	644/648 (99%)	614 (95%)	28 (4%)	2 (0%)	36	67

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	394	GLY
1	A	394	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	241/275 (88%)	231 (96%)	10 (4%)	27	60
1	B	275/275 (100%)	260 (94%)	15 (6%)	19	51
All	All	516/550 (94%)	491 (95%)	25 (5%)	23	55

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	283	THR
1	A	296	GLN
1	A	297	ASP
1	A	381	LYS
1	A	404	GLN
1	A	416	GLU
1	A	421	GLU
1	A	437	ASP
1	A	475	LEU
1	A	499	THR
1	B	216	ASP
1	B	252	GLN
1	B	266	LYS
1	B	296	GLN
1	B	330	SER
1	B	331	LYS
1	B	381	LYS
1	B	393	HIS
1	B	404	GLN
1	B	416	GLU
1	B	421	GLU
1	B	437	ASP
1	B	499	THR
1	B	524	ASP
1	B	531	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	254	ASN
1	A	272	ASN
1	A	392	ASN
1	A	393	HIS
1	A	464	HIS
1	B	272	ASN
1	B	275	GLN
1	B	392	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	324/324 (100%)	-0.43	0 100 100	27, 46, 76, 108	0
1	B	324/324 (100%)	-0.36	2 (0%) 85 72	26, 49, 81, 130	0
All	All	648/648 (100%)	-0.40	2 (0%) 90 82	26, 47, 79, 130	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	330	SER	2.0
1	B	415	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ZN	B	801	1/1	0.96	0.04	62,62,62,62	0
2	ZN	A	801	1/1	0.98	0.04	60,60,60,60	0

6.5 Other polymers [i](#)

There are no such residues in this entry.