



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 24, 2026 – 09:15 PM EDT

PDB ID : 1A8B / pdb_00001a8b
Title : RAT ANNEXIN V COMPLEXED WITH GLYCEROPHOSPHOETHANOL AMINE
Authors : Swairjo, M.A.; Concha, N.O.; Kaetzel, M.A.; Dedman, J.R.; Seaton, B.A.
Deposited on : 1998-03-23
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	319	5% 59% 34% 6%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2681 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ANNEXIN V.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	318	Total	C	N	O	S	0	0	0
			2503	1572	425	498	8			

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	10	Total	Ca	0	0
			10	10		

- Molecule 3 is L-ALPHA-GLYCEROPHOSPHORYLETHANOLAMINE (CCD ID: GPE) (formula: C₅H₁₄NO₆P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			13	5	1	6	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	155	Total 155	O 155	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ANNEXIN V



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	157.47Å 157.47Å 37.44Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	10.00 – 1.90 10.00 – 1.90	Depositor EDS
% Data completeness (in resolution range)	67.4 (10.00-1.90) 69.0 (10.00-1.90)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.57 (at 1.89Å)	Xtriage
Refinement program	X-PLOR	Depositor
R, R_{free}	0.200 , (Not available) 0.219 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	23.3	Xtriage
Anisotropy	0.422	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 69.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.018 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2681	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: CA, GPE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.27	5/2535 (0.2%)	2.02	76/3413 (2.2%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	185	TRP	CG-CD2	-6.35	1.32	1.43
1	A	265	HIS	CD2-NE2	-5.83	1.31	1.37
1	A	96	HIS	CD2-NE2	-5.80	1.31	1.37
1	A	314	CYS	CA-CB	-5.56	1.44	1.53
1	A	44	SER	CA-CB	-5.09	1.45	1.53

All (76) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	218	GLN	OE1-CD-NE2	-8.82	113.78	122.60
1	A	160	ASP	CA-CB-CG	8.54	121.14	112.60
1	A	104	ASP	CA-CB-CG	8.29	120.89	112.60
1	A	203	HIS	CA-C-O	-8.18	112.27	120.70
1	A	318	ASP	CA-CB-CG	7.68	120.28	112.60
1	A	212	MET	N-CA-C	-7.59	102.94	111.14
1	A	274	ARG	NE-CZ-NH2	-7.53	112.42	119.20
1	A	128	GLU	CB-CG-CD	6.92	124.37	112.60
1	A	9	ASP	CA-CB-CG	6.91	119.51	112.60
1	A	227	THR	N-CA-C	6.87	120.32	109.81
1	A	74	LYS	CA-CB-CG	6.86	127.83	114.10
1	A	263	ASP	CA-CB-CG	6.86	119.46	112.60
1	A	274	ARG	NE-CZ-NH1	6.84	128.34	121.50
1	A	259	GLY	O-C-N	6.64	128.97	122.92
1	A	240	LYS	CA-CB-CG	-6.42	101.27	114.10
1	A	116	THR	CA-C-N	6.38	125.60	118.97

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	116	THR	C-N-CA	6.38	125.60	118.97
1	A	13	PHE	CA-CB-CG	6.38	120.18	113.80
1	A	305	ASP	CA-CB-CG	6.30	118.90	112.60
1	A	201	VAL	CA-C-O	-6.25	114.11	121.05
1	A	143	THR	CA-CB-OG1	-6.21	100.29	109.60
1	A	47	GLN	CA-CB-CG	-6.18	101.74	114.10
1	A	203	HIS	CA-CB-CG	6.16	119.96	113.80
1	A	261	GLY	O-C-N	6.14	130.68	122.70
1	A	46	ALA	N-CA-C	-6.01	104.64	111.07
1	A	210	LYS	O-C-N	-5.98	115.78	122.12
1	A	175	GLN	O-C-N	-5.97	115.79	122.12
1	A	278	ASP	CA-CB-CG	5.91	118.50	112.60
1	A	137	ASP	CA-CB-CG	5.78	118.38	112.60
1	A	72	THR	N-CA-CB	-5.76	102.47	111.56
1	A	281	ASN	CB-CG-ND2	5.73	125.00	116.40
1	A	233	ASN	CB-CA-C	-5.73	101.64	110.81
1	A	76	GLU	N-CA-C	-5.71	104.96	111.07
1	A	218	GLN	N-CA-C	-5.70	101.26	110.32
1	A	15	GLY	O-C-N	5.70	127.66	122.19
1	A	314	CYS	N-CA-C	5.63	117.09	111.07
1	A	205	ARG	N-CA-C	-5.62	105.31	111.82
1	A	265	HIS	CA-CB-CG	5.59	119.39	113.80
1	A	162	ASP	N-CA-C	-5.57	101.41	109.59
1	A	233	ASN	N-CA-C	5.55	117.77	111.11
1	A	65	ASN	CB-CG-ND2	5.50	124.65	116.40
1	A	56	LYS	N-CA-C	-5.44	104.88	112.45
1	A	197	GLY	O-C-N	-5.44	116.35	122.28
1	A	53	GLU	CA-C-N	5.44	127.83	120.38
1	A	53	GLU	C-N-CA	5.44	127.83	120.38
1	A	209	ASP	CA-CB-CG	5.40	118.00	112.60
1	A	128	GLU	N-CA-C	-5.39	105.05	111.03
1	A	96	HIS	CB-CG-CD2	-5.39	124.20	131.20
1	A	317	GLU	CA-C-N	5.38	131.82	121.54
1	A	317	GLU	C-N-CA	5.38	131.82	121.54
1	A	230	ASN	O-C-N	-5.38	115.22	122.43
1	A	230	ASN	OD1-CG-ND2	-5.37	117.23	122.60
1	A	142	ASP	N-CA-CB	-5.35	102.66	110.47
1	A	203	HIS	O-C-N	-5.34	116.33	122.09
1	A	37	LEU	CA-CB-CG	5.30	134.84	116.30
1	A	273	SER	N-CA-C	5.28	117.78	111.71
1	A	37	LEU	N-CA-CB	-5.25	102.19	110.06
1	A	212	MET	CA-CB-CG	5.23	124.55	114.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	220	GLU	CB-CG-CD	5.21	121.45	112.60
1	A	8	THR	OG1-CB-CG2	5.18	119.67	109.30
1	A	31	THR	N-CA-C	-5.17	99.78	110.80
1	A	214	ILE	CB-CG1-CD1	-5.16	102.95	113.80
1	A	200	SER	CA-CB-OG	5.15	121.39	111.10
1	A	60	GLY	CA-C-O	5.14	126.27	119.58
1	A	193	ILE	N-CA-CB	-5.13	105.23	110.62
1	A	289	ASN	OD1-CG-ND2	-5.12	117.48	122.60
1	A	107	VAL	N-CA-C	-5.12	105.72	110.53
1	A	18	ASP	CA-CB-CG	5.11	117.71	112.60
1	A	34	ASP	N-CA-C	5.11	116.93	111.36
1	A	318	ASP	N-CA-C	-5.10	99.95	110.80
1	A	196	LEU	CA-C-N	5.09	126.68	120.22
1	A	196	LEU	C-N-CA	5.09	126.68	120.22
1	A	84	LYS	O-C-N	-5.07	117.23	121.38
1	A	185	TRP	CE2-CD2-CG	-5.07	101.12	107.20
1	A	220	GLU	CA-CB-CG	5.04	124.19	114.10
1	A	259	GLY	N-CA-C	5.00	121.57	112.27

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2503	0	2503	37	0
2	A	10	0	0	0	0
3	A	13	0	13	0	0
4	A	155	0	0	3	0
All	All	2681	0	2516	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:314:CYS:HB3	4:A:716:HOH:O	1.73	0.89
1:A:147:TYR:CD1	1:A:234:LEU:HD12	2.29	0.67
1:A:283:ARG:HG3	1:A:294:LEU:HD23	1.86	0.57
1:A:124:LYS:HG2	1:A:135:LEU:HD23	1.88	0.55
1:A:307:LYS:O	1:A:311:LEU:HG	2.08	0.52
1:A:78:LEU:HD22	1:A:310:LEU:HD13	1.91	0.51
1:A:150:MET:HE3	1:A:238:VAL:HG21	1.92	0.51
1:A:196:LEU:HD23	1:A:235:LEU:HD22	1.94	0.49
1:A:16:ARG:NH1	1:A:58:LEU:HD21	2.27	0.48
1:A:254:TYR:HE2	1:A:258:LYS:HD2	1.78	0.48
1:A:166:ASP:OD2	1:A:169:GLN:HG2	2.14	0.47
1:A:192:PHE:O	1:A:196:LEU:HB2	2.15	0.46
1:A:209:ASP:O	1:A:212:MET:HB3	2.15	0.46
1:A:68:LYS:HB2	1:A:68:LYS:NZ	2.31	0.45
1:A:126:ALA:HA	1:A:129:GLU:HG2	1.99	0.45
1:A:147:TYR:HD1	1:A:234:LEU:HD12	1.78	0.45
1:A:266:THR:O	1:A:270:VAL:HG23	2.17	0.45
1:A:298:ILE:CD1	1:A:314:CYS:SG	3.04	0.45
1:A:67:MET:HE1	1:A:83:MET:HG3	1.97	0.45
1:A:280:PHE:HD1	1:A:283:ARG:NH1	2.16	0.44
1:A:235:LEU:O	1:A:239:VAL:HG23	2.18	0.44
1:A:212:MET:HA	1:A:217:PHE:O	2.18	0.44
1:A:218:GLN:NE2	4:A:633:HOH:O	2.51	0.43
1:A:295:TYR:HB2	4:A:607:HOH:O	2.17	0.43
1:A:254:TYR:CE2	1:A:258:LYS:HD2	2.54	0.43
1:A:298:ILE:HD13	1:A:314:CYS:SG	2.60	0.42
1:A:71:LEU:HD22	1:A:75:PHE:CE2	2.55	0.42
1:A:115:ARG:HG3	1:A:119:GLU:OE1	2.19	0.41
1:A:243:ARG:HH11	1:A:243:ARG:HG2	1.84	0.41
1:A:267:LEU:O	1:A:271:ILE:HG13	2.20	0.41
1:A:310:LEU:HD12	1:A:310:LEU:HA	1.96	0.41
1:A:152:VAL:O	1:A:156:GLN:HG2	2.20	0.41
1:A:257:MET:HE3	1:A:257:MET:HB2	1.93	0.41
1:A:99:LYS:HD3	1:A:99:LYS:HA	1.86	0.41
1:A:304:GLY:O	1:A:308:LYS:HD2	2.21	0.40
1:A:302:THR:O	1:A:307:LYS:HD2	2.22	0.40
1:A:115:ARG:NE	1:A:274:ARG:HH21	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	316/319 (99%)	292 (92%)	21 (7%)	3 (1%)	14 6

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	318	ASP
1	A	316	GLY
1	A	101	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	268/268 (100%)	229 (85%)	39 (15%)	3 1

All (39) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	THR
1	A	23	ARG
1	A	34	ASP
1	A	37	LEU
1	A	53	GLU
1	A	57	THR
1	A	65	ASN
1	A	67	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	68	LYS
1	A	69	SER
1	A	72	THR
1	A	74	LYS
1	A	82	LEU
1	A	87	ARG
1	A	120	LEU
1	A	125	GLN
1	A	128	GLU
1	A	149	ARG
1	A	154	LEU
1	A	163	THR
1	A	165	ILE
1	A	172	LEU
1	A	187	THR
1	A	196	LEU
1	A	234	LEU
1	A	244	SER
1	A	246	PRO
1	A	262	THR
1	A	267	LEU
1	A	271	ILE
1	A	288	LYS
1	A	292	THR
1	A	296	SER
1	A	297	MET
1	A	301	ASP
1	A	308	LYS
1	A	310	LEU
1	A	313	LEU
1	A	318	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	134	ASN
1	A	179	GLN
1	A	218	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 11 ligands modelled in this entry, 10 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GPE	A	999	2	12,12,12	1.18	0	13,15,15	1.71	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GPE	A	999	2	-	4/13/13/13	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	999	GPE	C1-C2-C3	-4.47	96.72	111.77

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	999	GPE	C11-O12-P-O11
3	A	999	GPE	C11-O12-P-O13
3	A	999	GPE	O12-C11-C12-N
3	A	999	GPE	C2-C1-O11-P

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	318/319 (99%)	0.28	15 (4%) 36 39	5, 26, 52, 68	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	100	GLY	3.3
1	A	318	ASP	3.3
1	A	259	GLY	3.1
1	A	316	GLY	3.1
1	A	96	HIS	2.9
1	A	260	ALA	2.7
1	A	261	GLY	2.6
1	A	292	THR	2.6
1	A	4	ARG	2.5
1	A	287	ARG	2.5
1	A	319	ASP	2.3
1	A	10	PHE	2.1
1	A	28	GLY	2.1
1	A	8	THR	2.0
1	A	254	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CA	A	403	1/1	0.76	0.16	22,22,22,22	1
2	CA	A	406	1/1	0.79	0.09	20,20,20,20	1
3	GPE	A	999	13/13	0.81	0.14	22,24,29,29	13
2	CA	A	404	1/1	0.82	0.18	16,16,16,16	1
2	CA	A	409	1/1	0.85	0.10	21,21,21,21	1
2	CA	A	405	1/1	0.87	0.15	25,25,25,25	1
2	CA	A	410	1/1	0.94	0.06	24,24,24,24	1
2	CA	A	408	1/1	0.95	0.03	20,20,20,20	1
2	CA	A	401	1/1	0.97	0.05	24,24,24,24	0
2	CA	A	402	1/1	0.98	0.08	23,23,23,23	0
2	CA	A	407	1/1	0.98	0.02	13,13,13,13	0

6.5 Other polymers [i](#)

There are no such residues in this entry.