



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 01:44 AM UTC

PDB ID : 2A8T / pdb_00002a8t
Title : 2.1 Angstrom Crystal Structure of the Complex Between the Nuclear U8
snoRNA Decapping Nudix Hydrolase X29, Manganese and m7G-PPP-A
Authors : Scarsdale, J.N.; Peculis, B.A.; Wright, H.T.
Deposited on : 2005-07-08
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	212	
1	B	212	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3133 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called U8 snoRNA-binding protein X29.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	192	Total	C	N	O	S	0	0	0
			1427	896	254	270	7			
1	B	191	Total	C	N	O	S	0	0	0
			1497	946	262	282	7			

- Molecule 2 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	4	Total	Mn	0	0
			4	4		
2	B	3	Total	Mn	0	2
			5	5		

- Molecule 3 is 7N-METHYL-8-HYDROGUANOSINE-5'-TRIPHOSPHATE (CCD ID: MGT) (formula: C₁₁H₂₀N₅O₁₄P₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			32	11	5	13		
3	B	1	Total	C	N	O	0	0
			31	11	5	12		

- Molecule 4 is ADENOSINE (CCD ID: ADN) (formula: $C_{10}H_{13}N_5O_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			19	10	5	4		
4	B	1	Total	C	N	O	0	0
			19	10	5	4		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	30	Total	O	0	0
			30	30		
5	B	69	Total	O	0	0
			69	69		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: U8 snoRNA-binding protein X29



• Molecule 1: U8 snoRNA-binding protein X29



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	50.70Å 81.71Å 111.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.17 – 2.10 46.17 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.9 (46.17-2.10) 99.9 (46.17-2.10)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.33 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.2, CNS	Depositor
R, R_{free}	0.212 , 0.264 (Not available) , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	48.3	Xtriage
Anisotropy	0.381	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 68.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3133	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MGT, ADN, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.69	2/1448 (0.1%)	0.86	2/1963 (0.1%)
1	B	0.50	0/1521	0.78	0/2052
All	All	0.60	2/2969 (0.1%)	0.82	2/4015 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	115	GLU	CD-OE1	13.92	1.51	1.25
1	A	115	GLU	CD-OE2	11.02	1.46	1.25

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	117	PRO	N-CA-CB	7.88	111.52	103.25
1	A	18	PRO	N-CA-CB	6.68	110.35	103.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1427	0	1335	15	0
1	B	1497	0	1488	15	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	4	0	0	0	0
2	B	5	0	0	0	0
3	A	32	0	16	1	0
3	B	31	0	16	0	0
4	A	19	0	12	3	0
4	B	19	0	12	1	0
5	A	30	0	0	1	0
5	B	69	0	0	0	0
All	All	3133	0	2879	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:61:MET:HE2	1:A:63:ARG:HG2	1.65	0.77
1:B:180:ASN:HD22	1:B:180:ASN:H	1.45	0.64
1:B:102:GLU:O	1:B:127:LYS:HE3	1.99	0.63
1:B:114:ARG:NH2	1:B:116:HIS:O	2.34	0.60
1:A:112:GLN:HE22	1:A:184:GLN:HE21	1.49	0.60
3:A:251:MGT:H4'	1:B:49:PHE:CE2	2.40	0.56
1:A:180:ASN:HD22	1:A:180:ASN:H	1.53	0.55
1:A:122:THR:HG21	4:A:252:ADN:C8	2.37	0.54
1:B:190:ARG:HA	1:B:190:ARG:HE	1.76	0.50
1:B:190:ARG:HG2	1:B:200:ILE:HD12	1.94	0.49
4:A:252:ADN:H2	5:A:639:HOH:O	2.11	0.49
1:A:109:ARG:HG2	1:A:126:ILE:HD12	1.95	0.48
1:A:184:GLN:OE1	4:A:252:ADN:N6	2.49	0.46
1:B:73:GLY:N	1:B:89:GLU:HG2	2.31	0.45
1:A:76:ASP:HB3	1:A:79:ASP:HB2	1.98	0.45
1:A:109:ARG:HD3	1:A:126:ILE:HB	1.99	0.45
1:A:180:ASN:HD22	1:A:180:ASN:N	2.15	0.44
1:B:72:GLY:C	1:B:89:GLU:HG2	2.42	0.44
1:A:26:SER:HB2	1:A:121:VAL:HG11	1.99	0.44
1:A:63:ARG:HB2	1:A:65:ASP:OD1	2.18	0.44
1:B:41:HIS:HA	1:B:57:VAL:O	2.18	0.43
1:B:70:PHE:HD1	4:B:252:ADN:HN62	1.65	0.43
1:A:61:MET:HE2	1:A:63:ARG:CG	2.42	0.42
1:B:22:SER:OG	1:B:25:GLU:HB2	2.19	0.42
1:B:20:ASN:ND2	1:B:112:GLN:OE1	2.52	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:62:MET:HE3	1:B:155:ILE:HD13	2.03	0.41
1:A:112:GLN:NE2	1:A:184:GLN:HE21	2.16	0.41
1:B:103:VAL:HG22	1:B:127:LYS:HD2	2.03	0.41
1:A:41:HIS:ND1	1:A:56:ARG:HD3	2.36	0.41
1:B:149:LEU:O	1:B:152:MET:HE2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	190/212 (90%)	177 (93%)	9 (5%)	4 (2%)	5	2
1	B	189/212 (89%)	182 (96%)	7 (4%)	0	100	100
All	All	379/424 (89%)	359 (95%)	16 (4%)	4 (1%)	11	8

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	117	PRO
1	A	79	ASP
1	A	102	GLU
1	A	103	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	138/185 (75%)	132 (96%)	6 (4%)	26	27
1	B	159/185 (86%)	152 (96%)	7 (4%)	25	26
All	All	297/370 (80%)	284 (96%)	13 (4%)	25	26

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	88	ARG
1	A	109	ARG
1	A	114	ARG
1	A	180	ASN
1	A	183	SER
1	A	201	GLN
1	B	47	LYS
1	B	102	GLU
1	B	132	GLU
1	B	134	ILE
1	B	180	ASN
1	B	190	ARG
1	B	203	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	112	GLN
1	A	180	ASN
1	B	20	ASN
1	B	112	GLN
1	B	180	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 9 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	ADN	A	252	3	21,21,21	1.21	4 (19%)	31,31,31	1.94	7 (22%)
3	MGT	A	251	4,2	31,34,35	3.68	5 (16%)	39,53,56	1.95	7 (17%)
3	MGT	B	251	4,2	29,33,35	4.05	7 (24%)	34,51,56	1.83	6 (17%)
4	ADN	B	252	3	21,21,21	1.11	2 (9%)	31,31,31	2.05	9 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ADN	A	252	3	-	0/6/22/22	0/3/3/3
3	MGT	A	251	4,2	-	8/19/49/50	0/3/3/3
3	MGT	B	251	4,2	-	7/19/49/50	0/3/3/3
4	ADN	B	252	3	-	1/6/22/22	0/3/3/3

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	251	MGT	C8-N9	-18.18	1.34	1.45
3	B	251	MGT	C8-N9	-18.05	1.34	1.45
3	B	251	MGT	C6-C5	-7.30	1.40	1.49
3	B	251	MGT	C2-N1	6.93	1.41	1.34
3	A	251	MGT	PA-O3A	5.82	1.65	1.59
3	A	251	MGT	PB-O3A	4.09	1.63	1.59

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	252	ADN	C5-N7	-3.34	1.33	1.39
3	B	251	MGT	C4-N9	-3.28	1.33	1.37
4	B	252	ADN	C5-N7	-3.17	1.33	1.39
3	B	251	MGT	C6-N1	-3.16	1.40	1.45
3	A	251	MGT	C4-N9	-3.01	1.34	1.37
3	B	251	MGT	PA-O3A	2.71	1.62	1.59
4	A	252	ADN	O4'-C1'	2.29	1.47	1.42
3	A	251	MGT	C1'-N9	2.25	1.50	1.46
3	B	251	MGT	PB-O3A	2.17	1.61	1.59
4	A	252	ADN	C8-N9	-2.10	1.34	1.37
4	B	252	ADN	C8-N9	-2.02	1.34	1.37
4	A	252	ADN	C4-N9	-2.01	1.33	1.37

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	252	ADN	C5-C4-N3	-5.51	119.12	126.72
4	A	252	ADN	C5-C4-N3	-5.48	119.17	126.72
3	A	251	MGT	C2-N3-C4	5.47	121.72	112.30
3	B	251	MGT	C2-N3-C4	5.24	121.33	112.30
3	A	251	MGT	C5-C6-N1	4.60	119.03	110.94
4	B	252	ADN	N3-C2-N1	-4.47	121.82	128.58
3	A	251	MGT	N9-C4-N3	4.35	131.83	125.46
4	A	252	ADN	N3-C2-N1	-4.32	122.04	128.58
3	B	251	MGT	N9-C4-N3	4.08	131.44	125.46
3	A	251	MGT	C5-C4-N3	-4.04	120.54	128.13
4	B	252	ADN	N3-C4-N9	3.99	133.95	127.17
4	A	252	ADN	N3-C4-N9	3.90	133.80	127.17
3	A	251	MGT	N9-C8-N7	3.62	108.50	103.37
4	A	252	ADN	C2-N3-C4	3.61	120.64	111.83
4	B	252	ADN	C2-N3-C4	3.56	120.53	111.83
3	B	251	MGT	N9-C8-N7	3.56	108.41	103.37
3	B	251	MGT	C5-C4-N3	-3.35	120.90	127.16
3	B	251	MGT	N1-C2-N3	-3.35	120.30	122.97
3	A	251	MGT	C2-N1-C6	-3.29	119.14	125.11
4	A	252	ADN	N9-C8-N7	-2.92	109.79	113.94
4	B	252	ADN	N9-C8-N7	-2.75	110.04	113.94
4	A	252	ADN	C5-N7-C8	2.62	107.56	103.45
3	B	251	MGT	N2-C2-N1	2.59	120.02	117.93
4	B	252	ADN	C5'-C4'-C3'	-2.51	109.17	115.10
3	A	251	MGT	O6-C6-C5	-2.49	121.51	127.62
4	B	252	ADN	C5-N7-C8	2.48	107.35	103.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	252	ADN	C4-C5-N7	-2.38	107.86	110.58
4	B	252	ADN	C4-C5-N7	-2.30	107.95	110.58
4	B	252	ADN	C4'-O4'-C1'	-2.10	104.83	109.47

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	251	MGT	C5'-O5'-PA-O3A
3	A	251	MGT	C5'-O5'-PA-O1A
3	B	251	MGT	C5'-O5'-PA-O3A
3	B	251	MGT	C5'-O5'-PA-O1A
3	B	251	MGT	C5'-O5'-PA-O2A
3	A	251	MGT	PG-O3B-PB-O1B
3	B	251	MGT	PG-O3B-PB-O1B
3	A	251	MGT	C5'-O5'-PA-O2A
3	A	251	MGT	PG-O3B-PB-O3A
3	B	251	MGT	PG-O3B-PB-O2B
3	B	251	MGT	PA-O3A-PB-O1B
3	B	251	MGT	PG-O3B-PB-O3A
3	A	251	MGT	PG-O3B-PB-O2B
3	A	251	MGT	PA-O3A-PB-O3B
3	A	251	MGT	PA-O3A-PB-O1B
4	B	252	ADN	O4'-C4'-C5'-O5'



There are no ring outliers.


3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	252	ADN	3	0
3	A	251	MGT	1	0
4	B	252	ADN	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	192/212 (90%)	2.03	94 (48%) 0 0	36, 57, 71, 82	1 (0%)
1	B	191/212 (90%)	0.85	8 (4%) 40 42	49, 57, 71, 78	0
All	All	383/424 (90%)	1.45	102 (26%) 1 1	36, 57, 71, 82	1 (0%)

All (102) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	76	ASP	5.8
1	A	32	TYR	5.2
1	A	48	LEU	4.7
1	A	117	PRO	4.3
1	A	163	ARG	4.3
1	A	113	VAL	4.1
1	A	78	ARG	4.0
1	A	103	VAL	3.9
1	B	208	HIS	3.8
1	A	171	ALA	3.7
1	A	82	LEU	3.6
1	A	52	VAL	3.6
1	A	161	THR	3.6
1	A	19	ARG	3.6
1	A	49	PHE	3.6
1	A	169	LEU	3.5
1	A	154	LEU	3.5
1	A	79	ASP	3.4
1	A	75	VAL	3.4
1	A	157	VAL	3.4
1	A	66	GLY	3.4
1	A	139	ALA	3.3
1	A	166	VAL	3.3
1	A	77	THR	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	167	GLY	3.2
1	A	100	THR	3.2
1	A	29	LEU	3.2
1	B	18	PRO	3.1
1	A	168	GLY	3.1
1	A	121	VAL	3.1
1	A	94	LEU	3.1
1	A	44	SER	3.0
1	A	64	PHE	3.0
1	A	159	LEU	2.9
1	A	153	GLY	2.9
1	A	156	ARG	2.9
1	A	99	ALA	2.9
1	A	208	HIS	2.9
1	A	28	GLN	2.8
1	A	45	GLN	2.8
1	A	144	ALA	2.8
1	A	59	LEU	2.8
1	A	26	SER	2.8
1	A	132	GLU	2.8
1	A	22	SER	2.8
1	A	195	LEU	2.8
1	A	101	VAL	2.7
1	A	21	ILE	2.7
1	A	164	ASP	2.7
1	A	172	PHE	2.7
1	A	54	ILE	2.7
1	A	80	ILE	2.7
1	A	170	PRO	2.6
1	A	36	CYS	2.6
1	A	155	ILE	2.6
1	A	25	GLU	2.6
1	A	74	PHE	2.6
1	A	200	ILE	2.6
1	A	116	HIS	2.6
1	A	27	LEU	2.6
1	A	98	LEU	2.6
1	A	120	CYS	2.5
1	A	141	ALA	2.5
1	B	99	ALA	2.5
1	A	51	ARG	2.5
1	A	35	ALA	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	114	ARG	2.5
1	A	151	VAL	2.5
1	A	43	PRO	2.5
1	A	134	ILE	2.5
1	A	46	ALA	2.4
1	A	34	HIS	2.4
1	A	174	CYS	2.4
1	A	204	LEU	2.4
1	A	42	ALA	2.4
1	A	148	GLY	2.4
1	A	142	VAL	2.4
1	A	20	ASN	2.3
1	A	162	LEU	2.3
1	B	117	PRO	2.3
1	A	192	LEU	2.3
1	A	24	GLU	2.3
1	A	207	SER	2.3
1	A	57	VAL	2.3
1	A	152	MET	2.2
1	B	97	ALA	2.2
1	A	131	LEU	2.2
1	A	108	TYR	2.2
1	A	125	TYR	2.2
1	A	104	THR	2.2
1	B	118	GLN	2.2
1	A	158	PRO	2.2
1	A	136	ARG	2.2
1	A	85	GLY	2.1
1	B	27	LEU	2.1
1	A	50	ASP	2.1
1	A	160	TYR	2.1
1	A	203	VAL	2.1
1	A	97	ALA	2.1
1	A	58	LEU	2.0
1	A	73	GLY	2.0
1	B	116	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MN	B	306[A]	1/1	0.53	0.30	65,65,65,65	1
2	MN	B	306[B]	1/1	0.53	0.30	85,85,85,85	1
4	ADN	A	252	19/19	0.65	0.23	61,85,90,91	19
3	MGT	B	251	31/33	0.73	0.33	47,84,91,92	31
4	ADN	B	252	19/19	0.76	0.33	55,64,69,71	19
2	MN	A	303	1/1	0.83	0.13	56,56,56,56	1
3	MGT	A	251	32/33	0.85	0.16	50,84,90,91	32
2	MN	A	302	1/1	0.88	0.08	74,74,74,74	0
2	MN	A	300	1/1	0.93	0.08	60,60,60,60	0
2	MN	A	301	1/1	0.93	0.06	62,62,62,62	0
2	MN	B	305[A]	1/1	0.95	0.07	64,64,64,64	1
2	MN	B	305[B]	1/1	0.95	0.07	45,45,45,45	1
2	MN	B	304	1/1	0.97	0.07	64,64,64,64	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ADN A 252:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MGT B 251:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ADN B 252:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around MGT A 251:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.