



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 25, 2026 – 04:49 AM EDT

PDB ID : 2A8Y / pdb_00002a8y
Title : Crystal structure of 5'-deoxy-5'methylthioadenosine phosphorylase complexed with 5'-deoxy-5'methylthioadenosine and sulfate
Authors : Zhang, Y.; Porcelli, M.; Cacciapuoti, G.; Ealick, S.E.
Deposited on : 2005-07-10
Resolution : 1.45 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1756 (1.46-1.46)
Clashscore	190562	1795 (1.46-1.46)
Ramachandran outliers	187476	1776 (1.46-1.46)
Sidechain outliers	187428	1776 (1.46-1.46)
RSRZ outliers	180081	1756 (1.46-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	270	
1	B	270	
1	C	270	
1	D	270	
1	E	270	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	270	 78%18%..
1	G	270	 75%21%..
1	H	270	 72%23%..
1	I	270	 78%18%..
1	J	270	 73%23%..
1	K	270	 71%24%..
1	L	270	 74%23%..

2 Entry composition


There are 4 unique types of molecules in this entry. The entry contains 29330 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 5'-methylthioadenosine phosphorylase (mtaP).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	263	Total	C	N	O	S	0	16	0
			2113	1361	350	384	18			
1	B	263	Total	C	N	O	S	0	16	0
			2135	1370	354	393	18			
1	C	263	Total	C	N	O	S	0	16	0
			2136	1372	356	391	17			
1	D	263	Total	C	N	O	S	0	16	0
			2116	1363	350	385	18			
1	E	263	Total	C	N	O	S	0	16	0
			2138	1376	358	386	18			
1	F	263	Total	C	N	O	S	0	16	0
			2127	1370	352	388	17			
1	G	263	Total	C	N	O	S	0	16	0
			2130	1372	352	388	18			
1	H	263	Total	C	N	O	S	0	16	0
			2129	1370	355	386	18			
1	I	263	Total	C	N	O	S	0	16	0
			2146	1380	358	390	18			
1	J	263	Total	C	N	O	S	0	16	0
			2127	1369	356	385	17			
1	K	263	Total	C	N	O	S	0	16	0
			2136	1373	356	389	18			
1	L	263	Total	C	N	O	S	0	16	0
			2132	1369	355	391	17			

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		
2	E	1	Total	O	S	0	0
			5	4	1		
2	F	1	Total	O	S	0	0
			5	4	1		
2	G	1	Total	O	S	0	0
			5	4	1		
2	G	1	Total	O	S	0	0
			5	4	1		
2	H	1	Total	O	S	0	0
			5	4	1		
2	H	1	Total	O	S	0	0
			5	4	1		
2	H	1	Total	O	S	0	0
			5	4	1		
2	I	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	I	1	Total	O	S	0	0
			5	4	1		
2	I	1	Total	O	S	0	0
			5	4	1		
2	J	1	Total	O	S	0	0
			5	4	1		
2	J	1	Total	O	S	0	0
			5	4	1		
2	K	1	Total	O	S	0	0
			5	4	1		
2	K	1	Total	O	S	0	0
			5	4	1		
2	K	1	Total	O	S	0	0
			5	4	1		
2	L	1	Total	O	S	0	0
			5	4	1		
2	L	1	Total	O	S	0	0
			5	4	1		
2	L	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is 5'-DEOXY-5'-METHYLTHIOADENOSINE (CCD ID: MTA) (formula: $C_{11}H_{15}N_5O_3S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			20	11	5	3	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	N	O	S	0	0
			20	11	5	3	1		
3	C	1	Total	C	N	O	S	0	0
			20	11	5	3	1		
3	D	1	Total	C	N	O	S	0	0
			20	11	5	3	1		
3	E	1	Total	C	N	O	S	0	0
			20	11	5	3	1		
3	F	1	Total	C	N	O	S	0	0
			20	11	5	3	1		
3	G	1	Total	C	N	O	S	0	0
			20	11	5	3	1		
3	H	1	Total	C	N	O	S	0	0
			20	11	5	3	1		
3	I	1	Total	C	N	O	S	0	0
			20	11	5	3	1		
3	J	1	Total	C	N	O	S	0	0
			20	11	5	3	1		
3	K	1	Total	C	N	O	S	0	0
			20	11	5	3	1		
3	L	1	Total	C	N	O	S	0	0
			20	11	5	3	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	266	Total	O	0	0
			266	266		
4	B	311	Total	O	0	0
			311	311		
4	C	265	Total	O	0	0
			265	265		
4	D	293	Total	O	0	0
			293	293		
4	E	281	Total	O	0	0
			281	281		
4	F	275	Total	O	0	0
			275	275		
4	G	273	Total	O	0	0
			273	273		
4	H	283	Total	O	0	0
			283	283		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	I	288	Total 288	O 288	0	0
4	J	261	Total 261	O 261	0	0
4	K	293	Total 293	O 293	0	0
4	L	316	Total 316	O 316	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 5'-methylthioadenosine phosphorylase (mtaP)




- Molecule 1: 5'-methylthioadenosine phosphorylase (mtaP)



- Molecule 1: 5'-methylthioadenosine phosphorylase (mtaP)




- Molecule 1: 5'-methylthioadenosine phosphorylase (mtaP)






- Molecule 1: 5'-methylthioadenosine phosphorylase (mtaP)

Chain E: 74% 19%




- Molecule 1: 5'-methylthioadenosine phosphorylase (mtaP)

Chain F: 78% 18%




- Molecule 1: 5'-methylthioadenosine phosphorylase (mtaP)

Chain G: 75% 21%




- Molecule 1: 5'-methylthioadenosine phosphorylase (mtaP)


Chain H: 72% 23%




- Molecule 1: 5'-methylthioadenosine phosphorylase (mtaP)




- Molecule 1: 5'-methylthioadenosine phosphorylase (mtaP)



- Molecule 1: 5'-methylthioadenosine phosphorylase (mtaP)



- Molecule 1: 5'-methylthioadenosine phosphorylase (mtaP)



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	96.60Å 96.56Å 96.63Å 91.57° 91.23° 91.52°	Depositor
Resolution (Å)	48.27 – 1.45 48.27 – 1.45	Depositor EDS
% Data completeness (in resolution range)	89.0 (48.27-1.45) 89.1 (48.27-1.45)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.41 (at 1.45Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.182 , 0.202 0.179 , 0.199	Depositor DCC
R_{free} test set	40550 reflections (6.88%)	wwPDB-VP
Wilson B-factor (Å ²)	11.9	Xtriage
Anisotropy	0.126	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 39.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.53$, $\langle L^2 \rangle = 0.36$	Xtriage

Continued on next page...

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

Continued from previous page...

Property	Value	Source
Estimated twinning fraction	0.000 for l,k,-h 0.000 for -l,k,h 0.000 for h,-l,k 0.000 for h,l,-k 0.000 for -k,h,l 0.000 for k,-h,l 0.034 for l,h,k 0.034 for k,l,h 0.000 for -k,l,-h 0.000 for -l,-h,k 0.000 for -k,-l,h 0.000 for l,-h,-k 0.000 for -l,h,-k 0.000 for k,-l,-h 0.011 for -h,k,-l 0.000 for h,-k,-l 0.000 for -h,-k,l 0.000 for k,h,-l 0.033 for -k,-h,-l 0.000 for -h,l,k 0.036 for -h,-l,-k 0.011 for l,-k,h 0.467 for -l,-k,-h	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	29330	wwPDB-VP
Average B, all atoms ($\text{\AA}^2$)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 45.63 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.2853e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, MTA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.75	1/2158 (0.0%)	1.04	9/2929 (0.3%)
1	B	0.75	1/2180 (0.0%)	1.05	10/2958 (0.3%)
1	C	0.72	0/2181	1.05	10/2958 (0.3%)
1	D	0.75	0/2161	1.08	12/2933 (0.4%)
1	E	0.76	1/2183 (0.0%)	1.08	15/2959 (0.5%)
1	F	0.71	0/2172	1.06	13/2947 (0.4%)
1	G	0.71	0/2175	1.05	13/2950 (0.4%)
1	H	0.72	0/2174	1.06	14/2948 (0.5%)
1	I	0.74	1/2191 (0.0%)	1.04	8/2969 (0.3%)
1	J	0.70	0/2172	1.07	10/2947 (0.3%)
1	K	0.73	1/2181 (0.0%)	1.04	11/2957 (0.4%)
1	L	0.77	0/2177	1.05	9/2954 (0.3%)
All	All	0.73	5/26105 (0.0%)	1.06	134/35409 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	E	0	1
1	H	0	1
1	I	0	1
1	J	0	1
1	K	0	1
All	All	0	6

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	114	MET	SD-CE	-7.47	1.60	1.79
1	K	114	MET	SD-CE	-7.26	1.61	1.79
1	E	114	MET	SD-CE	-6.91	1.62	1.79
1	I	114	MET	SD-CE	-6.45	1.63	1.79
1	B	39	PRO	CA-C	5.10	1.58	1.52

The worst 5 of 134 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	117	ASN	N-CA-C	11.64	127.27	111.90
1	K	38	GLN	N-CA-C	10.47	124.91	109.42
1	F	38	GLN	N-CA-C	9.88	124.05	109.42
1	J	117	ASN	N-CA-C	9.68	124.68	111.90
1	G	38	GLN	N-CA-C	9.64	123.69	109.42

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	162	TYR	Sidechain
1	E	33	TYR	Sidechain
1	H	162	TYR	Sidechain
1	I	33	TYR	Sidechain
1	J	33	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2113	0	2080	82	0
1	B	2135	0	2099	72	0
1	C	2136	0	2110	57	0
1	D	2116	0	2085	64	0
1	E	2138	0	2131	71	0
1	F	2127	0	2106	48	0
1	G	2130	0	2109	62	0
1	H	2129	0	2107	65	0
1	I	2146	0	2139	47	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	J	2127	0	2107	65	0
1	K	2136	0	2115	61	0
1	L	2132	0	2099	72	0
2	A	10	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
2	D	10	0	0	0	0
2	E	5	0	0	0	0
2	F	5	0	0	0	0
2	G	10	0	0	0	0
2	H	15	0	0	0	0
2	I	15	0	0	0	0
2	J	10	0	0	0	0
2	K	15	0	0	0	0
2	L	15	0	0	0	0
3	A	20	0	15	1	0
3	B	20	0	15	2	0
3	C	20	0	15	2	0
3	D	20	0	15	2	0
3	E	20	0	15	1	0
3	F	20	0	15	1	0
3	G	20	0	15	1	0
3	H	20	0	15	1	0
3	I	20	0	15	2	0
3	J	20	0	15	2	0
3	K	20	0	15	1	0
3	L	20	0	15	0	0
4	A	266	0	0	13	0
4	B	311	0	0	19	0
4	C	265	0	0	8	0
4	D	293	0	0	13	0
4	E	281	0	0	22	0
4	F	275	0	0	9	0
4	G	273	0	0	6	0
4	H	283	0	0	10	0
4	I	288	0	0	16	0
4	J	261	0	0	12	0
4	K	293	0	0	13	0
4	L	316	0	0	11	0
All	All	29330	0	25467	754	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 754 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:202:ALA:HB3	1:L:204[A]:MET:HE2	1.25	1.13
1:L:48:ILE:HG22	1:L:248[B]:GLN:HE22	1.02	1.13
1:G:48:ILE:HG22	1:G:248[B]:GLN:HE22	1.15	1.11
1:D:48:ILE:HG22	1:D:248[B]:GLN:HE22	1.15	1.10
1:D:202:ALA:HB3	1:D:204[B]:MET:HE2	1.29	1.09

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	275/270 (102%)	270 (98%)	5 (2%)	0	100	100
1	B	275/270 (102%)	271 (98%)	4 (2%)	0	100	100
1	C	275/270 (102%)	269 (98%)	6 (2%)	0	100	100
1	D	275/270 (102%)	269 (98%)	5 (2%)	1 (0%)	30	11
1	E	275/270 (102%)	268 (98%)	6 (2%)	1 (0%)	30	11
1	F	275/270 (102%)	270 (98%)	5 (2%)	0	100	100
1	G	275/270 (102%)	269 (98%)	6 (2%)	0	100	100
1	H	275/270 (102%)	267 (97%)	7 (2%)	1 (0%)	30	11
1	I	275/270 (102%)	270 (98%)	5 (2%)	0	100	100
1	J	275/270 (102%)	268 (98%)	6 (2%)	1 (0%)	30	11
1	K	275/270 (102%)	268 (98%)	6 (2%)	1 (0%)	30	11
1	L	275/270 (102%)	270 (98%)	5 (2%)	0	100	100
All	All	3300/3240 (102%)	3229 (98%)	66 (2%)	5 (0%)	43	21

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	263	ASN
1	H	263	ASN
1	J	263	ASN
1	K	263	ASN
1	E	48	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	221/231 (96%)	217 (98%)	4 (2%)	51	19
1	B	226/231 (98%)	222 (98%)	4 (2%)	51	19
1	C	226/231 (98%)	222 (98%)	4 (2%)	51	19
1	D	222/231 (96%)	218 (98%)	4 (2%)	51	19
1	E	227/231 (98%)	219 (96%)	8 (4%)	32	5
1	F	225/231 (97%)	221 (98%)	4 (2%)	51	19
1	G	225/231 (97%)	222 (99%)	3 (1%)	61	31
1	H	224/231 (97%)	219 (98%)	5 (2%)	45	13
1	I	229/231 (99%)	223 (97%)	6 (3%)	40	10
1	J	224/231 (97%)	218 (97%)	6 (3%)	39	9
1	K	226/231 (98%)	221 (98%)	5 (2%)	45	13
1	L	225/231 (97%)	223 (99%)	2 (1%)	70	46
All	All	2700/2772 (97%)	2645 (98%)	55 (2%)	51	16

5 of 55 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	34	THR
1	H	221	ILE
1	L	221	ILE
1	K	102	LEU
1	G	211[A]	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 49 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	235	ASN
1	I	153	ASN
1	H	75	ASN
1	H	157	HIS
1	J	4	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

36 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	MTA	J	4030	-	22,22,22	1.30	3 (13%)	32,32,32	1.61	6 (18%)
2	SO4	K	4032	-	4,4,4	0.47	0	6,6,6	0.07	0
2	SO4	K	4008	-	4,4,4	0.32	0	6,6,6	0.31	0
2	SO4	I	4025	-	4,4,4	0.16	0	6,6,6	0.73	0
2	SO4	L	4035	-	4,4,4	0.50	0	6,6,6	0.22	0
3	MTA	F	4018	-	22,22,22	1.36	3 (13%)	32,32,32	1.67	6 (18%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	MTA	A	4003	-	22,22,22	1.19	1 (4%)	32,32,32	1.71	6 (18%)
2	SO4	I	4014	-	4,4,4	0.31	0	6,6,6	0.26	0
3	MTA	K	4033	-	22,22,22	1.33	4 (18%)	32,32,32	1.72	5 (15%)
2	SO4	D	4010	-	4,4,4	0.47	0	6,6,6	0.63	0
3	MTA	L	4036	-	22,22,22	1.23	1 (4%)	32,32,32	1.70	7 (21%)
2	SO4	J	4028	-	4,4,4	0.37	0	6,6,6	0.42	0
2	SO4	L	4034	-	4,4,4	0.35	0	6,6,6	0.56	0
2	SO4	E	4013	-	4,4,4	0.12	0	6,6,6	0.71	0
2	SO4	F	4016	-	4,4,4	0.34	0	6,6,6	0.67	0
3	MTA	B	4006	-	22,22,22	1.33	2 (9%)	32,32,32	1.71	6 (18%)
2	SO4	J	4029	-	4,4,4	0.38	0	6,6,6	0.09	0
2	SO4	K	4031	-	4,4,4	0.29	0	6,6,6	0.72	0
3	MTA	D	4012	-	22,22,22	1.22	1 (4%)	32,32,32	1.59	6 (18%)
3	MTA	H	4024	-	22,22,22	1.18	1 (4%)	32,32,32	1.68	7 (21%)
3	MTA	C	4009	-	22,22,22	1.39	3 (13%)	32,32,32	1.71	7 (21%)
2	SO4	G	4019	-	4,4,4	0.16	0	6,6,6	0.66	0
2	SO4	C	4007	-	4,4,4	0.41	0	6,6,6	0.65	0
2	SO4	B	4004	-	4,4,4	0.30	0	6,6,6	0.55	0
3	MTA	I	4027	-	22,22,22	1.23	3 (13%)	32,32,32	1.81	6 (18%)
2	SO4	G	4020	-	4,4,4	0.43	0	6,6,6	0.17	0
3	MTA	G	4021	-	22,22,22	1.38	4 (18%)	32,32,32	1.61	6 (18%)
2	SO4	I	4026	-	4,4,4	0.47	0	6,6,6	0.29	0
3	MTA	E	4015	-	22,22,22	1.31	2 (9%)	32,32,32	1.80	6 (18%)
2	SO4	A	4000	-	4,4,4	0.30	0	6,6,6	0.47	0
2	SO4	H	4023	-	4,4,4	0.42	0	6,6,6	0.12	0
2	SO4	H	4017	-	4,4,4	0.39	0	6,6,6	0.19	0
2	SO4	A	4001	-	4,4,4	0.41	0	6,6,6	0.38	0
2	SO4	L	4005	-	4,4,4	0.28	0	6,6,6	0.27	0
2	SO4	H	4022	-	4,4,4	0.55	0	6,6,6	0.67	0
2	SO4	D	4011	-	4,4,4	0.37	0	6,6,6	0.16	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MTA	E	4015	-	-	2/7/23/23	0/3/3/3
3	MTA	C	4009	-	-	2/7/23/23	0/3/3/3
3	MTA	J	4030	-	-	2/7/23/23	0/3/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MTA	F	4018	-	-	2/7/23/23	0/3/3/3
3	MTA	I	4027	-	-	2/7/23/23	0/3/3/3
3	MTA	A	4003	-	-	2/7/23/23	0/3/3/3
3	MTA	B	4006	-	-	2/7/23/23	0/3/3/3
3	MTA	D	4012	-	-	2/7/23/23	0/3/3/3
3	MTA	K	4033	-	-	2/7/23/23	0/3/3/3
3	MTA	H	4024	-	-	2/7/23/23	0/3/3/3
3	MTA	G	4021	-	-	2/7/23/23	0/3/3/3
3	MTA	L	4036	-	-	2/7/23/23	0/3/3/3

The worst 5 of 28 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	4015	MTA	O4'-C1'	4.00	1.51	1.42
3	B	4006	MTA	O4'-C1'	3.77	1.50	1.42
3	K	4033	MTA	O4'-C1'	3.67	1.50	1.42
3	G	4021	MTA	O4'-C1'	3.58	1.50	1.42
3	L	4036	MTA	O4'-C1'	3.38	1.49	1.42

The worst 5 of 74 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	I	4027	MTA	O4'-C1'-N9	-6.29	96.01	108.09
3	E	4015	MTA	O4'-C1'-N9	-6.24	96.10	108.09
3	A	4003	MTA	O4'-C1'-N9	-6.06	96.46	108.09
3	K	4033	MTA	O4'-C1'-N9	-6.00	96.56	108.09
3	B	4006	MTA	O4'-C1'-N9	-5.66	97.22	108.09

There are no chirality outliers.

5 of 24 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	A	4003	MTA	C2'-C1'-N9-C4
3	B	4006	MTA	C2'-C1'-N9-C4
3	C	4009	MTA	C2'-C1'-N9-C4
3	D	4012	MTA	C2'-C1'-N9-C4
3	E	4015	MTA	C2'-C1'-N9-C8

There are no ring outliers.


11 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	J	4030	MTA	2	0
3	F	4018	MTA	1	0
3	A	4003	MTA	1	0
3	K	4033	MTA	1	0
3	B	4006	MTA	2	0
3	D	4012	MTA	2	0
3	H	4024	MTA	1	0
3	C	4009	MTA	2	0
3	I	4027	MTA	2	0
3	G	4021	MTA	1	0
3	E	4015	MTA	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand MTA A 4003




Ligand MTA K 4033




Ligand MTA L 4036




Ligand MTA B 4006




Ligand MTA D 4012




Ligand MTA H 4024




Ligand MTA C 4009



Ligand MTA I 4027





5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	263/270 (97%)	-1.11	0 100 100	6, 14, 33, 38	16 (6%)
1	B	263/270 (97%)	-1.11	0 100 100	5, 13, 33, 44	16 (6%)
1	C	263/270 (97%)	-1.09	0 100 100	5, 14, 33, 40	16 (6%)
1	D	263/270 (97%)	-1.08	0 100 100	6, 14, 33, 37	16 (6%)
1	E	263/270 (97%)	-1.12	0 100 100	5, 13, 32, 43	16 (6%)
1	F	263/270 (97%)	-1.08	0 100 100	5, 15, 33, 42	16 (6%)
1	G	263/270 (97%)	-1.08	0 100 100	5, 15, 33, 42	16 (6%)
1	H	263/270 (97%)	-1.08	0 100 100	5, 14, 33, 38	16 (6%)
1	I	263/270 (97%)	-1.14	0 100 100	5, 13, 33, 43	16 (6%)
1	J	263/270 (97%)	-1.10	0 100 100	5, 14, 32, 40	16 (6%)
1	K	263/270 (97%)	-1.10	0 100 100	5, 14, 32, 43	16 (6%)
1	L	263/270 (97%)	-1.12	0 100 100	5, 13, 33, 43	16 (6%)
All	All	3156/3240 (97%)	-1.10	0 100 100	5, 14, 33, 44	192 (6%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	G	4020	5/5	0.97	0.06	28,28,29,29	5
2	SO4	D	4011	5/5	0.98	0.07	50,51,51,52	0
2	SO4	A	4001	5/5	0.98	0.07	45,45,46,47	0
2	SO4	H	4017	5/5	0.98	0.07	51,52,52,53	0
2	SO4	H	4023	5/5	0.98	0.04	54,54,54,55	0
2	SO4	I	4014	5/5	0.98	0.07	44,44,45,46	0
2	SO4	I	4026	5/5	0.98	0.05	22,22,25,27	0
2	SO4	J	4029	5/5	0.98	0.05	46,47,47,48	5
2	SO4	K	4008	5/5	0.98	0.06	43,43,44,45	0
2	SO4	K	4032	5/5	0.98	0.05	57,58,58,59	0
2	SO4	L	4005	5/5	0.98	0.07	46,47,48,49	0
2	SO4	L	4035	5/5	0.98	0.06	21,22,25,27	0
3	MTA	C	4009	20/20	0.99	0.03	9,11,18,18	0
3	MTA	D	4012	20/20	0.99	0.03	10,11,18,18	0
3	MTA	F	4018	20/20	0.99	0.03	9,12,18,18	0
3	MTA	H	4024	20/20	0.99	0.03	10,12,18,19	0
3	MTA	I	4027	20/20	0.99	0.03	8,10,14,16	0
3	MTA	L	4036	20/20	0.99	0.03	8,10,15,15	0
2	SO4	G	4019	5/5	1.00	0.03	12,12,15,15	0
2	SO4	K	4031	5/5	1.00	0.03	11,11,14,15	0
2	SO4	B	4004	5/5	1.00	0.02	10,11,14,15	0
2	SO4	C	4007	5/5	1.00	0.02	12,13,15,16	0
2	SO4	L	4034	5/5	1.00	0.02	11,11,14,15	0
2	SO4	H	4022	5/5	1.00	0.02	11,12,15,15	0
3	MTA	A	4003	20/20	1.00	0.02	9,10,16,17	0
3	MTA	B	4006	20/20	1.00	0.02	8,10,15,16	0
2	SO4	D	4010	5/5	1.00	0.03	11,12,15,16	0
2	SO4	A	4000	5/5	1.00	0.02	11,11,15,16	0
3	MTA	E	4015	20/20	1.00	0.02	8,10,14,15	0
2	SO4	I	4025	5/5	1.00	0.02	9,9,13,14	0
3	MTA	G	4021	20/20	1.00	0.02	10,12,18,19	0
2	SO4	E	4013	5/5	1.00	0.02	10,11,13,14	0
2	SO4	J	4028	5/5	1.00	0.02	10,11,14,14	0
3	MTA	J	4030	20/20	1.00	0.02	9,11,17,19	0
3	MTA	K	4033	20/20	1.00	0.02	8,11,17,18	0
2	SO4	F	4016	5/5	1.00	0.03	12,13,16,16	0

6.5 Other polymers [i](#)

There are no such residues in this entry.