



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 10:17 PM UTC

PDB ID : 8A85 / pdb_00008a85
Title : Structure of the Reconstructed Ancestor of Phenolic Acid Decarboxylase An-cPAD134
Authors : Mokos, D.; Schrufer, A.; Gruber, K.; Daniel, B.
Deposited on : 2022-06-22
Resolution : 2.67 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.67 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5070 (2.70-2.66)
Clashscore	190562	5409 (2.70-2.66)
Ramachandran outliers	187476	5324 (2.70-2.66)
Sidechain outliers	187428	5324 (2.70-2.66)
RSRZ outliers	180081	5070 (2.70-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	190	 81% 6% • 13%
1	B	190	 82% 6% • 12%
1	C	190	 2% 82% 6% 12%
1	D	190	 84% • • 13%
1	E	190	 84% • 13%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	190	 A horizontal bar chart showing the quality of chain. The bar is divided into four segments: a red segment on the left labeled '5%', a large green segment labeled '83%', a small yellow segment, and a grey segment on the right labeled '13%'. There are two small black dots between the yellow and grey segments.

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 8286 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phenolic acid decarboxylase N134.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	166	1330	848	222	255	5	0	0	0
1	B	167	1352	860	230	257	5	0	1	0
1	C	167	1341	854	226	256	5	0	0	0
1	D	166	1330	848	222	255	5	0	0	0
1	E	166	1330	848	222	255	5	0	0	0
1	F	166	1330	848	222	255	5	0	0	0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	46	Total	O	0	0
			46	46		
2	B	66	Total	O	0	0
			66	66		
2	C	55	Total	O	0	0
			55	55		
2	D	51	Total	O	0	0
			51	51		
2	E	29	Total	O	0	0
			29	29		
2	F	26	Total	O	0	0
			26	26		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Phenolic acid decarboxylase N134




- Molecule 1: Phenolic acid decarboxylase N134




- Molecule 1: Phenolic acid decarboxylase N134




- Molecule 1: Phenolic acid decarboxylase N134



- Molecule 1: Phenolic acid decarboxylase N134



- Molecule 1: Phenolic acid decarboxylase N134



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	97.32Å 106.68Å 114.75Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.95 – 2.67 35.95 – 2.67	Depositor EDS
% Data completeness (in resolution range)	99.9 (35.95-2.67) 99.9 (35.95-2.67)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.73 (at 2.68Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.191 , 0.229 0.191 , 0.229	Depositor DCC
R_{free} test set	1774 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å ²)	36.5	Xtrriage
Anisotropy	0.068	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 25.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8286	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.03	0/1367	1.25	2/1870 (0.1%)
1	B	1.04	1/1389 (0.1%)	1.24	1/1899 (0.1%)
1	C	1.02	1/1378 (0.1%)	1.28	1/1885 (0.1%)
1	D	1.02	0/1367	1.28	1/1870 (0.1%)
1	E	0.99	0/1367	1.27	1/1870 (0.1%)
1	F	1.02	0/1367	1.26	1/1870 (0.1%)
All	All	1.02	2/8235 (0.0%)	1.26	7/11264 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	124	HIS	CE1-NE2	5.91	1.38	1.32
1	B	124	HIS	CE1-NE2	5.60	1.38	1.32

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	117	PHE	CA-CB-CG	6.96	120.76	113.80
1	A	117	PHE	CA-CB-CG	6.64	120.44	113.80
1	B	117	PHE	CA-CB-CG	6.55	120.35	113.80
1	D	117	PHE	CA-CB-CG	6.36	120.16	113.80
1	E	117	PHE	CA-CB-CG	6.21	120.01	113.80
1	F	117	PHE	CA-CB-CG	5.75	119.55	113.80
1	A	34	ILE	CA-C-O	-5.15	115.66	120.34

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1330	0	1262	5	0
1	B	1352	0	1286	6	0
1	C	1341	0	1274	3	0
1	D	1330	0	1262	3	0
1	E	1330	0	1262	4	0
1	F	1330	0	1262	3	0
2	A	46	0	0	0	0
2	B	66	0	0	2	0
2	C	55	0	0	0	0
2	D	51	0	0	0	0
2	E	29	0	0	0	0
2	F	26	0	0	0	0
All	All	8286	0	7608	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:188[B]:ARG:NE	2:B:201:HOH:O	2.44	0.49
1:A:40:ILE:HD13	1:A:186:ILE:HD12	1.93	0.49
1:B:39:LEU:N	1:B:39:LEU:HD23	2.27	0.49
1:F:39:LEU:N	1:F:39:LEU:HD23	2.27	0.49
1:D:39:LEU:HD23	1:D:39:LEU:N	2.27	0.48
1:C:47:TRP:CH2	1:C:154:ILE:HD13	2.50	0.47
1:F:64:HIS:CE1	1:F:186:ILE:HD13	2.50	0.46
1:A:186:ILE:HD12	1:A:186:ILE:HA	1.74	0.44
1:B:64:HIS:CE1	1:B:186:ILE:HD13	2.53	0.43
1:B:188[B]:ARG:CZ	2:B:201:HOH:O	2.65	0.43
1:F:111:LEU:C	1:F:111:LEU:HD23	2.43	0.43
1:A:37:LYS:HE2	1:A:106:LEU:HD13	2.00	0.43
1:E:186:ILE:HD12	1:E:186:ILE:HA	1.90	0.43
1:E:64:HIS:CE1	1:E:186:ILE:HD13	2.55	0.42
1:B:58:THR:CG2	1:B:75:ASP:HA	2.50	0.42
1:B:111:LEU:C	1:B:111:LEU:HD23	2.45	0.41
1:D:64:HIS:CE1	1:D:186:ILE:HD13	2.54	0.41
1:A:111:LEU:C	1:A:111:LEU:HD23	2.46	0.41
1:D:111:LEU:C	1:D:111:LEU:HD23	2.45	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:64:HIS:CD2	1:E:186:ILE:HD13	2.55	0.41
1:C:58:THR:CG2	1:C:75:ASP:HA	2.51	0.40
1:C:111:LEU:C	1:C:111:LEU:HD23	2.46	0.40
1:E:111:LEU:C	1:E:111:LEU:HD23	2.46	0.40
1:A:58:THR:CG2	1:A:75:ASP:HA	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	164/190 (86%)	159 (97%)	5 (3%)	0	100	100
1	B	166/190 (87%)	163 (98%)	3 (2%)	0	100	100
1	C	165/190 (87%)	161 (98%)	4 (2%)	0	100	100
1	D	164/190 (86%)	159 (97%)	5 (3%)	0	100	100
1	E	164/190 (86%)	160 (98%)	4 (2%)	0	100	100
1	F	164/190 (86%)	160 (98%)	4 (2%)	0	100	100
All	All	987/1140 (87%)	962 (98%)	25 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	146/166 (88%)	142 (97%)	4 (3%)	39	67
1	B	148/166 (89%)	144 (97%)	4 (3%)	39	67
1	C	147/166 (89%)	143 (97%)	4 (3%)	39	67
1	D	146/166 (88%)	143 (98%)	3 (2%)	47	73
1	E	146/166 (88%)	143 (98%)	3 (2%)	47	73
1	F	146/166 (88%)	141 (97%)	5 (3%)	32	59
All	All	879/996 (88%)	856 (97%)	23 (3%)	40	68

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	39	LEU
1	A	78	VAL
1	A	171	ASP
1	A	186	ILE
1	B	24	ARG
1	B	39	LEU
1	B	78	VAL
1	B	171	ASP
1	C	39	LEU
1	C	78	VAL
1	C	85	ASN
1	C	171	ASP
1	D	39	LEU
1	D	78	VAL
1	D	171	ASP
1	E	39	LEU
1	E	78	VAL
1	E	171	ASP
1	F	39	LEU
1	F	78	VAL
1	F	165	ASN
1	F	168	ILE
1	F	171	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	26	GLN
1	A	85	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	85	ASN
1	B	123	GLN
1	C	123	GLN
1	C	151	GLN
1	E	26	GLN
1	E	48	GLN
1	F	26	GLN
1	F	123	GLN
1	F	151	GLN
1	F	165	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	166/190 (87%)	-0.41	1 (0%) 85 84	20, 31, 51, 88	0
1	B	167/190 (87%)	-0.43	0 100 100	15, 30, 49, 80	1 (0%)
1	C	167/190 (87%)	-0.37	3 (1%) 67 65	20, 30, 52, 95	0
1	D	166/190 (87%)	-0.29	1 (0%) 85 84	22, 35, 54, 79	0
1	E	166/190 (87%)	-0.35	1 (0%) 85 84	22, 36, 54, 83	0
1	F	166/190 (87%)	0.50	9 (5%) 31 27	24, 50, 76, 90	0
All	All	998/1140 (87%)	-0.23	15 (1%) 72 69	15, 34, 65, 95	1 (0%)

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	25	PRO	5.3
1	A	25	PRO	4.5
1	E	25	PRO	3.9
1	F	25	PRO	3.7
1	C	25	PRO	3.4
1	F	168	ILE	3.4
1	C	117	PHE	3.1
1	F	165	ASN	3.0
1	F	173	VAL	2.9
1	F	148	VAL	2.7
1	C	24	ARG	2.4
1	F	166	CYS	2.2
1	F	32	SER	2.2
1	F	72	TRP	2.1
1	F	167	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.