



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 04:06 AM UTC

PDB ID : 5A9I / pdb_00005a9i
Title : Crystal structure of the extracellular domain of PepT2
Authors : Beale, J.H.; Newstead, S.
Deposited on : 2015-07-21
Resolution : 2.84 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1520 (2.86-2.82)
Clashscore	190562	1559 (2.86-2.82)
Ramachandran outliers	187476	1517 (2.86-2.82)
Sidechain outliers	187428	1518 (2.86-2.82)
RSRZ outliers	180081	1521 (2.86-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	194	10% 82% 16% ..
1	B	194	% 79% 19% .
1	C	194	10% 71% 26% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	CIT	B	1602	-	X	-	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4678 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called SOLUTE CARRIER FAMILY 15 MEMBER 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	194	Total	C	N	O	S	Se	0	0	0
			1515	955	257	298	2	3			
1	B	194	Total	C	N	O	S	Se	0	0	0
			1515	955	257	298	2	3			
1	C	194	Total	C	N	O	S	Se	0	1	0
			1523	960	258	299	2	4			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	408	MSE	-	expression tag	UNP Q63424
A	409	ALA	-	expression tag	UNP Q63424
B	408	MSE	-	expression tag	UNP Q63424
B	409	ALA	-	expression tag	UNP Q63424
C	408	MSE	-	expression tag	UNP Q63424
C	409	ALA	-	expression tag	UNP Q63424

- Molecule 2 is CITRIC ACID (CCD ID: CIT) (formula: C₆H₈O₇).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			13	6	7		
2	B	1	Total	C	O	0	0
			13	6	7		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	43	Total 43	O 43	0	0
4	B	36	Total 36	O 36	0	0
4	C	14	Total 14	O 14	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: SOLUTE CARRIER FAMILY 15 MEMBER 2



• Molecule 1: SOLUTE CARRIER FAMILY 15 MEMBER 2



• Molecule 1: SOLUTE CARRIER FAMILY 15 MEMBER 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	95.75Å 95.75Å 165.93Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.87 – 2.84 47.87 – 2.85	Depositor EDS
% Data completeness (in resolution range)	99.7 (47.87-2.84) 99.9 (47.87-2.85)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.99 (at 2.86Å)	Xtriage
Refinement program	BUSTER 2.11.5	Depositor
R, R_{free}	0.193 , 0.246 0.202 , 0.251	Depositor DCC
R_{free} test set	1090 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å ²)	65.3	Xtriage
Anisotropy	0.275	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 76.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.028 for -h,-k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4678	wwPDB-VP
Average B, all atoms (Å ²)	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CIT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.87	2/1540 (0.1%)	1.28	8/2083 (0.4%)
1	B	0.86	0/1540	1.25	3/2083 (0.1%)
1	C	0.79	0/1548	1.31	7/2093 (0.3%)
All	All	0.84	2/4628 (0.0%)	1.28	18/6259 (0.3%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	514	MSE	SE-CE	-6.49	1.75	1.95
1	A	450	THR	CA-C	5.03	1.58	1.53

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	435	ASN	CA-C-N	7.10	131.89	122.30
1	C	435	ASN	C-N-CA	7.10	131.89	122.30
1	A	449	THR	CA-C-N	6.95	132.38	122.24
1	A	449	THR	C-N-CA	6.95	132.38	122.24
1	C	577	PHE	CA-CB-CG	6.44	120.24	113.80
1	C	532	ASP	CA-CB-CG	6.00	118.60	112.60
1	A	600	ASP	CA-CB-CG	5.99	118.59	112.60
1	A	450	THR	CB-CA-C	5.86	119.77	111.74
1	C	563	THR	N-CA-C	-5.84	100.88	109.18
1	A	518	ARG	CB-CG-CD	-5.73	98.13	111.30
1	B	420	ASN	CA-CB-CG	5.72	118.32	112.60
1	A	436	ASN	CA-CB-CG	5.29	117.89	112.60
1	C	539	VAL	CA-C-N	5.28	125.12	120.10
1	C	539	VAL	C-N-CA	5.28	125.12	120.10
1	B	600	ASP	CA-CB-CG	5.22	117.82	112.60
1	A	496	GLU	N-CA-C	-5.21	106.77	113.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	496	GLU	N-CA-C	-5.10	107.06	113.28
1	A	451	HIS	CA-CB-CG	5.05	118.85	113.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1515	0	1493	13	0
1	B	1515	0	1493	16	0
1	C	1523	0	1501	27	0
2	B	26	0	10	1	0
3	B	6	0	8	0	0
4	A	43	0	0	1	0
4	B	36	0	0	1	0
4	C	14	0	0	2	0
All	All	4678	0	4505	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:521:ASN:HB2	1:B:539:VAL:HG22	1.62	0.79
1:C:527:LEU:HD21	1:C:561:CYS:HB3	1.68	0.76
1:C:559:VAL:HG21	1:C:572:LEU:HD22	1.67	0.75
1:A:559:VAL:HG21	1:A:572:LEU:HD22	1.69	0.73
1:B:514:MSE:HG2	1:B:550:THR:HG22	1.81	0.62
1:C:416:LEU:HD11	1:C:489:LEU:HB2	1.83	0.60
1:B:517:ILE:HG22	1:B:581:TYR:HB2	1.84	0.60
1:C:553:ARG:HG2	1:C:577:PHE:HB2	1.84	0.60
1:B:562:GLU:HB3	1:B:567:VAL:HG12	1.82	0.59
1:A:523:LEU:HD11	1:A:593:LEU:HD21	1.85	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:517:ILE:HG22	1:C:581:TYR:HB2	1.83	0.59
1:C:418:VAL:O	1:C:449:THR:HG21	2.03	0.58
1:C:505:ASP:HA	1:C:518:ARG:NH2	2.21	0.56
1:C:421:LEU:HD12	1:C:586:THR:HG21	1.88	0.55
1:B:435:ASN:HB3	2:B:1603:CIT:H22	1.88	0.54
1:C:505:ASP:HA	1:C:518:ARG:HH22	1.73	0.53
1:C:530:SER:HB2	1:C:560:HIS:HB3	1.89	0.53
1:C:503:VAL:HG21	1:C:520:ILE:HD11	1.90	0.53
1:C:562:GLU:HG2	1:C:567:VAL:HG12	1.90	0.53
1:A:466:PHE:CE1	1:A:479:HIS:HB2	2.44	0.52
1:A:429:THR:HG23	1:A:440:VAL:HG22	1.92	0.52
1:C:429:THR:HG23	1:C:440:VAL:HG22	1.91	0.52
1:C:450:THR:HA	4:C:2003:HOH:O	2.09	0.52
1:C:424:GLY:HA3	1:C:470:TYR:CE1	2.46	0.51
4:A:2009:HOH:O	1:B:451:HIS:HB3	2.11	0.50
1:C:600:ASP:O	1:C:601:ILE:HB	2.11	0.49
1:A:451:HIS:CD2	1:B:451:HIS:CD2	3.00	0.49
1:B:539:VAL:HB	1:B:546:SER:HB3	1.94	0.49
1:B:429:THR:HG23	1:B:440:VAL:HG22	1.94	0.48
1:C:492:HIS:CE1	1:C:499:SER:HB3	2.48	0.48
1:B:421:LEU:HG	1:B:446:PHE:CE1	2.49	0.47
1:C:527:LEU:CD2	1:C:561:CYS:HB3	2.40	0.47
1:B:503:VAL:HG21	1:B:520:ILE:HD11	1.97	0.46
1:B:421:LEU:HD22	1:B:490:LEU:HD11	1.98	0.46
1:B:527:LEU:HB3	1:B:539:VAL:HG13	1.96	0.45
1:A:521:ASN:HB2	1:A:539:VAL:HG22	1.98	0.45
1:C:448:ASN:HA	4:C:2002:HOH:O	2.16	0.45
1:C:415:PHE:CE1	1:C:454:LYS:HG2	2.52	0.44
1:C:420:ASN:HD22	1:C:444:SER:C	2.25	0.44
1:C:414:ILE:HG23	1:C:457:LEU:HD11	1.99	0.44
1:A:419:LEU:HD22	1:A:596:TRP:HZ3	1.82	0.43
1:A:466:PHE:HE1	1:A:479:HIS:HB2	1.83	0.43
1:B:506:THR:HB	1:B:548:TYR:CE2	2.54	0.43
1:A:521:ASN:HB3	1:A:540:GLY:O	2.19	0.43
1:A:451:HIS:CD2	1:A:451:HIS:C	2.96	0.43
1:B:514:MSE:HG3	1:B:551:VAL:O	2.19	0.43
1:A:408:MSE:SE	4:B:2001:HOH:O	2.87	0.42
1:C:501[A]:MSE:SE	1:C:543:TYR:HB3	2.70	0.42
1:C:503:VAL:HG21	1:C:520:ILE:CD1	2.49	0.41
1:A:514:MSE:HE2	1:A:550:THR:HG22	2.02	0.41
1:C:501[B]:MSE:HE2	1:C:543:TYR:HB2	2.03	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:420:ASN:ND2	1:C:445:SER:HA	2.36	0.40
1:A:413:GLU:CD	1:A:454:LYS:HD2	2.46	0.40
1:B:548:TYR:CE1	1:B:580:THR:HG23	2.56	0.40
1:C:417:GLN:HB2	1:C:452:TYR:CE1	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	192/194 (99%)	177 (92%)	15 (8%)	0	100	100
1	B	192/194 (99%)	182 (95%)	10 (5%)	0	100	100
1	C	193/194 (100%)	179 (93%)	14 (7%)	0	100	100
All	All	577/582 (99%)	538 (93%)	39 (7%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	171/168 (102%)	159 (93%)	12 (7%)	14	30
1	B	171/168 (102%)	152 (89%)	19 (11%)	6	12
1	C	172/168 (102%)	155 (90%)	17 (10%)	7	16

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	514/504 (102%)	466 (91%)	48 (9%)	8 19

All (48) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	437	SER
1	A	441	GLU
1	A	442	SER
1	A	451	HIS
1	A	527	LEU
1	A	531	LEU
1	A	555	LYS
1	A	560	HIS
1	A	567	VAL
1	A	569	SER
1	A	572	LEU
1	A	586	THR
1	B	408	MSE
1	B	437	SER
1	B	441	GLU
1	B	442	SER
1	B	448	ASN
1	B	449	THR
1	B	469	LYS
1	B	489	LEU
1	B	514	MSE
1	B	518	ARG
1	B	527	LEU
1	B	531	LEU
1	B	539	VAL
1	B	559	VAL
1	B	562	GLU
1	B	568	PHE
1	B	569	SER
1	B	572	LEU
1	B	593	LEU
1	C	408	MSE
1	C	425	ASP
1	C	437	SER
1	C	441	GLU
1	C	442	SER
1	C	476	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	501[A]	MSE
1	C	501[B]	MSE
1	C	518	ARG
1	C	527	LEU
1	C	531	LEU
1	C	549	ARG
1	C	553	ARG
1	C	566	LYS
1	C	572	LEU
1	C	588	ILE
1	C	601	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	447	GLN
1	A	451	HIS
1	A	477	ASN
1	A	594	GLN
1	B	435	ASN
1	B	447	GLN
1	B	477	ASN
1	B	492	HIS
1	B	560	HIS
1	B	594	GLN
1	C	412	GLN
1	C	417	GLN
1	C	420	ASN
1	C	448	ASN
1	C	488	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	CIT	B	1602	-	12,12,12	1.64	4 (33%)	17,17,17	1.97	5 (29%)
2	CIT	B	1603	-	12,12,12	1.63	4 (33%)	17,17,17	2.09	5 (29%)
3	GOL	B	1604	-	5,5,5	0.12	0	5,5,5	0.17	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CIT	B	1602	-	-	11/16/16/16	-
2	CIT	B	1603	-	-	6/16/16/16	-
3	GOL	B	1604	-	-	0/4/4/4	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1603	CIT	O5-C6	3.41	1.32	1.22
2	B	1602	CIT	O5-C6	3.37	1.32	1.22
2	B	1603	CIT	O1-C1	2.98	1.31	1.22
2	B	1602	CIT	O2-C1	-2.91	1.21	1.30
2	B	1602	CIT	O1-C1	2.77	1.31	1.22
2	B	1603	CIT	O2-C1	-2.49	1.22	1.30
2	B	1603	CIT	O6-C6	-2.18	1.22	1.30
2	B	1602	CIT	O6-C6	-2.07	1.23	1.30

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1602	CIT	O6-C6-C3	4.85	122.45	113.14
2	B	1603	CIT	O6-C6-C3	4.84	122.43	113.14
2	B	1602	CIT	O5-C6-C3	-4.22	113.91	122.09
2	B	1603	CIT	O5-C6-C3	-3.93	114.47	122.09
2	B	1603	CIT	O1-C1-C2	-3.77	112.27	122.95
2	B	1602	CIT	O1-C1-C2	-3.15	114.04	122.95
2	B	1603	CIT	O2-C1-C2	2.86	123.39	114.35
2	B	1602	CIT	O2-C1-C2	2.61	122.63	114.35
2	B	1603	CIT	C3-C4-C5	2.36	120.38	113.92
2	B	1602	CIT	C3-C4-C5	2.36	120.36	113.92

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1602	CIT	O7-C3-C4-C5
2	B	1602	CIT	C6-C3-C4-C5
2	B	1602	CIT	O7-C3-C6-O5
2	B	1602	CIT	O7-C3-C6-O6
2	B	1603	CIT	O7-C3-C6-O5
2	B	1603	CIT	O7-C3-C6-O6
2	B	1603	CIT	C4-C3-C6-O5
2	B	1603	CIT	C4-C3-C6-O6
2	B	1602	CIT	C2-C3-C4-C5
2	B	1602	CIT	C2-C3-C6-O5
2	B	1602	CIT	C2-C3-C6-O6
2	B	1602	CIT	C4-C3-C6-O6
2	B	1602	CIT	C4-C3-C6-O5
2	B	1603	CIT	C1-C2-C3-C4
2	B	1602	CIT	C3-C4-C5-O3
2	B	1603	CIT	C1-C2-C3-O7
2	B	1602	CIT	C3-C4-C5-O4

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1603	CIT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	191/194 (98%)	-0.23	0 100 100	41, 60, 101, 122	0
1	B	191/194 (98%)	-0.25	1 (0%) 87 85	40, 59, 98, 129	0
1	C	191/194 (98%)	0.80	20 (10%) 11 9	58, 96, 135, 161	0
All	All	573/582 (98%)	0.11	21 (3%) 45 35	40, 71, 122, 161	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	459	ALA	3.6
1	C	513	GLY	3.5
1	C	601	ILE	3.4
1	C	498	ILE	3.4
1	C	502	LEU	3.1
1	C	508	ILE	2.9
1	B	450	THR	2.9
1	C	433	SER	2.9
1	C	510	PRO	2.7
1	C	477	ASN	2.6
1	C	467	HIS	2.6
1	C	511	ALA	2.5
1	C	475	VAL	2.5
1	C	449	THR	2.4
1	C	596	TRP	2.3
1	C	489	LEU	2.3
1	C	468	LEU	2.3
1	C	506	THR	2.2
1	C	470	TYR	2.2
1	C	447	GLN	2.2
1	C	525	LYS	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CIT	B	1603	13/13	0.64	0.16	107,111,119,120	0
3	GOL	B	1604	6/6	0.82	0.18	84,87,88,88	0
2	CIT	B	1602	13/13	0.93	0.09	94,101,105,105	0

6.5 Other polymers [i](#)

There are no such residues in this entry.