



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 05:52 AM UTC

PDB ID : 1ABW / pdb_00001abw
Title : DEOXY RHB1.1 (RECOMBINANT HEMOGLOBIN)
Authors : Kundrot, C.E.; Kroeger, K.S.
Deposited on : 1997-01-29
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	283	 87% 11% .
2	B	146	 84% 12% . .
2	D	146	 80% 16% .

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5017 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HEMOGLOBIN-BASED BLOOD SUBSTITUTE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	283	Total	C	N	O	S	21	9	0
			2178	1395	383	392	8			

- Molecule 2 is a protein called HEMOGLOBIN-BASED BLOOD SUBSTITUTE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	146	Total	C	N	O	S	0	1	0
			1131	730	197	200	4			
2	D	146	Total	C	N	O	S	0	1	0
			1129	729	195	200	5			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	108	LYS	ASN	conflict	UNP P68871
D	108	LYS	ASN	conflict	UNP P68871

- Molecule 3 is METHIONINE (CCD ID: MET) (formula: C₅H₁₁NO₂S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 8	C 5	N 1	O 1	S 1	4	1
3	A	1	Total 8	C 5	N 1	O 1	S 1	4	1

- Molecule 4 is LEUCINE (CCD ID: LEU) (formula: $C_6H_{13}NO_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O		1
			9	6	1	2	9	
4	A	1	Total	C	N	O		1
			9	6	1	2	9	

- Molecule 5 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
5	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
5	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
5	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
5	D	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	212	Total O 212 212	0	0
6	B	86	Total O 86 86	0	0
6	D	75	Total O 75 75	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: HEMOGLOBIN-BASED BLOOD SUBSTITUTE

Chain A:  87% 11% .



• Molecule 2: HEMOGLOBIN-BASED BLOOD SUBSTITUTE

Chain B:  84% 12% ..



• Molecule 2: HEMOGLOBIN-BASED BLOOD SUBSTITUTE

Chain D:  80% 16% .



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	62.90Å 81.98Å 53.92Å 90.00° 98.99° 90.00°	Depositor
Resolution (Å)	5.00 – 2.00	Depositor
% Data completeness (in resolution range)	86.2 (5.00-2.00)	Depositor
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.164 , 0.232	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	5017	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.78	0/2248	1.04	9/3054 (0.3%)
2	B	0.86	4/1166 (0.3%)	1.11	12/1581 (0.8%)
2	D	0.85	0/1163	1.07	7/1576 (0.4%)
All	All	0.82	4/4577 (0.1%)	1.06	28/6211 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	3	LEU	CA-CB	-6.59	1.42	1.53
2	B	2	HIS	CA-C	6.35	1.61	1.52
2	B	55	MET	SD-CE	-5.94	1.64	1.79
2	B	3	LEU	CA-C	5.43	1.60	1.52

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	47	ASP	N-CA-C	-8.77	94.29	108.41
1	A	189	ASP	N-CA-C	-7.79	96.58	108.96
2	D	94	ASP	N-CA-C	7.59	119.63	111.36
2	B	3	LEU	N-CA-C	-7.16	95.54	110.80
2	B	2	HIS	N-CA-CB	6.87	122.10	110.49
1	A	236	ASP	CA-C-N	6.81	127.08	119.32

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	236	ASP	C-N-CA	6.81	127.08	119.32
2	B	2	HIS	CB-CA-C	6.78	123.91	110.42
2	B	49	SER	N-CA-C	6.71	119.60	111.82
2	D	2	HIS	N-CA-C	6.57	119.31	109.25
2	B	2	HIS	CA-C-N	6.37	133.70	121.54
2	B	2	HIS	C-N-CA	6.37	133.70	121.54
2	B	42	PHE	N-CA-C	6.32	120.79	111.87
1	A	178	PHE	CA-C-N	6.05	126.49	119.47
1	A	178	PHE	C-N-CA	6.05	126.49	119.47
2	B	19	ASN	N-CA-C	-6.00	97.92	108.23
2	D	36	PRO	N-CA-C	5.85	121.44	113.84
2	D	49	SER	N-CA-C	5.77	117.57	111.28
1	A	89	HIS	N-CA-C	5.57	120.21	113.41
2	B	143	HIS	N-CA-C	5.46	118.73	111.75
2	B	94	ASP	N-CA-C	5.43	117.28	111.36
2	B	34	VAL	N-CA-C	5.33	116.11	110.72
1	A	105	LEU	N-CA-C	-5.24	105.65	111.36
1	A	15	GLY	N-CA-C	-5.19	106.48	112.50
2	D	143	HIS	N-CA-C	5.18	117.59	111.33
2	D	90	GLU	CB-CA-C	-5.12	102.30	110.79
2	D	20	VAL	N-CA-C	5.08	116.42	110.62
2	B	18	VAL	N-CA-C	5.03	115.82	108.58

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	2	HIS	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2178	0	2181	24	0
2	B	1131	0	1128	13	0
2	D	1129	0	1128	12	0
3	A	16	0	16	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	18	0	22	0	0
5	A	86	0	60	1	0
5	B	43	0	30	1	0
5	D	43	0	30	0	0
6	A	212	0	0	2	3
6	B	86	0	0	2	1
6	D	75	0	0	4	2
All	All	5017	0	4595	45	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:278:LEU:HD12	5:A:285:HEM:HBB2	1.50	0.93
1:A:1[B]:MET:HG3	1:A:131:SER:OG	1.84	0.78
1:A:255:LEU:HB3	1:A:258:GLU:HG2	1.71	0.71
2:B:143:HIS:HB3	6:B:192:HOH:O	1.93	0.68
2:B:1:MET:HB2	2:B:78:LEU:O	1.95	0.65
1:A:21:ALA:HB1	1:A:63:ALA:HB1	1.84	0.59
1:A:1[B]:MET:HE2	1:A:131:SER:HA	1.84	0.59
1:A:1[B]:MET:HE2	1:A:131:SER:OG	2.03	0.58
5:B:147:HEM:HMC2	5:B:147:HEM:HBC2	1.85	0.56
1:A:143[A]:VAL:HG12	1:A:144[A]:LEU:H	1.71	0.56
2:D:114:LEU:HD21	6:D:174:HOH:O	2.05	0.56
1:A:236:ASP:OD1	1:A:238:VAL:HG22	2.06	0.56
2:D:51:PRO:O	2:D:55:MET:HG2	2.07	0.54
1:A:143[A]:VAL:HG12	1:A:144[A]:LEU:N	2.23	0.54
1:A:169:GLU:OE1	1:A:254:HIS:HE1	1.92	0.53
2:D:20:VAL:HG13	2:D:68:LEU:HB3	1.91	0.53
1:A:234[A]:ARG:HD3	2:B:37:TRP:HA	1.93	0.50
2:D:144:LYS:HD3	6:D:191:HOH:O	2.10	0.50
2:B:1:MET:CE	2:B:133:VAL:HA	2.43	0.49
1:A:92:ARG:HB3	2:D:37:TRP:HB2	1.96	0.47
2:B:144:LYS:HD3	6:B:225:HOH:O	2.15	0.47
2:D:100:PRO:HB3	2:D:142:ALA:HB2	1.96	0.47
1:A:83:LEU:HD12	1:A:83:LEU:HA	1.77	0.47
1:A:234[A]:ARG:HD2	2:B:40:ARG:HB3	1.97	0.46
2:D:95:LYS:HD3	2:D:95:LYS:HA	1.78	0.45
2:D:124:PRO:HB2	2:D:125:PRO:HD3	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:7:LYS:O	1:A:11:LYS:HG2	2.17	0.45
1:A:172:GLU:HG3	6:A:486:HOH:O	2.17	0.44
1:A:1[B]:MET:HE2	1:A:131:SER:CA	2.47	0.44
1:A:283:ARG:HG3	2:B:37:TRP:HZ3	1.81	0.44
1:A:256:PRO:HD2	6:A:409:HOH:O	2.18	0.44
2:B:38:THR:HG22	2:B:102:ASN:OD1	2.17	0.43
1:A:234[B]:ARG:HD3	2:B:37:TRP:HA	2.00	0.43
2:B:51:PRO:O	2:B:55:MET:HG2	2.17	0.43
2:D:97:HIS:CE1	6:D:198:HOH:O	2.72	0.43
1:A:6:ASP:O	1:A:10:VAL:HG23	2.19	0.42
1:A:170:ALA:CB	1:A:247:LEU:HD13	2.50	0.42
2:B:100:PRO:HB3	2:B:142:ALA:HB2	2.02	0.41
2:B:24:GLY:HA2	2:B:68:LEU:HG	2.02	0.41
1:A:68:ASN:OD1	1:A:72:HIS:HD2	2.02	0.41
2:D:18:VAL:HG11	6:D:174:HOH:O	2.21	0.41
2:B:1:MET:HE1	2:B:133:VAL:HA	2.02	0.41
2:D:82:LYS:HE3	2:D:143:HIS:CD2	2.56	0.40
2:D:144:LYS:HA	2:D:144:LYS:HD2	1.98	0.40
1:A:278:LEU:HD23	1:A:278:LEU:HA	1.88	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:480:HOH:O	6:B:231:HOH:O[1_554]	0.01	2.19
6:A:468:HOH:O	6:D:208:HOH:O[1_556]	0.05	2.15
6:A:426:HOH:O	6:D:168:HOH:O[2_556]	2.15	0.05

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	285/283 (101%)	280 (98%)	5 (2%)	0	100	100
2	B	145/146 (99%)	140 (97%)	4 (3%)	1 (1%)	18	14
2	D	144/146 (99%)	142 (99%)	2 (1%)	0	100	100
All	All	574/575 (100%)	562 (98%)	11 (2%)	1 (0%)	43	42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	2	HIS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/226 (102%)	223 (96%)	8 (4%)	32	32
2	B	119/118 (101%)	114 (96%)	5 (4%)	26	25
2	D	119/118 (101%)	107 (90%)	12 (10%)	7	4
All	All	469/462 (102%)	444 (95%)	25 (5%)	20	18

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	83	LEU
1	A	92	ARG
1	A	105	LEU
1	A	109	LEU
1	A	137	THR
1	A	247	LEU
1	A	248	LEU
1	A	258	GLU
2	B	1	MET
2	B	2	HIS
2	B	5	PRO
2	B	51	PRO

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	78	LEU
2	D	1[A]	MET
2	D	1[B]	MET
2	D	6	GLU
2	D	14	LEU
2	D	26	GLU
2	D	32	LEU
2	D	68	LEU
2	D	75	LEU
2	D	78	LEU
2	D	120	LYS
2	D	124	PRO
2	D	144	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	72	HIS
1	A	89	HIS
1	A	192	HIS
1	A	214	HIS
1	A	254	HIS
2	B	143	HIS
2	D	63	HIS
2	D	139	ASN
2	D	143	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	HEM	B	147	2	50,50,50	1.56	10 (20%)	67,82,82	0.88	2 (2%)
5	HEM	D	147	2	50,50,50	1.49	8 (16%)	67,82,82	1.06	4 (5%)
4	LEU	A	2002[C]	3	6,8,8	0.74	0	5,10,10	0.25	0
3	MET	A	2001[C]	4	6,7,8	0.58	0	2,7,9	0.56	0
5	HEM	A	284	6,1	50,50,50	1.52	9 (18%)	67,82,82	0.89	1 (1%)
5	HEM	A	285	6,1	50,50,50	1.53	9 (18%)	67,82,82	0.94	4 (5%)
4	LEU	A	1002[B]	3	6,8,8	0.74	0	5,10,10	0.31	0
3	MET	A	1001[B]	4	6,7,8	0.53	0	2,7,9	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	HEM	B	147	2	-	0/14/54/54	-
5	HEM	D	147	2	-	6/14/54/54	-
4	LEU	A	2002[C]	3	-	2/8/8/8	-
3	MET	A	2001[C]	4	-	1/5/6/8	-
5	HEM	A	284	6,1	-	6/14/54/54	-
5	HEM	A	285	6,1	-	4/14/54/54	-
4	LEU	A	1002[B]	3	-	2/8/8/8	-
3	MET	A	1001[B]	4	-	1/5/6/8	-

All (36) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	285	HEM	FE-NA	4.71	2.10	1.95
5	A	284	HEM	FE-NA	4.68	2.10	1.95

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	D	147	HEM	FE-NA	4.04	2.08	1.95
5	A	284	HEM	FE-NB	4.04	2.07	1.94
5	D	147	HEM	FE-NC	3.81	2.07	1.95
5	B	147	HEM	FE-NC	3.78	2.07	1.95
5	B	147	HEM	FE-ND	3.54	2.05	1.94
5	D	147	HEM	FE-NB	3.50	2.05	1.94
5	B	147	HEM	CAB-C3B	-3.48	1.38	1.47
5	B	147	HEM	FE-NA	3.40	2.06	1.95
5	B	147	HEM	FE-NB	3.40	2.05	1.94
5	A	284	HEM	CAB-C3B	-3.27	1.38	1.47
5	D	147	HEM	CBC-CAC	3.26	1.46	1.30
5	A	285	HEM	FE-NB	3.26	2.04	1.94
5	A	285	HEM	CBC-CAC	3.21	1.45	1.30
5	A	285	HEM	CMC-C2C	3.17	1.57	1.50
5	B	147	HEM	CBB-CAB	3.07	1.45	1.30
5	D	147	HEM	CMD-C2D	2.95	1.56	1.50
5	B	147	HEM	CAC-C3C	-2.93	1.39	1.47
5	A	284	HEM	CAC-C3C	-2.84	1.39	1.47
5	A	285	HEM	CMB-C2B	2.81	1.56	1.50
5	A	285	HEM	CAB-C3B	-2.81	1.39	1.47
5	A	285	HEM	CAC-C3C	-2.76	1.40	1.47
5	B	147	HEM	CMC-C2C	2.75	1.56	1.50
5	B	147	HEM	CAD-C3D	2.50	1.57	1.51
5	A	284	HEM	FE-NC	2.48	2.03	1.95
5	D	147	HEM	CBB-CAB	2.46	1.42	1.30
5	A	284	HEM	CMD-C2D	2.44	1.55	1.50
5	B	147	HEM	CBC-CAC	2.43	1.42	1.30
5	A	285	HEM	O2D-CGD	-2.31	1.23	1.30
5	A	284	HEM	FE-ND	2.31	2.02	1.94
5	D	147	HEM	CAB-C3B	-2.25	1.41	1.47
5	A	284	HEM	CBC-CAC	2.14	1.40	1.30
5	D	147	HEM	FE-ND	2.13	2.01	1.94
5	A	285	HEM	CMD-C2D	2.13	1.55	1.50
5	A	284	HEM	CMB-C2B	2.09	1.55	1.50

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	D	147	HEM	CAD-C3D-C4D	4.01	131.69	124.70
5	D	147	HEM	CAD-C3D-C2D	-3.75	120.84	127.87
5	A	285	HEM	CMB-C2B-C1B	2.88	129.54	125.03
5	D	147	HEM	CMD-C2D-C1D	2.61	129.12	125.03

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	285	HEM	CMA-C3A-C2A	2.40	130.72	125.62
5	B	147	HEM	O1D-CGD-CBD	-2.39	115.51	123.09
5	A	285	HEM	CMA-C3A-C4A	-2.18	122.10	125.42
5	A	284	HEM	CBD-CAD-C3D	-2.18	106.52	112.53
5	D	147	HEM	CBB-CAB-C3B	-2.13	116.89	127.53
5	B	147	HEM	O2D-CGD-CBD	2.10	120.62	114.00
5	A	285	HEM	CMB-C2B-C3B	-2.02	123.55	128.43

There are no chirality outliers.

All (22) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	1002[B]	LEU	O-C-CA-N
5	A	284	HEM	C2B-C3B-CAB-CBB
5	A	284	HEM	C4B-C3B-CAB-CBB
5	A	284	HEM	C2C-C3C-CAC-CBC
5	A	285	HEM	C2B-C3B-CAB-CBB
5	A	285	HEM	C4B-C3B-CAB-CBB
5	D	147	HEM	C2B-C3B-CAB-CBB
4	A	2002[C]	LEU	OXT-C-CA-N
5	D	147	HEM	C4B-C3B-CAB-CBB
3	A	2001[C]	MET	CB-CG-SD-CE
4	A	2002[C]	LEU	O-C-CA-N
4	A	1002[B]	LEU	OXT-C-CA-N
5	D	147	HEM	C2C-C3C-CAC-CBC
5	A	284	HEM	C4C-C3C-CAC-CBC
3	A	1001[B]	MET	CA-CB-CG-SD
5	A	284	HEM	CAD-CBD-CGD-O1D
5	A	285	HEM	CAD-CBD-CGD-O1D
5	D	147	HEM	CAD-CBD-CGD-O1D
5	A	284	HEM	CAD-CBD-CGD-O2D
5	D	147	HEM	CAD-CBD-CGD-O2D
5	A	285	HEM	CAD-CBD-CGD-O2D
5	D	147	HEM	C4C-C3C-CAC-CBC


There are no ring outliers.


2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	147	HEM	1	0
5	A	285	HEM	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.