



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:02 PM UTC

PDB ID : 8ABX / pdb_00008abx
Title : Crystal structure of IDO1 in complex with Apoxidole-1
Authors : Dotsch, L.; Ziegler, S.; Waldmann, H.; Gasper, R.
Deposited on : 2022-07-05
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.66-1.66)
Clashscore	190562	2662 (1.66-1.66)
Ramachandran outliers	187476	2621 (1.66-1.66)
Sidechain outliers	187428	2621 (1.66-1.66)
RSRZ outliers	180081	2564 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	404	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 6615 atoms, of which 3258 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Indoleamine 2,3-dioxygenase 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	369	6074	1943	3050	510	550	21	0	20	0

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP P14902
A	-2	PRO	-	expression tag	UNP P14902
A	-1	LEU	-	expression tag	UNP P14902
A	0	GLY	-	expression tag	UNP P14902
A	1	SER	-	expression tag	UNP P14902
A	2	GLY	-	expression tag	UNP P14902
A	3	ILE	-	expression tag	UNP P14902
A	4	GLN	-	expression tag	UNP P14902

- Molecule 2 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: C₈H₁₈O₅).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			31	8	18	5		
2	A	1	Total	C	H	O	0	0
			31	8	18	5		

- Molecule 3 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: $C_6H_{14}O_4$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			24	6	14	4		
3	A	1	Total	C	H	O	0	0
			24	6	14	4		
3	A	1	Total	C	H	O	0	0
			24	6	14	4		
3	A	1	Total	C	H	O	0	0
			24	6	14	4		
3	A	1	Total	C	H	O	0	0
			24	6	14	4		

- Molecule 4 is O1-tert-butyl O2-ethyl O5-methyl (E,5R)-5-(1-methylindol-2-yl)-5-[(4-methylphenyl)sulfonylamino]pent-2-ene-1,2,5-tricarboxylate (CCD ID: N39) (formula: $C_{31}H_{36}N_2O_8S$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
4	A	1	Total	C	H	N	O	S	0	1
			156	62	72	4	16	2		

- Molecule 5 is O2-tert-butyl O3-ethyl O6-methyl (2S,6R)-6-(1-methylindol-2-yl)-2,5-dihydro-1H-pyridine-2,3,6-tricarboxylate (CCD ID: N2U) (formula: $C_{24}H_{30}N_2O_6$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	H	N	O		
			62	24	30	2	6	0	1


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	141	Total 141	O 141	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Indoleamine 2,3-dioxygenase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	106.48Å 109.49Å 36.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.17 – 1.65 38.17 – 1.65	Depositor EDS
% Data completeness (in resolution range)	99.6 (38.17-1.65) 99.6 (38.17-1.65)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.22 (at 1.65Å)	Xtriage
Refinement program	PHENIX 1.20.1-4487_4487	Depositor
R, R_{free}	0.192 , 0.210 0.192 , 0.210	Depositor DCC
R_{free} test set	2596 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	26.4	Xtriage
Anisotropy	0.287	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 31.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.008 for k,h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6615	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.26% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: N39, N2U, PG4, PGE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.19	0/3168	0.43	0/4279

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3024	3050	2968	26	0
2	A	26	36	36	0	0
3	A	50	70	70	4	0
4	A	84	72	0	11	0
5	A	32	30	0	1	0
6	A	141	0	0	1	0
All	All	3357	3258	3074	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:508[B]:N39:C24	4:A:508[B]:N39:N32	1.70	1.53
4:A:508[A]:N39:C24	4:A:508[A]:N39:N32	1.71	1.51
4:A:508[A]:N39:C24	4:A:508[A]:N39:C11	2.54	0.86
4:A:508[B]:N39:C24	4:A:508[B]:N39:C11	2.57	0.82
1:A:355:LEU:HD11	1:A:385:MET:HG3	1.73	0.70
1:A:84:GLY:O	1:A:88:MET:HG2	1.96	0.65
1:A:339:LEU:O	1:A:343:ARG:HG3	2.02	0.59
4:A:508[B]:N39:N32	4:A:508[B]:N39:C25	2.61	0.57
4:A:508[A]:N39:C24	4:A:508[A]:N39:S33	2.92	0.57
1:A:265:GLY:H	4:A:508[A]:N39:C40	2.19	0.56
1:A:57:GLU:H	1:A:57:GLU:CD	2.15	0.55
1:A:95:GLY:HA3	1:A:232:ILE:HD13	1.90	0.53
1:A:44:LYS:HD3	1:A:152:PHE:CE1	2.45	0.51
4:A:508[A]:N39:N32	4:A:508[A]:N39:C25	2.64	0.51
1:A:175:ALA:HB2	6:A:661:HOH:O	2.11	0.50
1:A:393:SER:OG	1:A:397:LYS:HE2	2.11	0.50
1:A:227:PHE:HA	1:A:231:ARG:HD3	1.93	0.50
1:A:296:ARG:O	1:A:299:MET:HB2	2.13	0.49
1:A:319:PHE:CZ	1:A:323:LYS:HE2	2.48	0.49
1:A:19:GLU:OE1	1:A:19:GLU:N	2.27	0.48
4:A:508[B]:N39:C24	4:A:508[B]:N39:S33	2.97	0.46
1:A:58:ARG:HD3	3:A:507:PGE:H52	1.97	0.46
1:A:58:ARG:HD3	3:A:507:PGE:O4	2.15	0.46
1:A:267:SER:HB3	4:A:508[A]:N39:C40	2.47	0.45
1:A:3:ILE:HD12	1:A:306:PHE:CE1	2.52	0.44
1:A:58:ARG:NH1	3:A:507:PGE:O2	2.51	0.44
1:A:115:SER:HB3	1:A:120:LEU:O	2.18	0.43
1:A:103:LEU:HD11	1:A:107:ILE:HD11	1.99	0.43
1:A:58:ARG:O	3:A:507:PGE:H5	2.19	0.42
1:A:279:ILE:O	1:A:280:GLN:C	2.63	0.41
1:A:190[A]:MET:HB2	1:A:192:GLU:HG3	2.03	0.41
5:A:509[C]:N2U:O26	5:A:509[C]:N2U:C19	2.68	0.41
1:A:26:PRO:O	1:A:27:ASN:C	2.64	0.41
1:A:57:GLU:CD	1:A:57:GLU:N	2.78	0.41
1:A:270:PHE:HB2	4:A:508[A]:N39:C37	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	382/404 (95%)	371 (97%)	11 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	343/349 (98%)	341 (99%)	2 (1%)	78	68

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	222[A]	ASN
1	A	222[B]	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	76	GLN
1	A	212	GLN
1	A	215	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PGE	A	503	-	9,9,9	0.32	0	8,8,8	0.27	0
3	PGE	A	507	-	9,9,9	0.34	0	8,8,8	0.30	0
5	N2U	A	509[C]	-	31,34,34	5.32	13 (41%)	40,51,51	2.13	11 (27%)
3	PGE	A	504	-	9,9,9	0.31	0	8,8,8	0.32	0
2	PG4	A	505	-	12,12,12	0.16	0	11,11,11	0.53	0
4	N39	A	508[A]	-	42,45,45	5.91	16 (38%)	54,69,69	3.81	19 (35%)
4	N39	A	508[B]	-	42,45,45	5.83	16 (38%)	54,69,69	3.67	19 (35%)
3	PGE	A	502	-	9,9,9	0.33	0	8,8,8	0.25	0
3	PGE	A	506	-	9,9,9	0.35	0	8,8,8	0.31	0
2	PG4	A	501	-	12,12,12	0.14	0	11,11,11	0.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PGE	A	503	-	-	3/7/7/7	-
3	PGE	A	507	-	-	4/7/7/7	-
5	N2U	A	509[C]	-	-	8/25/45/45	0/3/3/3
3	PGE	A	504	-	-	4/7/7/7	-
2	PG4	A	505	-	-	5/10/10/10	-
4	N39	A	508[A]	-	-	19/37/62/62	0/4/4/4
4	N39	A	508[B]	-	-	19/37/62/62	0/4/4/4
3	PGE	A	502	-	-	5/7/7/7	-
3	PGE	A	506	-	-	4/7/7/7	-
2	PG4	A	501	-	-	1/10/10/10	-

All (45) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	508[A]	N39	C24-N32	30.59	1.71	1.48
4	A	508[B]	N39	C24-N32	30.09	1.70	1.48
5	A	509[C]	N2U	C24-N32	20.51	1.66	1.48
5	A	509[C]	N2U	C17-C18	17.92	1.59	1.33
4	A	508[A]	N39	C17-C18	13.41	1.52	1.33
4	A	508[B]	N39	C17-C18	13.16	1.52	1.33
4	A	508[B]	N39	C24-C18	-8.85	1.43	1.51
4	A	508[A]	N39	C24-C18	-7.78	1.44	1.51
4	A	508[A]	N39	S33-N32	6.83	1.76	1.64
4	A	508[A]	N39	C35-S33	6.49	1.85	1.76
4	A	508[B]	N39	C35-S33	6.21	1.84	1.76
4	A	508[B]	N39	S33-N32	5.95	1.75	1.64
4	A	508[B]	N39	O26-C25	5.39	1.44	1.33
4	A	508[A]	N39	O26-C25	5.34	1.44	1.33
5	A	509[C]	N2U	C16-C17	-4.79	1.40	1.50
5	A	509[C]	N2U	O26-C27	-4.64	1.40	1.48
4	A	508[A]	N39	O20-C19	4.61	1.42	1.33
4	A	508[B]	N39	O20-C19	4.51	1.42	1.33
4	A	508[B]	N39	O13-C12	4.46	1.41	1.33
4	A	508[B]	N39	C16-C17	-4.46	1.41	1.50
4	A	508[A]	N39	O13-C12	4.41	1.41	1.33
4	A	508[A]	N39	C16-C17	-4.27	1.41	1.50
4	A	508[A]	N39	C19-C18	3.91	1.55	1.48
4	A	508[A]	N39	C09-C10	3.75	1.40	1.35
5	A	509[C]	N2U	O13-C12	3.72	1.40	1.33
4	A	508[B]	N39	O34-S33	3.70	1.47	1.43
4	A	508[B]	N39	C19-C18	3.66	1.55	1.48

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	509[C]	N2U	O20-C19	3.57	1.40	1.33
4	A	508[A]	N39	O42-S33	3.54	1.47	1.43
4	A	508[A]	N39	C24-C25	-3.45	1.48	1.53
4	A	508[B]	N39	C09-C10	3.41	1.40	1.35
4	A	508[A]	N39	O34-S33	3.38	1.47	1.43
4	A	508[B]	N39	C24-C25	-3.34	1.48	1.53
4	A	508[B]	N39	O26-C27	-3.25	1.42	1.48
4	A	508[B]	N39	O42-S33	3.23	1.47	1.43
4	A	508[A]	N39	O26-C27	-3.19	1.42	1.48
5	A	509[C]	N2U	O26-C25	2.99	1.39	1.33
5	A	509[C]	N2U	C08-C03	-2.73	1.37	1.41
4	A	508[A]	N39	C04-C03	2.53	1.43	1.39
5	A	509[C]	N2U	C09-C10	2.51	1.38	1.35
5	A	509[C]	N2U	O13-C14	-2.46	1.39	1.45
5	A	509[C]	N2U	C24-C18	-2.43	1.46	1.51
5	A	509[C]	N2U	C19-C18	2.38	1.52	1.48
4	A	508[B]	N39	C04-C03	2.35	1.43	1.39
5	A	509[C]	N2U	O20-C21	-2.12	1.40	1.46

All (49) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	508[A]	N39	O42-S33-O34	-12.06	100.77	119.59
4	A	508[A]	N39	C30-C27-C28	11.63	140.15	111.13
4	A	508[B]	N39	C30-C27-C28	11.53	139.89	111.13
4	A	508[B]	N39	O42-S33-O34	-11.06	102.33	119.59
4	A	508[A]	N39	O26-C25-C24	10.20	120.04	109.25
4	A	508[B]	N39	O26-C25-C24	10.15	119.99	109.25
4	A	508[A]	N39	O26-C27-C29	8.52	140.83	107.21
4	A	508[B]	N39	O26-C27-C29	8.35	140.15	107.21
4	A	508[A]	N39	O13-C12-C11	7.97	119.81	110.82
5	A	509[C]	N2U	O26-C25-C24	7.19	117.50	109.52
4	A	508[B]	N39	O13-C12-C11	7.04	118.76	110.82
4	A	508[B]	N39	C27-O26-C25	-6.41	113.36	121.66
4	A	508[A]	N39	C30-C27-C29	-6.04	96.07	111.13
4	A	508[A]	N39	C27-O26-C25	-5.93	113.99	121.66
4	A	508[B]	N39	C30-C27-C29	-5.92	96.36	111.13
4	A	508[A]	N39	C29-C27-C28	-5.58	97.22	111.13
4	A	508[B]	N39	C29-C27-C28	-5.53	97.33	111.13
5	A	509[C]	N2U	O20-C19-C18	5.14	119.59	112.15
4	A	508[A]	N39	C11-C10-N02	5.12	127.17	123.89
4	A	508[A]	N39	O20-C19-C18	4.99	119.39	112.15

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	508[B]	N39	O20-C19-C18	4.68	118.93	112.15
5	A	509[C]	N2U	O23-C19-C18	-4.15	118.99	124.32
4	A	508[A]	N39	C25-C24-C18	-3.99	106.70	111.88
4	A	508[B]	N39	C25-C24-C18	-3.88	106.83	111.88
4	A	508[A]	N39	O23-C19-C18	-3.41	119.94	124.32
4	A	508[B]	N39	O23-C19-C18	-3.40	119.95	124.32
4	A	508[B]	N39	C11-C10-N02	3.34	126.03	123.89
4	A	508[B]	N39	O26-C25-O31	-3.27	119.71	125.69
5	A	509[C]	N2U	C16-C11-N32	3.26	112.06	108.01
4	A	508[A]	N39	O26-C25-O31	-3.20	119.84	125.69
5	A	509[C]	N2U	C25-C24-C18	-3.02	108.03	111.89
4	A	508[B]	N39	O26-C27-C30	-2.99	95.39	107.21
4	A	508[A]	N39	O31-C25-C24	-2.92	120.13	124.27
4	A	508[A]	N39	O26-C27-C30	-2.88	95.83	107.21
4	A	508[B]	N39	O31-C25-C24	-2.81	120.30	124.27
4	A	508[B]	N39	O42-S33-C35	2.69	111.45	108.10
5	A	509[C]	N2U	C11-C10-N02	2.67	125.91	123.79
5	A	509[C]	N2U	O26-C25-O31	-2.62	120.90	125.69
5	A	509[C]	N2U	O13-C12-O15	-2.60	119.49	123.95
4	A	508[A]	N39	O26-C27-C28	-2.56	97.11	107.21
4	A	508[A]	N39	O13-C12-O15	-2.53	119.60	123.95
5	A	509[C]	N2U	O26-C27-C29	2.47	116.94	107.21
5	A	509[C]	N2U	O15-C12-C11	-2.44	119.68	123.87
4	A	508[B]	N39	O26-C27-C28	-2.40	97.74	107.21
4	A	508[A]	N39	O15-C12-C11	-2.34	120.58	123.79
4	A	508[B]	N39	O15-C12-C11	-2.14	120.86	123.79
4	A	508[A]	N39	C35-S33-N32	2.11	112.42	105.09
4	A	508[B]	N39	O13-C12-O15	-2.07	120.38	123.95
5	A	509[C]	N2U	O31-C25-C24	-2.07	121.60	124.31

There are no chirality outliers.

All (72) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	508[B]	N39	C10-C11-C12-O13
4	A	508[B]	N39	C10-C11-C12-O15
4	A	508[B]	N39	C18-C24-C25-O26
5	A	509[C]	N2U	C16-C11-C12-O13
5	A	509[C]	N2U	C16-C11-C12-O15
5	A	509[C]	N2U	N32-C11-C12-O13
5	A	509[C]	N2U	N32-C11-C12-O15
4	A	508[A]	N39	C30-C27-O26-C25

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	A	508[A]	N39	C41-C35-S33-N32
4	A	508[B]	N39	C18-C19-O20-C21
4	A	508[B]	N39	C29-C27-O26-C25
4	A	508[B]	N39	C30-C27-O26-C25
4	A	508[A]	N39	C36-C35-S33-N32
4	A	508[A]	N39	C29-C27-O26-C25
4	A	508[A]	N39	O31-C25-O26-C27
4	A	508[B]	N39	O31-C25-O26-C27
4	A	508[B]	N39	O23-C19-O20-C21
4	A	508[B]	N39	C28-C27-O26-C25
4	A	508[A]	N39	C24-C25-O26-C27
4	A	508[A]	N39	C28-C27-O26-C25
3	A	502	PGE	O2-C3-C4-O3
3	A	506	PGE	O2-C3-C4-O3
4	A	508[B]	N39	C24-C25-O26-C27
3	A	503	PGE	O1-C1-C2-O2
3	A	507	PGE	O3-C5-C6-O4
2	A	505	PG4	O4-C7-C8-O5
3	A	506	PGE	O1-C1-C2-O2
4	A	508[B]	N39	O15-C12-O13-C14
3	A	502	PGE	O3-C5-C6-O4
3	A	507	PGE	C4-C3-O2-C2
3	A	507	PGE	O2-C3-C4-O3
4	A	508[B]	N39	C41-C35-S33-O34
4	A	508[B]	N39	C36-C35-S33-O34
4	A	508[B]	N39	C11-C12-O13-C14
4	A	508[A]	N39	O15-C12-O13-C14
5	A	509[C]	N2U	O15-C12-O13-C14
2	A	505	PG4	O1-C1-C2-O2
4	A	508[B]	N39	C41-C35-S33-N32
3	A	504	PGE	C6-C5-O3-C4
4	A	508[A]	N39	C11-C12-O13-C14
4	A	508[B]	N39	C36-C35-S33-N32
3	A	506	PGE	C6-C5-O3-C4
2	A	505	PG4	C4-C3-O2-C2
3	A	504	PGE	C1-C2-O2-C3
4	A	508[B]	N39	C18-C24-C25-O31
4	A	508[B]	N39	N32-C11-C12-O13
4	A	508[B]	N39	N32-C11-C12-O15
4	A	508[A]	N39	C36-C35-S33-O42
2	A	501	PG4	O3-C5-C6-O4
5	A	509[C]	N2U	C18-C24-C25-O26

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	A	508[A]	N39	C41-C35-S33-O42
3	A	502	PGE	O1-C1-C2-O2
3	A	506	PGE	C4-C3-O2-C2
3	A	503	PGE	O2-C3-C4-O3
2	A	505	PG4	C6-C5-O3-C4
3	A	504	PGE	O1-C1-C2-O2
4	A	508[A]	N39	C24-N32-S33-O42
3	A	504	PGE	O2-C3-C4-O3
3	A	503	PGE	C4-C3-O2-C2
5	A	509[C]	N2U	C11-C12-O13-C14
2	A	505	PG4	C8-C7-O4-C6
5	A	509[C]	N2U	C18-C24-C25-O31
4	A	508[A]	N39	N32-C24-C25-O26
4	A	508[A]	N39	N32-C24-C25-O31
4	A	508[A]	N39	C41-C35-S33-O34
4	A	508[A]	N39	C24-N32-S33-O34
3	A	502	PGE	C3-C4-O3-C5
4	A	508[A]	N39	C36-C35-S33-O34
4	A	508[A]	N39	C18-C24-C25-O26
4	A	508[A]	N39	C18-C24-C25-O31
3	A	502	PGE	C1-C2-O2-C3
3	A	507	PGE	C6-C5-O3-C4


There are no ring outliers.

4 monomers are involved in 16 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	507	PGE	4	0
5	A	509[C]	N2U	1	0
4	A	508[A]	N39	7	0
4	A	508[B]	N39	4	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



Ligand N39 A 508 (A)





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	369/404 (91%)	0.18	14 (3%)	44 48	11, 29, 51, 78	11 (2%)

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	324	GLY	4.4
1	A	288	ALA	4.0
1	A	384	LEU	3.9
1	A	289	ALA	3.9
1	A	3	ILE	3.7
1	A	383	ASP	3.2
1	A	280	GLN	3.1
1	A	57	GLU	3.0
1	A	323	LYS	2.7
1	A	19	GLU	2.6
1	A	107	ILE	2.5
1	A	96	HIS	2.2
1	A	99	VAL	2.1
1	A	322	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	N39	A	508[A]	42/42	0.34	0.34	18,29,36,36	78
4	N39	A	508[B]	42/42	0.34	0.34	23,30,36,37	78
3	PGE	A	504	10/10	0.70	0.19	53,66,75,76	0
3	PGE	A	503	10/10	0.72	0.20	49,68,85,97	0
3	PGE	A	507	10/10	0.77	0.18	50,64,79,80	0
2	PG4	A	505	13/13	0.82	0.16	35,49,68,80	0
3	PGE	A	506	10/10	0.83	0.14	42,55,71,76	0
3	PGE	A	502	10/10	0.85	0.13	47,58,68,69	0
5	N2U	A	509[C]	32/32	0.91	0.09	23,29,36,36	62
2	PG4	A	501	13/13	0.92	0.10	30,40,56,63	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around N39 A 508 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around N39 A 508 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around N2U A 509 (C):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.