



Full wwPDB NMR Structure Validation Report ⓘ

Mar 5, 2026 – 05:21 PM UTC

PDB ID : 1AC3 / pdb_00001ac3
Title : SOLUTION STRUCTURE OF AN RNA-DNA HYBRID DUPLEX CONTAINING A 3'-THIOFORMACETAL LINKER AND AN RNA A-TRACT, NMR, 8 STRUCTURES
Authors : Cross, C.W.; Rice, J.S.; Gao, X.
Deposited on : 1997-02-11

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR


The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
RNA backbone	8273	777

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	12	 83% 17%
2	B	12	 100%

2 Ensemble composition and analysis ⓘ

This entry contains 8 models. This entry does not contain polypeptide chains, therefore identification of well-defined residues and clustering analysis are not possible. All residues are included in the validation scores.

3 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 770 atoms, of which 274 are hydrogens and 0 are deuteriums.

- Molecule 1 is a DNA chain called ANTISENSE HYBRID DUPLEX.

Mol	Chain	Residues	Atoms							Trace
1	A	12	Total	C	H	N	O	P	S	0
			379	117	140	40	71	10	1	

- Molecule 2 is a RNA chain called DNA (5'-D(*(TCP)P*TP*GP*CP*GP*C)-3').


Mol	Chain	Residues	Atoms						Trace
2	B	12	Total	C	H	N	O	P	0
			391	116	134	52	78	11	

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: ANTISENSE HYBRID DUPLEX



- Molecule 2: DNA (5'-D(*(TCP)P*TP*GP*CP*GP*C)-3')




There are no outlier residues in this chain.

4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

- Molecule 1: ANTISENSE HYBRID DUPLEX



- Molecule 2: DNA (5'-D(*(TCP)P*TP*GP*CP*GP*C)-3')




There are no outlier residues in this chain.

4.2.2 Score per residue for model 2

- Molecule 1: ANTISENSE HYBRID DUPLEX

Chain A:  83% 17%



- Molecule 2: DNA (5'-D(*(TCP)P*TP*GP*CP*GP*C)-3')


Chain B:  92% 8%



4.2.3 Score per residue for model 3

- Molecule 1: ANTISENSE HYBRID DUPLEX

Chain A:  83% 17%



- Molecule 2: DNA (5'-D(*(TCP)P*TP*GP*CP*GP*C)-3')


Chain B:  92% 8%




4.2.4 Score per residue for model 4

- Molecule 1: ANTISENSE HYBRID DUPLEX

Chain A:  83% 17%



- Molecule 2: DNA (5'-D(*(TCP)P*TP*GP*CP*GP*C)-3')


Chain B:  100%

There are no outlier residues in this chain.

4.2.5 Score per residue for model 5

- Molecule 1: ANTISENSE HYBRID DUPLEX

Chain A:  83% 17%



- Molecule 2: DNA (5'-D(*(TCP)P*TP*GP*CP*GP*C)-3')


Chain B:  100%

There are no outlier residues in this chain.

4.2.6 Score per residue for model 6

- Molecule 1: ANTISENSE HYBRID DUPLEX

Chain A:  83% 17%



- Molecule 2: DNA (5'-D(*(TCP)P*TP*GP*CP*GP*C)-3')


Chain B:  92% 8%



4.2.7 Score per residue for model 7


- Molecule 1: ANTISENSE HYBRID DUPLEX

Chain A:  83% 17%



- Molecule 2: DNA (5'-D(*(TCP)P*TP*GP*CP*GP*C)-3')


Chain B:  92% 8%




4.2.8 Score per residue for model 8

- Molecule 1: ANTISENSE HYBRID DUPLEX

Chain A:  83% 17%



- Molecule 2: DNA (5'-D(*(TCP)P*TP*GP*CP*GP*C')-3')

Chain B:  100%

There are no outlier residues in this chain.

5 Refinement protocol and experimental data overview

The models were refined using the following method: *SIMULATED ANNEALING, MOLECULAR DYNAMICS SIMULATIONS AND RELAXATION MATRIX REFINEMENT*.

Of the 10 calculated structures, 8 were deposited, based on the following criterion: *IDEAL GEOMETRY, NOE VIOLATIONS, COORDINATE RMSD*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	refinement	
X-PLOR	structure solution	

No chemical shift data was provided.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TCP, TSP

There are no covalent bond-length or bond-angle outliers.

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
2	B	0.0±0.0	0.5±0.5
All	All	0	4

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
2	B	21	C	Sidechain	3
2	B	15	G	Sidechain	1

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
All	All	3968	2192	2192	-

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is -.

There are no clashes.

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

6.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

6.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers	Suiteness
2	B	11/12 (92%)	0±0 (0±0%)	0±0 (0±0%)	0.35±0.00
All	All	88/96 (92%)	0 (0%)	0 (0%)	0.35

The overall RNA backbone suiteness is 0.39.

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
1	TCP	A	7	2,1	19,19,19	0.53±0.01	0±0 (0±0%)
1	TSP	A	6	2,1	17,21,22	0.56±0.01	0±0 (0±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is

considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
1	TCP	A	7	2,1	27,27,27	1.06±0.03	2±1 (6±2%)
1	TSP	A	6	2,1	21,30,33	1.11±0.05	2±0 (8±2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TSP	A	6	2,1	-	0±0,7,21,22	0±0,2,2,2
1	TCP	A	7	2,1	-	0±0,7,19,19	0±0,2,2,2

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	6	TSP	C6-C5-C4	3.06	120.54	118.02	4	8
1	A	7	TCP	C6-C5-C4	2.94	120.45	118.02	3	8
1	A	7	TCP	O4'-C1'-N1	2.50	112.30	107.86	4	5
1	A	6	TSP	O4'-C1'-N1	2.41	112.15	107.86	6	6
1	A	7	TCP	C5M-C5-C6	2.11	119.99	122.85	7	2

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided