



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 06:26 AM UTC

PDB ID : 2AC0 / pdb_00002ac0
Title : Structural Basis of DNA Recognition by p53 Tetramers (complex I)
Authors : Kitayner, M.; Rozenberg, H.; Kessler, N.; Rabinovich, D.; Shakked, Z.
Deposited on : 2005-07-18
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	12	33% 50% 17%
1	F	12	50% 33% 17%
1	G	12	75% 25%
1	H	12	50% 42% 8%
2	A	200	2% 90% 8%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	B	200	2% 88% 9% ..
2	C	200	% 88% 8% ..
2	D	200	3% 84% 12% ..

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8247 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(*CP*GP*GP*GP*CP*AP*TP*GP*CP*CP*CP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	E	12	Total	C	N	O	P	0	0	0
			243	115	47	70	11			
1	F	12	Total	C	N	O	P	0	0	0
			243	115	47	70	11			
1	G	12	Total	C	N	O	P	0	0	0
			243	115	47	70	11			
1	H	12	Total	C	N	O	P	0	0	0
			243	115	47	70	11			

- Molecule 2 is a protein called Cellular tumor antigen p53.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	199	Total	C	N	O	S	0	2	1
			1557	957	289	294	17			
2	B	198	Total	C	N	O	S	0	4	0
			1557	957	287	296	17			
2	C	195	Total	C	N	O	S	0	5	0
			1544	954	282	290	18			
2	D	198	Total	C	N	O	S	0	2	1
			1559	961	290	291	17			

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		
3	B	1	Total	Zn	0	0
			1	1		
3	C	1	Total	Zn	0	0
			1	1		
3	D	1	Total	Zn	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	E	51	Total 51	O 51	0	0
4	F	40	Total 40	O 40	0	0
4	G	31	Total 31	O 31	0	0
4	H	32	Total 32	O 32	0	0
4	A	255	Total 255	O 255	0	0
4	B	195	Total 195	O 195	0	0
4	C	253	Total 253	O 253	0	0
4	D	197	Total 197	O 197	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 5'-D(*CP*GP*GP*GP*CP*AP*TP*GP*CP*CP*CP*G)-3'

Chain E: 



- Molecule 1: 5'-D(*CP*GP*GP*GP*CP*AP*TP*GP*CP*CP*CP*G)-3'

Chain F: 



- Molecule 1: 5'-D(*CP*GP*GP*GP*CP*AP*TP*GP*CP*CP*CP*G)-3'

Chain G: 



- Molecule 1: 5'-D(*CP*GP*GP*GP*CP*AP*TP*GP*CP*CP*CP*G)-3'

Chain H: 





- Molecule 2: Cellular tumor antigen p53


Chain A: 





- Molecule 2: Cellular tumor antigen p53



• Molecule 2: Cellular tumor antigen p53



• Molecule 2: Cellular tumor antigen p53



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	54.44Å 58.15Å 77.54Å 82.93° 87.99° 73.60°	Depositor
Resolution (Å)	33.52 – 1.80 33.52 – 1.80	Depositor EDS
% Data completeness (in resolution range)	97.8 (33.52-1.80) 92.1 (33.52-1.80)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.86 (at 1.79Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.153 , 0.217 0.160 , 0.158	Depositor DCC
R_{free} test set	4002 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	18.5	Xtriage
Anisotropy	0.237	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 61.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8247	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	E	0.73	0/272	1.55	7/418 (1.7%)
1	F	0.63	0/272	1.47	5/418 (1.2%)
1	G	0.55	0/272	1.35	2/418 (0.5%)
1	H	0.50	0/272	1.37	2/418 (0.5%)
2	A	0.98	1/1599 (0.1%)	1.00	1/2168 (0.0%)
2	B	0.90	1/1607 (0.1%)	1.01	1/2179 (0.0%)
2	C	0.98	1/1599 (0.1%)	1.01	0/2169
2	D	0.85	0/1601	0.94	1/2168 (0.0%)
All	All	0.89	3/7494 (0.0%)	1.07	19/10356 (0.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	190	PRO	CA-C	7.26	1.55	1.51
2	B	190	PRO	CA-C	5.46	1.54	1.51
2	C	253	THR	N-CA	5.40	1.52	1.46

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	5	DC	P-O5'-C5'	-10.33	104.50	120.00
1	F	5	DC	P-O5'-C5'	-7.65	108.52	120.00
1	F	5	DC	C5'-C4'-C3'	-7.25	104.02	114.90
1	F	10	DC	O4'-C1'-N1	-6.58	98.53	108.40
1	E	5	DC	C4'-C3'-O3'	-6.51	100.24	110.00
1	G	6	DA	O4'-C1'-N9	-6.49	98.66	108.40
1	G	6	DA	N9-C1'-C2'	6.27	122.91	113.50
2	A	97	VAL	N-CA-C	6.06	114.64	107.73
1	E	3	DG	O4'-C1'-N9	-5.86	99.61	108.40
1	E	6	DA	O4'-C1'-N9	-5.78	99.73	108.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	6	DA	O4'-C1'-N9	-5.61	99.98	108.40
2	B	97	VAL	N-CA-C	5.56	114.61	108.05
1	F	6	DA	O4'-C1'-N9	-5.45	100.23	108.40
1	E	6	DA	N9-C1'-C2'	5.41	121.61	113.50
1	F	7	DT	C1'-O4'-C4'	-5.33	101.70	109.70
1	E	5	DC	C1'-O4'-C4'	-5.32	101.73	109.70
1	E	6	DA	C4'-C3'-O3'	-5.14	102.28	110.00
1	H	2	DG	O4'-C1'-N9	-5.12	100.71	108.40
2	D	97	VAL	N-CA-C	5.03	112.84	107.76

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	243	0	135	8	0
1	F	243	0	135	5	0
1	G	243	0	135	2	0
1	H	243	0	135	6	0
2	A	1557	0	1511	11	0
2	B	1557	0	1503	25	0
2	C	1544	0	1505	16	0
2	D	1559	0	1526	22	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	255	0	0	3	2
4	B	195	0	0	4	0
4	C	253	0	0	4	2
4	D	197	0	0	6	0
4	E	51	0	0	2	0
4	F	40	0	0	0	0
4	G	31	0	0	0	0
4	H	32	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	8247	0	6585	83	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (83) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:102:THR:HG23	2:B:268:ASN:HD21	1.18	1.02
2:B:102:THR:HG23	2:B:268:ASN:ND2	1.84	0.92
2:D:207:ASP:HB3	4:D:426:HOH:O	1.82	0.78
1:E:11:DC:H5	4:E:919:HOH:O	1.67	0.78
2:C:256[A]:THR:HG22	4:C:317:HOH:O	1.86	0.76
2:D:174:ARG:HD3	4:D:424:HOH:O	1.88	0.74
1:E:1:DC:H5'	1:G:12:DG:H2'	1.68	0.73
1:E:12:DG:OP1	4:E:982:HOH:O	2.08	0.71
4:A:435:HOH:O	2:B:243:MET:HE2	1.91	0.70
2:D:166:SER:HA	2:D:169:MET:HG3	1.73	0.69
2:B:119:ALA:O	2:B:279:GLY:HA3	1.92	0.69
1:F:1:DC:O5'	2:B:283:ARG:HD3	1.94	0.67
1:F:5:DC:H2''	1:F:6:DA:C8	2.30	0.66
2:C:286:GLU:OE1	4:C:475:HOH:O	2.13	0.65
2:C:124:CYS:SG	2:C:133[B]:MET:SD	2.94	0.65
2:C:178:HIS:CD2	2:D:243:MET:HG3	2.33	0.63
2:B:247:ASN:ND2	4:B:450:HOH:O	2.31	0.63
2:C:123[B]:THR:CG2	2:C:139:LYS:HB3	2.31	0.61
2:A:100:GLN:HG2	4:A:377:HOH:O	2.03	0.59
2:B:110:ARG:HB2	4:B:383:HOH:O	2.01	0.59
1:E:5:DC:H2''	1:E:6:DA:C8	2.40	0.57
1:H:1:DC:H5''	2:C:119:ALA:HB1	1.86	0.57
2:A:268:ASN:HD22	2:A:269:SER:N	2.02	0.56
1:H:11:DC:H2''	1:H:12:DG:C8	2.41	0.56
2:D:288:ASN:ND2	4:D:463:HOH:O	2.39	0.55
2:C:123[B]:THR:HG23	2:C:139:LYS:HB3	1.88	0.55
2:A:178:HIS:CD2	2:B:243:MET:HE3	2.41	0.55
2:A:175:ARG:HD3	2:A:191:PRO:O	2.07	0.55
2:B:192:GLN:HB3	4:B:341:HOH:O	2.07	0.54
2:C:204:GLU:HB3	2:C:217[A]:VAL:HG13	1.90	0.54
1:H:11:DC:C2'	1:H:12:DG:C8	2.91	0.53
2:C:123[B]:THR:HG23	2:C:139:LYS:CB	2.38	0.53
2:D:259:ASP:HA	4:D:354:HOH:O	2.09	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:1:DC:H2'	1:H:2:DG:C8	2.44	0.52
2:A:287:GLU:O	2:A:290:ARG:HG2	2.10	0.52
2:B:243:MET:HA	2:B:247:ASN:HD22	1.75	0.51
2:D:158:ARG:NH1	2:D:217:VAL:CG2	2.73	0.51
1:E:1:DC:H3'	2:A:121:SER:HB2	1.93	0.51
1:G:11:DC:H2'	1:G:12:DG:C8	2.46	0.51
2:B:198:GLU:CG	2:B:233:HIS:HB3	2.41	0.51
2:B:198:GLU:CG	2:B:233:HIS:CB	2.89	0.50
2:B:192:GLN:H	2:B:192:GLN:CD	2.20	0.50
2:D:158:ARG:CZ	2:D:217:VAL:HG21	2.41	0.50
2:C:177:PRO:HB2	2:D:177:PRO:HB2	1.94	0.50
2:A:198:GLU:CD	4:A:488:HOH:O	2.54	0.49
2:D:102:THR:HG23	2:D:268:ASN:HD21	1.76	0.49
2:D:99:SER:OG	2:D:101:LYS:HG2	2.12	0.49
2:B:174:ARG:HD3	4:B:426:HOH:O	2.13	0.49
2:C:158:ARG:HB3	2:C:256[A]:THR:HG23	1.95	0.49
2:C:119:ALA:O	2:C:279:GLY:HA3	2.13	0.49
2:B:198:GLU:HG2	2:B:233:HIS:HB3	1.95	0.49
2:A:166:SER:HA	2:A:169:MET:HE2	1.94	0.48
1:E:1:DC:H2''	1:E:2:DG:C8	2.48	0.48
2:B:167:GLN:HB3	4:C:380:HOH:O	2.13	0.48
2:C:178:HIS:CD2	2:D:243:MET:CG	2.97	0.48
1:F:1:DC:O5'	2:B:283:ARG:CD	2.63	0.47
2:B:198:GLU:HG3	2:B:233:HIS:HB2	1.97	0.47
1:F:12:DG:H2''	1:H:1:DC:H5'	1.98	0.46
2:D:259:ASP:OD2	2:D:263:ASN:HB2	2.16	0.46
2:B:198:GLU:HG2	2:B:233:HIS:CB	2.45	0.45
2:D:102:THR:HG23	2:D:268:ASN:ND2	2.32	0.45
1:H:10:DC:H2''	1:H:11:DC:C6	2.52	0.45
2:C:158:ARG:NH1	2:C:217[B]:VAL:CG2	2.80	0.45
2:D:158:ARG:NH1	2:D:217:VAL:HG21	2.30	0.45
2:C:209:ARG:HD3	4:C:428:HOH:O	2.17	0.45
2:D:119:ALA:O	2:D:120:LYS:C	2.60	0.44
2:B:198:GLU:CG	2:B:233:HIS:HB2	2.47	0.44
2:C:158:ARG:CZ	2:C:217[B]:VAL:HG21	2.47	0.44
2:A:178:HIS:HD2	2:B:243:MET:HE3	1.82	0.43
2:D:118:THR:HG22	2:D:282:ARG:HD3	1.99	0.43
2:B:175:ARG:HD3	2:B:191:PRO:O	2.19	0.43
2:D:202:ARG:NH1	2:D:221:GLU:OE2	2.53	0.42
2:B:151:PRO:HD2	2:B:220:TYR:CE2	2.54	0.42
1:E:1:DC:C2'	1:E:2:DG:C8	3.03	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:145:LEU:HD11	2:D:232:ILE:HD11	2.02	0.41
2:D:97:VAL:HG21	2:D:169:MET:HB3	2.03	0.41
1:F:1:DC:C5'	2:B:283:ARG:HD3	2.50	0.41
2:B:267:ARG:O	2:B:268:ASN:ND2	2.46	0.41
2:A:100:GLN:HG2	2:A:100:GLN:H	1.30	0.41
2:D:152:PRO:HG2	4:D:354:HOH:O	2.21	0.41
1:E:9:DC:O2	1:E:12:DG:H2'	2.21	0.40
2:D:128:PRO:HD2	4:D:407:HOH:O	2.21	0.40
2:A:204:GLU:HB3	2:A:217:VAL:HG13	2.03	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:344:HOH:O	4:C:537:HOH:O[1_665]	2.04	0.16
4:A:516:HOH:O	4:C:535:HOH:O[1_655]	2.12	0.08

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	199/200 (100%)	196 (98%)	3 (2%)	0	100	100
2	B	199/200 (100%)	196 (98%)	3 (2%)	0	100	100
2	C	198/200 (99%)	196 (99%)	2 (1%)	0	100	100
2	D	198/200 (99%)	192 (97%)	5 (2%)	1 (0%)	24	14
All	All	794/800 (99%)	780 (98%)	13 (2%)	1 (0%)	48	34

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	290	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	179/179 (100%)	174 (97%)	5 (3%)	38	26
2	B	180/179 (101%)	172 (96%)	8 (4%)	25	13
2	C	179/179 (100%)	171 (96%)	8 (4%)	24	12
2	D	179/179 (100%)	172 (96%)	7 (4%)	28	16
All	All	717/716 (100%)	689 (96%)	28 (4%)	30	16

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	100	GLN
2	A	224	GLU
2	A	225	VAL
2	A	264	LEU
2	A	268	ASN
2	B	110	ARG
2	B	149	SER
2	B	192	GLN
2	B	224	GLU
2	B	243	MET
2	B	268	ASN
2	B	283	ARG
2	B	290	ARG
2	C	169	MET
2	C	180	GLU
2	C	217[A]	VAL
2	C	217[B]	VAL
2	C	256[A]	THR
2	C	256[B]	THR
2	C	283	ARG
2	C	287	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	D	96	SER
2	D	97	VAL
2	D	174	ARG
2	D	243	MET
2	D	264	LEU
2	D	268	ASN
2	D	290	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	239	ASN
2	A	268	ASN
2	A	288	ASN
2	B	100	GLN
2	B	131	ASN
2	B	136	GLN
2	B	239	ASN
2	B	247	ASN
2	B	268	ASN
2	C	104	GLN
2	D	100	GLN
2	D	168	HIS
2	D	247	ASN
2	D	263	ASN
2	D	268	ASN
2	D	288	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	E	12/12 (100%)	-0.28	0 100 100	19, 25, 42, 49	0
1	F	12/12 (100%)	-0.39	0 100 100	18, 24, 45, 50	0
1	G	12/12 (100%)	-0.49	0 100 100	19, 27, 44, 46	0
1	H	12/12 (100%)	-0.15	0 100 100	20, 32, 64, 66	0
2	A	199/200 (99%)	-0.43	3 (1%) 72 72	11, 19, 39, 54	2 (1%)
2	B	198/200 (99%)	-0.06	3 (1%) 72 72	13, 26, 48, 64	4 (2%)
2	C	195/200 (97%)	-0.39	2 (1%) 79 80	9, 20, 39, 57	5 (2%)
2	D	198/200 (99%)	0.09	6 (3%) 52 52	15, 28, 55, 77	2 (1%)
All	All	838/848 (98%)	-0.21	14 (1%) 69 69	9, 23, 48, 77	13 (1%)

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	291	LYS	3.7
2	A	225	VAL	3.5
2	D	293	GLY	3.3
2	B	225	VAL	3.3
2	C	289	LEU	2.9
2	D	226	GLY	2.8
2	D	289	LEU	2.7
2	D	225	VAL	2.6
2	C	226	GLY	2.4
2	D	228	ASP	2.3
2	A	96	SER	2.2
2	A	183	SER	2.1
2	D	227	SER	2.1
2	B	106	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	ZN	A	1	1/1	0.99	0.03	17,17,17,17	0
3	ZN	B	1	1/1	1.00	0.04	20,20,20,20	0
3	ZN	C	1	1/1	1.00	0.04	17,17,17,17	0
3	ZN	D	1	1/1	1.00	0.02	19,19,19,19	0

6.5 Other polymers [i](#)

There are no such residues in this entry.