



# wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 25, 2026 – 04:32 AM EDT

PDB ID : 2ACX / pdb\_00002acx  
Title : Crystal Structure of G protein coupled receptor kinase 6 bound to AMPPNP  
Authors : Lodowski, D.T.; Tesmer, V.M.; Benovic, J.L.; Tesmer, J.J.  
Deposited on : 2005-07-19  
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	576	 2% 65% 17% 14%
1	B	576	 3% 68% 15% 15%

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8106 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called G protein-coupled receptor kinase 6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	495	4011	2548	714	721	28	0	0	0
1	B	492	3984	2535	705	717	27	0	0	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	60	ASN	GLN	SEE REMARK 999	UNP P43250
A	104	ASN	GLN	SEE REMARK 999	UNP P43250
A	561	SER	CYS	engineered mutation	UNP P43250
A	562	SER	CYS	engineered mutation	UNP P43250
A	565	SER	CYS	engineered mutation	UNP P43250
B	60	ASN	GLN	SEE REMARK 999	UNP P43250
B	104	ASN	GLN	SEE REMARK 999	UNP P43250
B	561	SER	CYS	engineered mutation	UNP P43250
B	562	SER	CYS	engineered mutation	UNP P43250
B	565	SER	CYS	engineered mutation	UNP P43250

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		
2	B	1	Total	Mg	0	0
			1	1		

- Molecule 3 is PHOSPHATE ION (CCD ID: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O P 5 4 1	0	0
3	B	1	Total O P 5 4 1	0	0

- Molecule 4 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (CCD ID: ANP) (formula: C<sub>10</sub>H<sub>17</sub>N<sub>6</sub>O<sub>12</sub>P<sub>3</sub>).



*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
4	B	1	31	10	6	12	3	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	30	Total	O	0	0
			30	30		
5	B	7	Total	O	0	0
			7	7		



GLY  
GLN  
PRO  
ALA  
PRO  
LYS  
LYS  
GLY  
LEU  
LEU  
GLN  
ARG  
LEU  
PHE  
SER  
ARG  
GLN  
ASP  
SER  
SER  
SER  
GLY  
ASN  
SER  
SER  
ASP  
SER  
GLU  
GLU  
LEU  
PRO  
THR  
ARG  
LEU

## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	120.16Å 59.27Å 221.09Å 90.00° 102.58° 90.00°	Depositor
Resolution (Å)	30.00 – 2.60 30.00 – 2.60	Depositor EDS
% Data completeness (in resolution range)	97.4 (30.00-2.60) 97.2 (30.00-2.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.52 (at 2.61Å)	Xtrriage
Refinement program	REFMAC 5.2.0003	Depositor
R, $R_{free}$	0.202 , 0.243 0.208 , 0.244	Depositor DCC
$R_{free}$ test set	2317 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	61.3	Xtrriage
Anisotropy	0.088	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 76.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	8106	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	69.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.99% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ANP, PO4, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	1.26	4/4101 (0.1%)	1.17	11/5521 (0.2%)
1	B	0.90	0/4074	1.02	2/5488 (0.0%)
All	All	1.10	4/8175 (0.0%)	1.10	13/11009 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	427	ASP	N-CA	5.63	1.52	1.46
1	A	362	TYR	CA-CB	5.57	1.63	1.53
1	A	308	VAL	CA-CB	-5.53	1.47	1.54
1	A	347	GLY	C-O	-5.16	1.18	1.23

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	438	SER	CB-CA-C	-7.70	106.63	115.79
1	A	168	ASN	N-CA-C	-7.31	103.25	111.07
1	A	404	VAL	CA-C-O	6.60	123.60	119.51
1	A	355	GLU	CB-CA-C	-6.48	99.99	110.74
1	A	349	VAL	N-CA-C	6.05	116.71	110.36

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4011	0	4008	67	0
1	B	3984	0	3974	57	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
4	A	31	0	13	0	0
4	B	31	0	13	1	0
5	A	30	0	0	2	0
5	B	7	0	0	0	0
All	All	8106	0	8008	119	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 119 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:132:ARG:NH2	1:A:138:CYS:SG	2.42	0.93
1:A:319:LEU:CD2	1:A:325:ILE:HG22	2.00	0.92
1:B:133:LEU:HB2	1:B:141:LEU:HD21	1.57	0.86
1:B:319:LEU:HD22	1:B:325:ILE:HG22	1.58	0.84
1:A:325:ILE:C	1:A:325:ILE:HD12	2.11	0.76

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	489/576 (85%)	458 (94%)	24 (5%)	7 (1%)	<b>9</b> <b>19</b>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	B	482/576 (84%)	459 (95%)	17 (4%)	6 (1%)	10 23
All	All	971/1152 (84%)	917 (94%)	41 (4%)	13 (1%)	9 21

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	141	LEU
1	A	449	LYS
1	A	526	VAL
1	B	141	LEU
1	B	196	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	432/503 (86%)	399 (92%)	33 (8%)	12 27
1	B	430/503 (86%)	401 (93%)	29 (7%)	15 33
All	All	862/1006 (86%)	800 (93%)	62 (7%)	13 30

5 of 62 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	473	TYR
1	B	395	GLU
1	B	60	ASN
1	B	394	ARG
1	B	472	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	387	GLN
1	B	335	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	131	GLN
1	B	322	HIS
1	B	108	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	PO4	A	579	-	4,4,4	0.85	0	6,6,6	1.65	1 (16%)
4	ANP	A	577	2	33,33,33	2.25	9 (27%)	45,52,52	2.31	15 (33%)
3	PO4	B	579	-	4,4,4	0.74	0	6,6,6	1.56	1 (16%)
4	ANP	B	577	2	33,33,33	1.97	9 (27%)	45,52,52	2.22	15 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ANP	A	577	2	-	3/18/38/38	0/3/3/3
4	ANP	B	577	2	-	3/18/38/38	0/3/3/3

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	577	ANP	PB-O3A	5.87	1.66	1.59
4	A	577	ANP	PG-O1G	4.59	1.53	1.46
4	A	577	ANP	PG-N3B	4.59	1.75	1.63
4	B	577	ANP	PG-N3B	4.58	1.75	1.63
4	B	577	ANP	PB-N3B	4.19	1.74	1.63

The worst 5 of 32 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	577	ANP	C5-C4-N3	-6.42	117.87	126.72
4	A	577	ANP	N3-C2-N1	-5.55	120.18	128.58
4	B	577	ANP	N3-C4-N9	5.30	136.18	127.17
4	A	577	ANP	C5-C4-N3	-5.01	119.82	126.72
4	A	577	ANP	O1B-PB-N3B	-4.77	104.75	111.77

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	577	ANP	PB-N3B-PG-O1G
4	A	577	ANP	PA-O3A-PB-O2B
4	B	577	ANP	PB-N3B-PG-O1G
4	B	577	ANP	C4'-C5'-O5'-PA
4	B	577	ANP	PG-N3B-PB-O1B


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	577	ANP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	495/576 (85%)	0.07	9 (1%) 67 63	56, 67, 85, 106	0
1	B	492/576 (85%)	0.37	19 (3%) 43 38	58, 68, 84, 111	0
All	All	987/1152 (85%)	0.22	28 (2%) 55 49	56, 68, 85, 111	0

The worst 5 of 28 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	473	TYR	4.4
1	B	472	ILE	3.7
1	B	531	GLY	3.3
1	B	533	VAL	3.0
1	B	471	ALA	2.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.


### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MG	B	578	1/1	0.53	0.24	87,87,87,87	0
2	MG	A	578	1/1	0.87	0.10	92,92,92,92	0
3	PO4	B	579	5/5	0.89	0.17	91,93,95,97	0
3	PO4	A	579	5/5	0.90	0.19	83,83,87,87	0
4	ANP	B	577	31/31	0.93	0.08	60,63,92,93	0
4	ANP	A	577	31/31	0.95	0.07	60,65,103,105	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.