



wwPDB NMR Structure Validation Summary Report ⓘ

Apr 15, 2026 – 01:11 PM UTC

PDB ID : 7ACS / pdb_00007acs
BMRB ID : 34511
Title : The SARS-CoV-2 nucleocapsid phosphoprotein N-terminal domain in complex with 7mer dsRNA
Authors : Veverka, V.
Deposited on : 2020-09-11

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
BMRB Restraints Analysis : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 79%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823
RNA backbone	8273	777

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	140	
2	B	7	
3	C	7	

2 Ensemble composition and analysis

This entry contains 30 models. Model 13 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:10-A:55, A:61-A:136 (122)	0.56	13

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 11 single-model clusters were found.

Cluster number	Models
1	3, 4, 8, 9, 10, 11, 13, 14, 18, 20, 22, 23, 24, 25, 27
2	6, 30
3	7, 26
Single-model clusters	1; 2; 5; 12; 15; 16; 17; 19; 21; 28; 29

3 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2564 atoms, of which 1195 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Nucleoprotein.

Mol	Chain	Residues	Atoms						Trace
1	A	140	Total	C	H	N	O	S	0
			2109	671	1040	195	201	2	

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP P0DTC9
A	2	ALA	-	expression tag	UNP P0DTC9
A	3	MET	-	expression tag	UNP P0DTC9

- Molecule 2 is a RNA chain called RNA (5'-R(P*CP*AP*CP*UP*GP*AP*C)-3').

Mol	Chain	Residues	Atoms						Trace
2	B	7	Total	C	H	N	O	P	0
			226	66	78	26	49	7	

- Molecule 3 is a RNA chain called RNA (5'-R(P*GP*UP*CP*AP*GP*UP*G)-3').


Mol	Chain	Residues	Atoms						Trace
3	C	7	Total	C	H	N	O	P	0
			229	67	77	27	51	7	

4 Residue-property plots [i](#)


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: Nucleoprotein



- Molecule 2: RNA (5'-R(P*CP*AP*CP*UP*GP*AP*C)-3')




- Molecule 3: RNA (5'-R(P*GP*UP*CP*AP*GP*UP*G)-3')



4.2 Residue scores for the representative (medoid) model from the NMR ensemble


The representative model is number 13. Colouring as in section 4.1 above.

- Molecule 1: Nucleoprotein



- Molecule 2: RNA (5'-R(P*CP*AP*CP*UP*GP*AP*C)-3')

Chain B:  43% 29% 29%



- Molecule 3: RNA (5'-R(P*GP*UP*CP*AP*GP*UP*G)-3')

Chain C:  14% 43% 43%



5 Refinement protocol and experimental data overview

The models were refined using the following method: *molecular dynamics*.

Of the 200 calculated structures, 30 were deposited, based on the following criterion: *structures with the least restraint violations*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
HADDOCK	structure calculation	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1665
Number of shifts mapped to atoms	1665
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	79%

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.33±0.01	0±0/984 (0.0± 0.0%)	0.67±0.02	0±0/1341 (0.0± 0.0%)
2	B	1.83±0.00	2±0/164 (1.2± 0.0%)	2.03±0.00	8±0/251 (3.1± 0.2%)
3	C	1.78±0.00	1±0/169 (0.6± 0.0%)	2.07±0.00	8±0/260 (3.2± 0.2%)
All	All	0.95	90/39510 (0.2%)	1.22	482/55560 (0.9%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
2	B	0.0±0.0	3.0±0.0
3	C	0.0±0.0	4.0±0.0
All	All	0	210

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
3	C	8	G	OP3-P	5.69	1.59	1.48	24	30
2	B	1	C	OP3-P	5.66	1.59	1.48	17	30
2	B	7	C	C3'-O3'	5.47	1.50	1.42	6	30

5 of 17 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
3	C	8	G	O4'-C4'-C3'	11.05	115.05	104.00	3	30
3	C	12	G	C4'-C3'-C2'	-8.34	94.26	102.60	3	30
3	C	9	U	O3'-P-O5'	8.28	116.42	104.00	21	30
2	B	6	A	O4'-C4'-C3'	7.57	111.57	104.00	20	30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
3	C	11	A	C1'-C2'-O2'	6.83	118.64	108.40	30	30

There are no chirality outliers.

5 of 7 unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
2	B	2	A	Sidechain	30
2	B	5	G	Sidechain	30
2	B	7	C	Sidechain	30
3	C	10	C	Sidechain	30
3	C	11	A	Sidechain	30

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	955	931	929	10±3
2	B	148	78	73	1±1
3	C	152	77	72	1±1
All	All	37650	32580	32220	325

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

5 of 107 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:14:THR:HG22	3:C:8:G:H5'	0.84	1.48	23	4
1:A:42:ASP:HA	1:A:103:LYS:HE3	0.81	1.52	11	20
1:A:14:THR:HB	1:A:134:GLU:HB3	0.75	1.59	15	4
1:A:44:ILE:HB	1:A:98:ALA:HB1	0.72	1.61	19	14
1:A:54:ILE:HG12	1:A:62:LYS:HB2	0.71	1.61	12	6

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	122/140 (87%)	111±3 (91±2%)	10±3 (8±2%)	1±1 (1±1%)	16	66
All	All	3660/4200 (87%)	3322 (91%)	304 (8%)	34 (1%)	16	66

5 of 7 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	63	ASP	21
1	A	24	LEU	5
1	A	37	ASN	3
1	A	66	PRO	2
1	A	113	ASN	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	97/107 (91%)	85±3 (88±3%)	12±3 (12±3%)	7	49
All	All	2910/3210 (91%)	2558 (88%)	352 (12%)	7	49

5 of 46 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	36	THR	28
1	A	126	THR	25
1	A	38	SER	24
1	A	75	THR	23
1	A	103	LYS	20

6.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers	Suiteness
2	B	7/7 (100%)	0±0 (0±0%)	1±0 (14±0%)	0.48±0.00
3	C	6/7 (86%)	0±0 (0±0%)	0±0 (0±0%)	0.35±0.00
All	All	390/420 (93%)	0 (0%)	30 (8%)	0.41

The overall RNA backbone suiteness is 0.41.

There are no RNA backbone outliers to report.

All unique RNA pucker outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
2	B	1	C	30

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.6 Ligand geometry ⓘ

There are no ligands in this entry.

6.7 Other polymers ⓘ

There are no such molecules in this entry.

6.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 79% for the well-defined parts and 79% for the entire structure.

7.1 Chemical shift list 1

File name: `working_cs.cif`

Chemical shift list name: *starch_output*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1665
Number of shifts mapped to atoms	1665
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	24

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	140	0.17 ± 0.13	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	119	0.03 ± 0.16	None needed (< 0.5 ppm)
$^{13}\text{C}'$	137	0.10 ± 0.15	None needed (< 0.5 ppm)
^{15}N	127	-0.56 ± 0.42	None needed (imprecise)

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 79%, i.e. 1495 atoms were assigned a chemical shift out of a possible 1901. 0 out of 12 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	600/604 (99%)	247/248 (100%)	242/244 (99%)	111/112 (99%)
Sidechain	791/878 (90%)	535/569 (94%)	236/267 (88%)	20/42 (48%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	104/155 (67%)	63/74 (85%)	38/74 (51%)	3/7 (43%)
Sugar	0/154 (0%)	0/84 (0%)	0/70 (0%)	0/0 (—%)
Base	0/110 (0%)	0/68 (0%)	0/24 (0%)	0/18 (0%)
Overall	1495/1901 (79%)	845/1043 (81%)	516/679 (76%)	134/179 (75%)

7.1.4 Statistically unusual chemical shifts ⓘ

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.


List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	72	TYR	CE1	182.76	111.24 – 124.66	48.3
1	A	14	THR	HG1	6.34	0.08 – 2.19	24.6
1	A	125	THR	HG1	5.79	0.08 – 2.19	22.1
1	A	101	THR	HG1	5.78	0.08 – 2.19	22.0
1	A	47	TYR	CE1	129.73	111.24 – 124.66	8.8
1	A	109	ARG	HB3	-0.55	0.43 – 3.11	-8.6
1	A	49	ARG	HB3	-0.51	0.43 – 3.11	-8.5
1	A	47	TYR	HE1	5.00	5.59 – 7.82	-7.6
1	A	47	TYR	HE2	5.00	5.58 – 7.83	-7.5
1	A	24	LEU	HD11	-1.21	-0.61 – 2.12	-7.2
1	A	24	LEU	HD12	-1.21	-0.61 – 2.12	-7.2
1	A	24	LEU	HD13	-1.21	-0.61 – 2.12	-7.2
1	A	33	PRO	HD3	0.96	1.76 – 5.48	-7.1
1	A	24	LEU	HB3	-0.85	-0.26 – 3.31	-6.7
1	A	78	GLU	HB2	0.69	1.00 – 3.05	-6.5
1	A	33	PRO	HG3	-0.10	0.33 – 3.48	-6.3
1	A	31	GLY	HA2	1.75	2.15 – 5.77	-6.1
1	A	24	LEU	HB2	-0.33	-0.07 – 3.30	-5.8
1	A	83	TYR	HA	1.62	1.87 – 7.33	-5.5
1	A	47	TYR	HD1	5.43	5.49 – 8.39	-5.2
1	A	47	TYR	HD2	5.43	5.48 – 8.39	-5.2
1	A	24	LEU	HD21	-0.67	-0.65 – 2.13	-5.1
1	A	24	LEU	HD22	-0.67	-0.65 – 2.13	-5.1
1	A	24	LEU	HD23	-0.67	-0.65 – 2.13	-5.1

7.1.5 Random Coil Index (RCI) plots ⓘ

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication

of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:



8 NMR restraints analysis

8.1 Conformationally restricting restraints

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	17
Intra-residue ($ i-j =0$)	0
Sequential ($ i-j =1$)	0
Medium range ($ i-j >1$ and $ i-j <5$)	0
Long range ($ i-j \geq 5$)	0
Inter-chain	17
Hydrogen bond restraints	0
Disulfide bond restraints	0
Total dihedral-angle restraints	0
Number of unmapped restraints	0
Number of restraints per residue	0.1
Number of long range restraints per residue ¹	0.0

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	0.7	0.2
0.2-0.5 (Medium)	1.3	0.49
>0.5 (Large)	12.4	11.07

8.2.2 Average number of dihedral-angle violations per model [i](#)

Dihedral-angle violations less than 1° are not included in the calculation. There are no dihedral-angle violations

9 Distance violation analysis ⓘ


9.1 Summary of distance violations ⓘ

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Restrains type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
Intra-residue (i-j =0)	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sequential (i-j =1)	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Medium range (i-j >1 & i-j <5)	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Long range (i-j ≥5)	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Inter-chain	17	100.0	17	100.0	100.0	8	47.1	47.1
Backbone-Backbone	17	100.0	17	100.0	100.0	8	47.1	47.1
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	17	100.0	17	100.0	100.0	8	47.1	47.1
Backbone-Backbone	17	100.0	17	100.0	100.0	8	47.1	47.1
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0

¹ percentage calculated with respect to the total number of distance restraints, ² percentage calculated with respect to the number of restraints in a particular restraint category, ³ violated in at least one model, ⁴ violated in all the models

9.1.1 Bar chart : Distribution of distance restraints and violations [i](#)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfid bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model [i](#)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Model ID	Number of violations						Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total				
1	0	0	0	0	14	14	3.3	8.91	2.29	2.88
2	0	0	0	0	13	13	3.64	8.33	2.08	3.88
3	0	0	0	0	10	10	4.31	7.78	2.05	4.48
4	0	0	0	0	14	14	3.12	8.42	2.63	3.19
5	0	0	0	0	12	12	3.86	9.35	2.53	3.41
6	0	0	0	0	15	15	3.89	8.43	2.4	3.9
7	0	0	0	0	14	14	2.93	8.28	2.54	2.76
8	0	0	0	0	15	15	3.32	9.04	2.57	3.76
9	0	0	0	0	14	14	3.58	8.87	2.46	3.6
10	0	0	0	0	16	16	3.43	9.31	2.65	3.22

Continued on next page...


Continued from previous page...

Model ID	Number of violations						Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total				
11	0	0	0	0	16	16	3.2	8.69	2.61	2.74
12	0	0	0	0	17	17	3.44	9.5	2.48	3.11
13	0	0	0	0	14	14	3.89	9.97	2.66	3.74
14	0	0	0	0	16	16	2.94	7.9	2.58	2.19
15	0	0	0	0	11	11	3.48	8.44	2.73	2.74
16	0	0	0	0	15	15	3.13	8.58	2.44	3.01
17	0	0	0	0	16	16	3.12	9.04	2.64	2.1
18	0	0	0	0	16	16	2.84	10.0	2.78	2.0
19	0	0	0	0	16	16	3.19	8.46	2.65	3.08
20	0	0	0	0	15	15	3.61	11.07	2.96	2.95
21	0	0	0	0	14	14	3.53	10.24	2.75	2.96
22	0	0	0	0	15	15	3.66	8.51	2.72	4.15
23	0	0	0	0	15	15	3.04	9.02	2.42	3.07
24	0	0	0	0	14	14	3.86	8.68	2.4	4.19
25	0	0	0	0	12	12	3.16	7.46	2.45	3.22
26	0	0	0	0	15	15	3.22	9.47	2.72	2.5
27	0	0	0	0	15	15	4.04	11.02	2.9	3.13
28	0	0	0	0	16	16	2.47	8.16	2.48	1.3
29	0	0	0	0	13	13	3.81	8.79	2.54	3.84
30	0	0	0	0	15	15	3.67	10.34	2.85	3.34

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints,

⁵Inter-chain restraints, ⁶Standard deviation

9.2.1 Bar graph : Distance Violation statistics for each model [i](#)



The mean(dot), median(x) and the standard deviation are shown in blue with respect to the y axis on the right

9.3 Distance violation statistics for the ensemble [i](#)

Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 0(IR:0, SQ:0, MR:0, LR:0, IC:0) restraints are not violated in the ensemble.

Number of violated restraints						Fraction of the ensemble	
IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total	Count ⁶	%
0	0	0	0	1	1	1	3.3
0	0	0	0	0	0	2	6.7
0	0	0	0	0	0	3	10.0
0	0	0	0	0	0	4	13.3
0	0	0	0	0	0	5	16.7
0	0	0	0	0	0	6	20.0

Continued on next page...


Continued from previous page...

Number of violated restraints						Fraction of the ensemble	
IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total	Count ⁶	%
0	0	0	0	0	0	7	23.3
0	0	0	0	0	0	8	26.7
0	0	0	0	0	0	9	30.0
0	0	0	0	0	0	10	33.3
0	0	0	0	0	0	11	36.7
0	0	0	0	0	0	12	40.0
0	0	0	0	0	0	13	43.3
0	0	0	0	0	0	14	46.7
0	0	0	0	0	0	15	50.0
0	0	0	0	0	0	16	53.3
0	0	0	0	0	0	17	56.7
0	0	0	0	0	0	18	60.0
0	0	0	0	0	0	19	63.3
0	0	0	0	2	2	20	66.7
0	0	0	0	0	0	21	70.0
0	0	0	0	0	0	22	73.3
0	0	0	0	2	2	23	76.7
0	0	0	0	2	2	24	80.0
0	0	0	0	0	0	25	83.3
0	0	0	0	0	0	26	86.7
0	0	0	0	0	0	27	90.0
0	0	0	0	0	0	28	93.3
0	0	0	0	2	2	29	96.7
0	0	0	0	8	8	30	100.0

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints,

⁵Inter-chain restraints, ⁶ Number of models with violations


9.3.1 Bar graph : Distance violation statistics for the ensemble [i](#)



9.4 Most violated distance restraints in the ensemble [i](#)

9.4.1 Histogram : Distribution of mean distance violations [i](#)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble



9.4.2 Table: Most violated distance restraints [i](#)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Models ¹	Mean (Å)	SD ¹ (Å)	Median (Å)
(1,1)	1:10:A:ALA:HB1	2:7:B:C:O2	30	8.95	0.92	8.83
(1,1)	1:10:A:ALA:HB3	2:7:B:C:O2	30	8.95	0.92	8.83
(1,1)	1:10:A:ALA:HB2	2:7:B:C:O2	30	8.95	0.92	8.83
(1,8)	1:109:A:ARG:HH21	2:7:B:C:O2	30	7.48	0.84	7.61
(1,8)	1:109:A:ARG:HH22	2:7:B:C:O2	30	7.48	0.84	7.61
(1,8)	1:109:A:ARG:HH12	2:7:B:C:O2	30	7.48	0.84	7.61
(1,8)	1:109:A:ARG:HH22	2:7:B:C:H41	30	7.48	0.84	7.61
(1,4)	1:52:A:ARG:O	2:1:B:C:H42	30	5.03	0.44	5.04
(1,4)	1:52:A:ARG:HB3	2:1:B:C:H5	30	5.03	0.44	5.04
(1,4)	1:52:A:ARG:O	2:1:B:C:H5	30	5.03	0.44	5.04
(1,4)	1:52:A:ARG:HB3	2:1:B:C:H42	30	5.03	0.44	5.04
(1,16)	2:6:B:A:H2	1:112:A:ALA:HA	30	4.99	0.54	4.81
(1,16)	2:6:B:A:H2	1:112:A:ALA:HB3	30	4.99	0.54	4.81
(1,16)	2:6:B:A:H2	1:112:A:ALA:HB1	30	4.99	0.54	4.81
(1,16)	2:6:B:A:H61	1:62:A:LYS:HZ1	30	4.99	0.54	4.81
(1,16)	2:6:B:A:H61	1:62:A:LYS:HZ2	30	4.99	0.54	4.81

Continued on next page...

Continued from previous page...


Key	Atom-1	Atom-2	Models ¹	Mean (Å)	SD ¹ (Å)	Median (Å)
(1,5)	1:54:A:ILE:HD13	2:2:B:A:H61	30	4.42	0.6	4.38
(1,5)	1:54:A:ILE:HD11	2:2:B:A:H61	30	4.42	0.6	4.38
(1,5)	1:54:A:ILE:HD12	2:2:B:A:H61	30	4.42	0.6	4.38
(1,5)	1:54:A:ILE:HD13	2:1:B:C:H42	30	4.42	0.6	4.38
(1,5)	1:54:A:ILE:HD11	2:1:B:C:H42	30	4.42	0.6	4.38
(1,15)	2:5:B:G:O6	1:62:A:LYS:HZ1	30	4.32	0.65	4.44
(1,15)	2:5:B:G:O6	1:62:A:LYS:HZ3	30	4.32	0.65	4.44
(1,15)	2:5:B:G:O6	1:62:A:LYS:HZ2	30	4.32	0.65	4.44
(1,3)	1:19:A:HIS:HD1	2:1:B:C:H5'	30	3.87	1.01	4.12
(1,3)	1:19:A:HIS:HB2	2:1:B:C:H5'	30	3.87	1.01	4.12
(1,3)	1:19:A:HIS:HD1	2:1:B:C:H1'	30	3.87	1.01	4.12
(1,3)	1:19:A:HIS:HE1	2:1:B:C:H4'	30	3.87	1.01	4.12
(1,3)	1:19:A:HIS:HD1	2:1:B:C:H4'	30	3.87	1.01	4.12
(1,14)	2:4:B:U:O4	1:62:A:LYS:HZ1	30	2.68	0.66	2.78
(1,14)	2:4:B:U:O4	1:62:A:LYS:HZ3	30	2.68	0.66	2.78
(1,14)	2:4:B:U:O4	1:62:A:LYS:HZ2	30	2.68	0.66	2.78
(1,6)	1:65:A:SER:HG	2:1:B:C:H5'	29	2.05	1.24	2.07
(1,6)	1:65:A:SER:OG	2:1:B:C:H5'	29	2.05	1.24	2.07
(1,6)	1:65:A:SER:H	2:1:B:C:H5	29	2.05	1.24	2.07
(1,6)	1:65:A:SER:HG	2:1:B:C:O4'	29	2.05	1.24	2.07
(1,6)	1:65:A:SER:H	2:1:B:C:H6	29	2.05	1.24	2.07
(1,6)	1:65:A:SER:OG	2:1:B:C:O4'	29	2.05	1.24	2.07
(1,13)	2:3:B:C:H41	1:62:A:LYS:HZ3	29	0.97	0.4	0.79

¹Number of violated models, ²Standard deviation

9.5 All violated distance restraints [i](#)

9.5.1 Histogram : Distribution of distance violations [i](#)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



9.5.2 Table : All distance violations [i](#)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,1)	1:10:A:ALA:HB1	2:7:B:C:O2	20	11.07
(1,1)	1:10:A:ALA:HB1	2:7:B:C:O2	27	11.02
(1,1)	1:10:A:ALA:HB3	2:7:B:C:O2	30	10.34
(1,1)	1:10:A:ALA:HB1	2:7:B:C:O2	21	10.24
(1,1)	1:10:A:ALA:HB1	2:7:B:C:O2	18	10.0
(1,1)	1:10:A:ALA:HB3	2:7:B:C:O2	13	9.97
(1,1)	1:10:A:ALA:HB3	2:7:B:C:O2	12	9.5
(1,1)	1:10:A:ALA:HB3	2:7:B:C:O2	26	9.47
(1,1)	1:10:A:ALA:HB1	2:7:B:C:O2	5	9.35
(1,1)	1:10:A:ALA:HB3	2:7:B:C:O2	10	9.31

10 Dihedral-angle violation analysis ⓘ

No dihedral-angle restraints found