



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 01:36 PM UTC

PDB ID : 8AC8 / pdb_00008ac8
Title : The nucleoprotein complex of Rep protein with DUE ssDNA
Authors : Nowacka, M.; Wegrzyn, K.; Oliwa, M.; Konieczny, I.
Deposited on : 2022-07-05
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	251	8% 80% 6% 14%
1	C	251	5% 81% 5% 14%
2	B	9	78% 22%
2	D	9	11% 89% 11%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4307 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Replication initiation protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	216	Total	C	N	O	S	0	6	0
			1799	1161	308	323	7			
1	A	217	Total	C	N	O	S	0	11	0
			1811	1172	308	324	7			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	118	PRO	ARG	conflict	UNP P03856
A	118	PRO	ARG	conflict	UNP P03856

- Molecule 2 is a DNA chain called DNA (5'-D(*AP*TP*TP*TP*TP*TP*AP*TP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	9	Total	C	N	O	P	0	1	0
			181	90	27	56	8			
2	D	9	Total	C	N	O	P	0	1	0
			189	95	28	58	8			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	134	Total	O	0	0
			134	134		
3	A	148	Total	O	0	0
			148	148		
3	B	25	Total	O	0	0
			25	25		
3	D	20	Total	O	0	0
			20	20		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Replication initiation protein




- Molecule 1: Replication initiation protein



- Molecule 2: DNA (5'-D(*AP*TP*TP*TP*TP*TP*AP*TP*A)-3')



- Molecule 2: DNA (5'-D(*AP*TP*TP*TP*TP*TP*AP*TP*A)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, α , β , γ	47.40Å 85.31Å 146.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.85 – 1.60 36.85 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.2 (36.85-1.60) 99.2 (36.85-1.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.07 (at 1.60Å)	Xtriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.212 , 0.243 0.213 , 0.242	Depositor DCC
R_{free} test set	2000 reflections (2.52%)	wwPDB-VP
Wilson B-factor (Å ²)	33.1	Xtriage
Anisotropy	0.283	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 55.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4307	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 38.88 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.4517e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.28	0/1886	0.52	0/2548
1	C	0.26	0/1861	0.48	0/2511
2	B	0.31	0/201	0.61	0/310
2	D	0.31	0/210	0.64	0/323
All	All	0.28	0/4158	0.52	0/5692

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1811	0	1786	14	1
1	C	1799	0	1768	8	0
2	B	181	0	107	2	0
2	D	189	0	115	1	0
3	A	148	0	0	1	0
3	B	25	0	0	0	0
3	C	134	0	0	2	1
3	D	20	0	0	0	0
All	All	4307	0	3776	23	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:214:ASN:ND2	3:C:301:HOH:O	2.14	0.80
2:D:1[A]:DA:N3	2:D:1[A]:DA:O5'	2.32	0.60
1:A:185:ILE:HD13	1:A:201:MET:HE1	1.85	0.59
1:A:227[B]:ILE:HD11	1:A:229:LYS:NZ	2.18	0.58
1:A:201:MET:HE2	1:A:201:MET:HA	1.89	0.55
1:C:57:ASP:O	1:C:130:HIS:NE2	2.40	0.53
1:C:117:LYS:NZ	1:C:118:PRO:O	2.40	0.52
1:A:157:ILE:HG23	1:A:215:GLU:HG2	1.94	0.48
1:A:15:SER:HB2	3:A:392:HOH:O	2.14	0.47
1:C:31:LEU:HD23	1:C:92:LYS:HD3	1.97	0.46
1:C:206:ARG:NH2	1:A:90:ALA:O	2.48	0.45
1:A:228:GLU:O	1:A:229:LYS:HD3	2.17	0.45
1:A:28:ALA:HB1	1:A:193[B]:GLN:OE1	2.18	0.44
1:A:227[B]:ILE:HD11	1:A:229:LYS:HZ3	1.80	0.44
1:C:45:GLN:NE2	1:C:61:GLU:O	2.33	0.43
2:B:1[B]:DA:C2	2:B:2:DT:C4	3.07	0.42
1:A:213:VAL:HG21	1:A:226:TYR:HD2	1.84	0.42
1:A:17:ARG:HH22	2:B:1[B]:DA:H3'	1.85	0.42
1:A:185:ILE:HG21	1:A:201:MET:HE1	2.02	0.41
1:A:201:MET:SD	1:A:235:THR:OG1	2.66	0.41
1:A:229:LYS:HD3	1:A:229:LYS:HA	1.80	0.40
1:C:58:GLY:CA	1:C:136[B]:ILE:HD11	2.52	0.40
1:C:205:ARG:NH1	3:C:308:HOH:O	2.54	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:50:ASP:OD2	3:C:366:HOH:O[4_555]	2.03	0.17

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	222/251 (88%)	221 (100%)	1 (0%)	0	100	100
1	C	216/251 (86%)	213 (99%)	3 (1%)	0	100	100
All	All	438/502 (87%)	434 (99%)	4 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	195/226 (86%)	195 (100%)	0	100	100
1	C	194/226 (86%)	194 (100%)	0	100	100
All	All	389/452 (86%)	389 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	63	HIS
1	C	84	GLN
1	C	214	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	217/251 (86%)	0.58	19 (8%) 15 15	21, 36, 62, 83	11 (5%)
1	C	216/251 (86%)	0.60	13 (6%) 27 28	21, 37, 65, 89	8 (3%)
2	B	9/9 (100%)	0.42	0 100 100	34, 42, 60, 63	1 (11%)
2	D	9/9 (100%)	0.55	1 (11%) 10 10	29, 42, 67, 103	1 (11%)
All	All	451/520 (86%)	0.59	33 (7%) 21 21	21, 37, 64, 103	21 (4%)

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	109	TYR	4.8
1	A	226	TYR	4.5
1	A	227[A]	ILE	4.4
1	C	245	ILE	4.3
1	A	14	ASN	4.1
1	A	197	SER	3.9
1	C	247	SER	3.6
1	C	109	TYR	3.6
1	A	202	PRO	3.5
1	C	56	HIS	3.3
1	C	97	TYR	3.3
1	A	50	ASP	3.2
1	C	246	THR	3.1
1	A	203	ASP	3.0
1	A	201	MET	3.0
2	D	9	DA	2.9
1	A	247	SER	2.7
1	A	232	GLY	2.7
1	C	62	ILE	2.6
1	A	228	GLU	2.6
1	C	232	GLY	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	202	PRO	2.6
1	A	49	SER	2.5
1	C	49	SER	2.5
1	A	62	ILE	2.5
1	C	207[A]	ARG	2.4
1	A	246	THR	2.4
1	A	97	TYR	2.2
1	C	211	VAL	2.1
1	A	230	LYS	2.1
1	C	233	ARG	2.1
1	A	198	TYR	2.1
1	A	196	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.