



wwPDB NMR Structure Validation Summary Report ⓘ

Mar 6, 2026 – 05:02 PM UTC

PDB ID : 2ADC / pdb_00002adc
Title : Solution structure of Polypyrimidine Tract Binding protein RBD34 complexed with CUCUCU RNA
Authors : Oberstrass, F.C.; Auweter, S.D.; Erat, M.; Hargous, Y.; Henning, A.; Wenter, P.; Reymond, L.; Pitsch, S.; Black, D.L.; Allain, F.H.T.
Deposited on : 2005-07-20

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823
RNA backbone	8273	777

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	B	6	17% 67% 17%
1	C	6	17% 67% 17%
2	A	229	72% 7% 12% 9%

2 Ensemble composition and analysis i

This entry contains 20 models. Model 11 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:336-A:417, A:426-A:438, A:446-A:531 (181)	1.19	11

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 4 single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 4, 5, 6, 9, 10, 11, 12, 15, 18, 20
2	13, 14, 19
Single-model clusters	7; 8; 16; 17

3 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 3623 atoms, of which 1771 are hydrogens and 0 are deuteriums.

- Molecule 1 is a RNA chain called 5'-R(*CP*UP*CP*UP*CP*U)-3'.

Mol	Chain	Residues	Atoms					Trace	
1	B	6	Total	C	H	N	O	P	0
			182	54	65	15	43	5	
1	C	6	Total	C	H	N	O	P	0
			182	54	65	15	43	5	

- Molecule 2 is a protein called Polypyrimidine tract-binding protein 1.

Mol	Chain	Residues	Atoms					Trace	
2	A	208	Total	C	H	N	O	S	0
			3259	1023	1641	296	295	4	

There are 21 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	303	MET	-	expression tag	UNP P26599
A	304	GLY	-	expression tag	UNP P26599
A	305	SER	-	expression tag	UNP P26599
A	306	SER	-	expression tag	UNP P26599
A	307	HIS	-	expression tag	UNP P26599
A	308	HIS	-	expression tag	UNP P26599
A	309	HIS	-	expression tag	UNP P26599
A	310	HIS	-	expression tag	UNP P26599
A	311	HIS	-	expression tag	UNP P26599
A	312	HIS	-	expression tag	UNP P26599
A	313	SER	-	expression tag	UNP P26599
A	314	SER	-	expression tag	UNP P26599
A	315	GLY	-	expression tag	UNP P26599
A	316	LEU	-	expression tag	UNP P26599
A	317	VAL	-	expression tag	UNP P26599
A	318	PRO	-	expression tag	UNP P26599
A	319	ARG	-	expression tag	UNP P26599
A	320	GLY	-	expression tag	UNP P26599
A	321	SER	-	expression tag	UNP P26599
A	322	HIS	-	expression tag	UNP P26599
A	323	MET	-	expression tag	UNP P26599

4 Residue-property plots


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: 5'-R(*CP*UP*CP*UP*CP*U)-3'



- Molecule 1: 5'-R(*CP*UP*CP*UP*CP*U)-3'




- Molecule 2: Polypyrimidine tract-binding protein 1




4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 11. Colouring as in section 4.1 above.

- Molecule 1: 5'-R(*CP*UP*CP*UP*CP*U)-3'




- Molecule 1: 5'-R(*CP*UP*CP*UP*CP*U)-3'

Chain C:  17% 50% 33%



- Molecule 2: Polypyrimidine tract-binding protein 1

Chain A:  66% 11% 12% 9%



5 Refinement protocol and experimental data overview

The models were refined using the following method: *distance geometry simulated annealing*.

Of the 40 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DYANA	structure solution	3.02
Amber	refinement	7

No chemical shift data was provided.

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	B	0.91±0.02	0±0/128 (0.0± 0.0%)	1.33±0.06	1±1/196 (0.6± 0.4%)
1	C	0.89±0.03	0±0/128 (0.0± 0.0%)	1.32±0.06	0±0/196 (0.2± 0.2%)
2	A	0.88±0.01	0±0/1461 (0.0± 0.0%)	1.55±0.02	7±3/1972 (0.3± 0.1%)
All	All	0.89	1/34340 (0.0%)	1.52	170/47280 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	B	0.0±0.0	1.4±1.1
1	C	0.0±0.0	1.5±0.6
2	A	0.0±0.0	3.0±1.6
All	All	0	118

All unique bond outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
2	A	451	PRO	CA-C	5.01	1.54	1.51	1	1

5 of 79 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	B	536	C	C4'-C3'-O3'	8.89	122.74	109.40	4	3
2	A	483	GLY	CA-C-N	8.01	136.11	121.70	15	2
2	A	483	GLY	C-N-CA	8.01	136.11	121.70	15	2
1	B	536	C	C3'-C2'-C1'	6.84	108.14	101.30	7	9
2	A	358	PHE	CA-C-N	6.81	128.65	120.14	8	6

There are no chirality outliers.

5 of 33 unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	C	540	C	Sidechain	20
2	A	450	PHE	Sidechain	12
2	A	527	SER	Peptide	10
2	A	492	LYS	Peptide	9
1	B	536	C	Sidechain	8

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	B	117	65	65	0±0
1	C	117	65	65	0±1
2	A	1430	1450	1450	1±1
All	All	33280	31600	31600	17

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

5 of 10 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:C:540:C:C5	2:A:340:LEU:HD22	0.74	2.18	7	2
2:A:338:VAL:HG23	2:A:409:SER:HB2	0.53	1.79	10	4
1:B:536:C:H2'	2:A:528:LYS:HE3	0.48	1.85	10	1
1:C:540:C:C4	2:A:340:LEU:HD22	0.46	2.46	19	1
1:B:536:C:N4	2:A:450:PHE:CE1	0.45	2.84	13	2

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation

was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	180/229 (79%)	152±4 (84±2%)	22±3 (12±2%)	6±1 (3±1%)	5	35
All	All	3600/4580 (79%)	3039 (84%)	441 (12%)	120 (3%)	5	35

5 of 42 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	A	528	LYS	13
2	A	362	GLY	11
2	A	484	PHE	7
2	A	371	PHE	7
2	A	480	VAL	5

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	162/197 (82%)	147±3 (91±2%)	15±3 (9±2%)	10	55
All	All	3240/3940 (82%)	2933 (91%)	307 (9%)	10	55

5 of 79 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	A	378	LEU	20
2	A	379	VAL	20
2	A	496	ILE	14
2	A	495	LEU	13
2	A	389	LEU	11

6.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers	Suiteness
1	B	5/6 (83%)	4±0 (80±0%)	1±1 (29±17%)	0.60±0.00

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers	Suiteness
1	C	5/6 (83%)	5±0 (100±0%)	1±1 (20±15%)	0.00±0.00
All	All	202/240 (84%)	180 (89%)	49 (24%)	0.15

The overall RNA backbone suiteness is 0.05.

5 of 9 unique RNA backbone outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
1	B	534	C	20
1	B	535	U	20
1	B	536	C	20
1	B	537	U	20
1	C	539	U	20

5 of 8 unique RNA pucker outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
1	B	536	C	16
1	C	542	C	9
1	B	534	C	8
1	C	539	U	5
1	C	540	C	4

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided