



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 09:27 PM UTC

PDB ID : 4ADQ / pdb_00004adq
Title : CRYSTAL STRUCTURE OF THE MOUSE COLONY-STIMULATING FACTOR 1 (MCSF-1) CYTOKINE IN COMPLEX WITH THE VIRAL RECEPTOR BARF1
Authors : Elegheert, J.; Bracke, N.; Savvides, S.N.
Deposited on : 2012-01-02
Resolution : 4.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 4.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1127 (5.10-3.90)
Clashscore	190562	1002 (5.06-3.94)
Ramachandran outliers	187476	1060 (5.10-3.90)
Sidechain outliers	187428	1043 (5.10-3.90)
RSRZ outliers	180081	1122 (5.10-3.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	208	 78% 11% 11%
1	B	208	 74% 15% 11%
1	C	208	 73% 17% 11%
1	D	208	 76% 13% 11%
2	E	153	 85% 9% 6%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	153	 79%14%7%
2	G	153	 3%84%9%7%
2	H	153	 80%13%7%
3	I	3	 33%67%
3	K	3	 33%67%
3	L	3	 67%33%
4	J	4	 75%25%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 10758 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SECRETED PROTEIN BARF1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	185	Total	C	N	O	S	0	0	0
			1477	955	251	264	7			
1	B	185	Total	C	N	O	S	0	0	0
			1477	955	251	264	7			
1	C	186	Total	C	N	O	S	0	0	0
			1483	958	252	266	7			
1	D	186	Total	C	N	O	S	0	0	0
			1483	958	252	266	7			

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	222	LYS	-	expression tag	UNP P0CW72
A	223	HIS	-	expression tag	UNP P0CW72
A	224	HIS	-	expression tag	UNP P0CW72
A	225	HIS	-	expression tag	UNP P0CW72
A	226	HIS	-	expression tag	UNP P0CW72
A	227	HIS	-	expression tag	UNP P0CW72
A	228	HIS	-	expression tag	UNP P0CW72
A	169	SER	THR	engineered mutation	UNP P0CW72
B	222	LYS	-	expression tag	UNP P0CW72
B	223	HIS	-	expression tag	UNP P0CW72
B	224	HIS	-	expression tag	UNP P0CW72
B	225	HIS	-	expression tag	UNP P0CW72
B	226	HIS	-	expression tag	UNP P0CW72
B	227	HIS	-	expression tag	UNP P0CW72
B	228	HIS	-	expression tag	UNP P0CW72
B	169	SER	THR	engineered mutation	UNP P0CW72
C	222	LYS	-	expression tag	UNP P0CW72
C	223	HIS	-	expression tag	UNP P0CW72
C	224	HIS	-	expression tag	UNP P0CW72
C	225	HIS	-	expression tag	UNP P0CW72
C	226	HIS	-	expression tag	UNP P0CW72

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	227	HIS	-	expression tag	UNP P0CW72
C	228	HIS	-	expression tag	UNP P0CW72
C	169	SER	THR	engineered mutation	UNP P0CW72
D	222	LYS	-	expression tag	UNP P0CW72
D	223	HIS	-	expression tag	UNP P0CW72
D	224	HIS	-	expression tag	UNP P0CW72
D	225	HIS	-	expression tag	UNP P0CW72
D	226	HIS	-	expression tag	UNP P0CW72
D	227	HIS	-	expression tag	UNP P0CW72
D	228	HIS	-	expression tag	UNP P0CW72
D	169	SER	THR	engineered mutation	UNP P0CW72

- Molecule 2 is a protein called MACROPHAGE COLONY-STIMULATING FACTOR 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	144	Total	C	N	O	S	0	0	0
			1174	734	198	232	10			
2	F	142	Total	C	N	O	S	0	0	0
			1159	725	197	227	10			
2	G	143	Total	C	N	O	S	0	0	0
			1172	734	198	230	10			
2	H	142	Total	C	N	O	S	0	0	0
			1166	731	197	228	10			

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
E	-3	GLY	-	expression tag	UNP P07141
E	-2	SER	-	expression tag	UNP P07141
E	-1	HIS	-	expression tag	UNP P07141
E	0	MET	-	expression tag	UNP P07141
F	-3	GLY	-	expression tag	UNP P07141
F	-2	SER	-	expression tag	UNP P07141
F	-1	HIS	-	expression tag	UNP P07141
F	0	MET	-	expression tag	UNP P07141
G	-3	GLY	-	expression tag	UNP P07141
G	-2	SER	-	expression tag	UNP P07141
G	-1	HIS	-	expression tag	UNP P07141
G	0	MET	-	expression tag	UNP P07141
H	-3	GLY	-	expression tag	UNP P07141
H	-2	SER	-	expression tag	UNP P07141
H	-1	HIS	-	expression tag	UNP P07141

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	0	MET	-	expression tag	UNP P07141

- Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	I	3	Total	C	N	O	0	0	0
			39	22	2	15			
3	K	3	Total	C	N	O	0	0	0
			39	22	2	15			
3	L	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	J	4	Total	C	N	O	0	0	0
			50	28	2	20			

- Molecule 1: SECRETED PROTEIN BARF1





• Molecule 2: MACROPHAGE COLONY-STIMULATING FACTOR 1

Chain E: 85% 9% 6%



• Molecule 2: MACROPHAGE COLONY-STIMULATING FACTOR 1

Chain F: 79% 14% 7%



• Molecule 2: MACROPHAGE COLONY-STIMULATING FACTOR 1

Chain G: 3% 84% 9% 7%



• Molecule 2: MACROPHAGE COLONY-STIMULATING FACTOR 1

Chain H: 80% 13% 7%



• Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I: 33% 67%



• Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K: 33% 67%



• Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain L:  67% 33%

HA01
HA02
BM03

- Molecule 4: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J:  75% 25%

HA01
HA02
BM03
NA04

4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	234.75Å 234.75Å 96.04Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	117.38 – 4.50 117.38 – 4.50	Depositor EDS
% Data completeness (in resolution range)	94.5 (117.38-4.50) 94.5 (117.38-4.50)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.69 (at 4.47Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.223 , 0.262 0.254 , 0.282	Depositor DCC
R_{free} test set	1112 reflections (10.03%)	wwPDB-VP
Wilson B-factor (Å ²)	138.7	Xtriage
Anisotropy	0.310	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 196.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	0.052 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	10758	wwPDB-VP
Average B, all atoms (Å ²)	187.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, BMA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.28	0/1520	0.67	0/2069
1	B	0.27	0/1520	0.66	0/2069
1	C	0.29	0/1526	0.68	0/2077
1	D	0.28	0/1526	0.67	0/2077
2	E	0.27	0/1195	0.63	0/1611
2	F	0.28	0/1179	0.65	0/1588
2	G	0.28	0/1193	0.63	0/1607
2	H	0.28	0/1187	0.63	0/1599
All	All	0.28	0/10846	0.66	0/14697

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1477	0	1445	18	0
1	B	1477	0	1445	23	0
1	C	1483	0	1450	25	0
1	D	1483	0	1450	18	0
2	E	1174	0	1113	11	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	F	1159	0	1107	17	0
2	G	1172	0	1119	11	0
2	H	1166	0	1114	16	0
3	I	39	0	34	1	0
3	K	39	0	34	1	0
3	L	39	0	34	0	0
4	J	50	0	43	0	0
All	All	10758	0	10388	132	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 132 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:G:52:LYS:NZ	2:G:98:GLN:O	2.24	0.71
2:E:52:LYS:NZ	2:E:98:GLN:O	2.24	0.70
1:B:193:LYS:HG3	1:B:194:PRO:HA	1.78	0.63
1:D:50:LYS:HD3	1:D:102:TYR:CE2	2.33	0.63
1:C:105:ARG:NH1	1:D:105:ARG:NH2	2.47	0.62

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	181/208 (87%)	177 (98%)	3 (2%)	1 (1%)	21 58
1	B	181/208 (87%)	177 (98%)	3 (2%)	1 (1%)	21 58
1	C	182/208 (88%)	178 (98%)	3 (2%)	1 (0%)	24 63
1	D	182/208 (88%)	177 (97%)	4 (2%)	1 (0%)	24 63

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	E	142/153 (93%)	139 (98%)	3 (2%)	0	100	100
2	F	140/153 (92%)	138 (99%)	2 (1%)	0	100	100
2	G	141/153 (92%)	138 (98%)	3 (2%)	0	100	100
2	H	140/153 (92%)	138 (99%)	2 (1%)	0	100	100
All	All	1289/1444 (89%)	1262 (98%)	23 (2%)	4 (0%)	36	71

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	41	GLY
1	B	41	GLY
1	C	41	GLY
1	D	41	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	165/183 (90%)	165 (100%)	0	100	100
1	B	165/183 (90%)	164 (99%)	1 (1%)	78	80
1	C	166/183 (91%)	166 (100%)	0	100	100
1	D	166/183 (91%)	165 (99%)	1 (1%)	78	80
2	E	135/144 (94%)	135 (100%)	0	100	100
2	F	133/144 (92%)	133 (100%)	0	100	100
2	G	135/144 (94%)	135 (100%)	0	100	100
2	H	134/144 (93%)	134 (100%)	0	100	100
All	All	1199/1308 (92%)	1197 (100%)	2 (0%)	87	85

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	180	ASP
1	D	180	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
2	H	6	HIS
2	H	107	HIS
2	H	85	ASN
2	F	6	HIS
2	G	13	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

13 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	I	1	3,1	14,14,15	0.50	0	17,19,21	0.79	0
3	NAG	I	2	3	14,14,15	0.53	0	17,19,21	0.58	0
3	BMA	I	3	3	11,11,12	0.57	0	15,15,17	0.59	0
4	NAG	J	1	1,4	14,14,15	0.54	0	17,19,21	0.79	0
4	NAG	J	2	4	14,14,15	0.48	0	17,19,21	0.79	0
4	BMA	J	3	4	11,11,12	0.61	0	15,15,17	0.87	1 (6%)
4	MAN	J	4	4	11,11,12	0.54	0	15,15,17	0.84	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	K	1	3,1	14,14,15	0.47	0	17,19,21	0.84	0
3	NAG	K	2	3	14,14,15	0.52	0	17,19,21	0.64	0
3	BMA	K	3	3	11,11,12	0.61	0	15,15,17	0.57	0
3	NAG	L	1	3,1	14,14,15	0.53	0	17,19,21	0.70	0
3	NAG	L	2	3	14,14,15	0.52	0	17,19,21	0.84	0
3	BMA	L	3	3	11,11,12	0.58	0	15,15,17	0.91	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	I	1	3,1	-	1/6/23/26	0/1/1/1
3	NAG	I	2	3	-	5/6/23/26	0/1/1/1
3	BMA	I	3	3	-	0/2/19/22	0/1/1/1
4	NAG	J	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	J	2	4	-	0/6/23/26	0/1/1/1
4	BMA	J	3	4	-	0/2/19/22	0/1/1/1
4	MAN	J	4	4	-	0/2/19/22	0/1/1/1
3	NAG	K	1	3,1	-	1/6/23/26	0/1/1/1
3	NAG	K	2	3	-	0/6/23/26	0/1/1/1
3	BMA	K	3	3	-	0/2/19/22	0/1/1/1
3	NAG	L	1	3,1	-	1/6/23/26	0/1/1/1
3	NAG	L	2	3	-	0/6/23/26	0/1/1/1
3	BMA	L	3	3	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	L	3	BMA	C1-C2-C3	2.57	113.39	109.64
4	J	3	BMA	C1-C2-C3	2.07	112.66	109.64

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	I	1	NAG	C1-C2-N2-C7
3	I	2	NAG	C1-C2-N2-C7
3	K	1	NAG	C1-C2-N2-C7

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	I	2	NAG	C4-C5-C6-O6
3	I	2	NAG	O5-C5-C6-O6



There are no ring outliers.


4 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	I	2	NAG	1	0
3	I	3	BMA	1	0
3	K	2	NAG	1	0
3	K	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	185/208 (88%)	-0.10	0	100	100	122, 162, 201, 219	0
1	B	185/208 (88%)	-0.03	0	100	100	121, 162, 196, 219	0
1	C	186/208 (89%)	-0.14	1 (0%)	87	75	129, 162, 201, 228	0
1	D	186/208 (89%)	0.00	0	100	100	122, 162, 199, 234	0
2	E	144/153 (94%)	0.13	0	100	100	142, 217, 254, 277	0
2	F	142/153 (92%)	-0.12	0	100	100	141, 216, 254, 279	0
2	G	143/153 (93%)	0.16	4 (2%)	55	43	139, 216, 251, 275	0
2	H	142/153 (92%)	-0.11	0	100	100	142, 218, 256, 280	0
All	All	1313/1444 (90%)	-0.03	5 (0%)	88	77	121, 179, 248, 280	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	G	65	MET	2.8
2	G	87	LEU	2.7
1	C	148	VAL	2.5
2	G	61	ILE	2.1
2	G	80	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	BMA	I	3	11/12	0.57	0.12	128,168,198,201	0
4	BMA	J	3	11/12	0.58	0.08	152,186,205,211	0
4	MAN	J	4	11/12	0.61	0.15	205,232,270,272	0
3	BMA	K	3	11/12	0.75	0.11	134,183,206,212	0
3	BMA	L	3	11/12	0.77	0.09	114,166,179,183	0
3	NAG	K	1	14/15	0.84	0.12	124,172,199,213	0
3	NAG	L	2	14/15	0.85	0.10	155,163,188,204	0
3	NAG	K	2	14/15	0.89	0.09	144,161,169,179	0
3	NAG	I	2	14/15	0.90	0.09	177,203,208,214	0
3	NAG	L	1	14/15	0.91	0.11	123,162,214,226	0
4	NAG	J	1	14/15	0.92	0.10	135,163,189,212	0
3	NAG	I	1	14/15	0.93	0.08	145,179,206,212	0
4	NAG	J	2	14/15	0.95	0.10	114,153,183,184	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain K:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain L:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain J:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.