



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 10:51 PM UTC

PDB ID : 8ADF / pdb_00008adf
Title : X-ray crystal structure of PPAR gamma ligand binding domain in complex with CZ39
Authors : Capelli, D.; Montanari, R.; Pochetti, G.; Meneghetti, F.; Villa, S.
Deposited on : 2022-07-08
Resolution : 2.13 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.13 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3689 (2.16-2.12)
Clashscore	190562	3812 (2.16-2.12)
Ramachandran outliers	187476	3773 (2.16-2.12)
Sidechain outliers	187428	3772 (2.16-2.12)
RSRZ outliers	180081	3691 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	303	6% 70% 14% 15%
1	B	303	10% 63% 12% 25%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4175 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peroxisome proliferator-activated receptor gamma.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	257	Total	C	N	O	S	0	0	0
			2053	1324	335	384	10			
1	B	228	Total	C	N	O	S	0	0	0
			1823	1182	296	336	9			

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	174	MET	-	initiating methionine	UNP P37231
A	175	GLY	-	expression tag	UNP P37231
A	176	SER	-	expression tag	UNP P37231
A	177	SER	-	expression tag	UNP P37231
A	178	HIS	-	expression tag	UNP P37231
A	179	HIS	-	expression tag	UNP P37231
A	180	HIS	-	expression tag	UNP P37231
A	181	HIS	-	expression tag	UNP P37231
A	182	HIS	-	expression tag	UNP P37231
A	183	HIS	-	expression tag	UNP P37231
A	184	SER	-	expression tag	UNP P37231
A	185	SER	-	expression tag	UNP P37231
A	186	GLY	-	expression tag	UNP P37231
A	187	LEU	-	expression tag	UNP P37231
A	188	VAL	-	expression tag	UNP P37231
A	189	PRO	-	expression tag	UNP P37231
A	190	ARG	-	expression tag	UNP P37231
A	191	GLY	-	expression tag	UNP P37231
A	192	SER	-	expression tag	UNP P37231
A	193	HIS	-	expression tag	UNP P37231
A	194	MET	-	expression tag	UNP P37231
B	174	MET	-	initiating methionine	UNP P37231
B	175	GLY	-	expression tag	UNP P37231
B	176	SER	-	expression tag	UNP P37231
B	177	SER	-	expression tag	UNP P37231

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	178	HIS	-	expression tag	UNP P37231
B	179	HIS	-	expression tag	UNP P37231
B	180	HIS	-	expression tag	UNP P37231
B	181	HIS	-	expression tag	UNP P37231
B	182	HIS	-	expression tag	UNP P37231
B	183	HIS	-	expression tag	UNP P37231
B	184	SER	-	expression tag	UNP P37231
B	185	SER	-	expression tag	UNP P37231
B	186	GLY	-	expression tag	UNP P37231
B	187	LEU	-	expression tag	UNP P37231
B	188	VAL	-	expression tag	UNP P37231
B	189	PRO	-	expression tag	UNP P37231
B	190	ARG	-	expression tag	UNP P37231
B	191	GLY	-	expression tag	UNP P37231
B	192	SER	-	expression tag	UNP P37231
B	193	HIS	-	expression tag	UNP P37231
B	194	MET	-	expression tag	UNP P37231

- Molecule 2 is (2R)-3-(4-bromophenyl)-2-(3-hydroxyphenyl)-4-oxidanyl-2H-furan-5-one (CCD ID: LRA) (formula: C₁₆H₁₁BrO₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	Br	C	O	0	0
			21	1	16	4		
2	B	1	Total	Br	C	O	0	0
			21	1	16	4		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	137	Total 137	O 137	0	0
3	B	120	Total 120	O 120	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Peroxisome proliferator-activated receptor gamma



- Molecule 1: Peroxisome proliferator-activated receptor gamma



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	92.88Å 59.95Å 117.37Å 90.00° 103.29° 90.00°	Depositor
Resolution (Å)	57.11 – 2.13 57.11 – 2.13	Depositor EDS
% Data completeness (in resolution range)	95.4 (57.11-2.13) 95.6 (57.11-2.13)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.04 (at 2.05Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.216 , 0.257 0.218 , 0.260	Depositor DCC
R_{free} test set	1901 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	54.0	Xtriage
Anisotropy	0.336	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 62.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4175	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: LRA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.34	0/2087	0.54	0/2813
1	B	0.30	0/1852	0.49	0/2494
All	All	0.33	0/3939	0.52	0/5307

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2053	0	2111	38	0
1	B	1823	0	1870	25	0
2	A	21	0	0	0	0
2	B	21	0	0	1	0
3	A	137	0	0	6	0
3	B	120	0	0	3	0
All	All	4175	0	3981	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:358:LYS:HD3	1:B:359:PRO:HA	1.45	0.98
1:B:325:ILE:HD12	1:B:388:ILE:HG23	1.67	0.75
1:A:239:GLY:C	1:A:241:THR:H	1.97	0.72
1:A:235:ALA:HA	1:A:240:LYS:HE3	1.75	0.69
1:A:277:VAL:N	3:A:605:HOH:O	2.26	0.68
1:A:422:LYS:NZ	3:A:603:HOH:O	2.24	0.66
1:B:468:LEU:N	3:B:602:HOH:O	2.28	0.66
1:A:330:LEU:HD21	1:A:364:MET:HE1	1.78	0.66
1:A:325:ILE:HD11	1:A:392:ILE:HG13	1.77	0.65
1:A:350:ARG:NH1	3:A:604:HOH:O	2.26	0.65
1:B:373:LYS:NZ	3:B:601:HOH:O	2.28	0.61
1:B:279:ILE:O	1:B:283:GLN:HG3	2.01	0.61
1:B:288:ARG:NH2	3:B:607:HOH:O	2.34	0.60
1:B:334:MET:HE1	1:B:367:LYS:HB3	1.84	0.60
1:A:293:VAL:HG22	1:A:322:VAL:HG11	1.84	0.58
1:A:466:HIS:ND1	1:A:467:PRO:HD2	2.19	0.58
1:A:440:THR:HG21	1:B:443:ARG:HD2	1.86	0.57
1:A:282:PHE:HZ	1:A:453:LEU:HD11	1.69	0.56
1:B:348:MET:HE1	2:B:501:LRA:CAQ	2.35	0.56
1:A:290:VAL:HG21	1:A:466:HIS:CD2	2.42	0.55
1:B:358:LYS:HD3	1:B:359:PRO:CA	2.30	0.55
1:A:359:PRO:HG2	1:A:456:ILE:CD1	2.40	0.51
1:A:216:LYS:HG2	3:A:678:HOH:O	2.08	0.51
1:A:467:PRO:HA	1:A:470:GLN:HB2	1.93	0.51
1:B:333:LEU:HB3	1:B:340:LEU:HB2	1.92	0.50
1:B:365:GLU:O	1:B:369:GLU:HG3	2.11	0.50
1:A:359:PRO:HG2	1:A:456:ILE:HD11	1.95	0.49
1:A:282:PHE:CE1	1:A:463:MET:HE3	2.48	0.49
1:A:457:LYS:HZ1	1:A:465:LEU:HB2	1.78	0.49
1:A:379:LEU:HD11	1:A:435:LEU:HD13	1.95	0.48
1:B:330:LEU:HG	1:B:334:MET:HE3	1.95	0.47
1:A:418:GLU:OE2	3:A:601:HOH:O	2.20	0.47
1:B:255:LEU:HD13	1:B:281:ILE:HD11	1.96	0.47
1:A:457:LYS:NZ	1:A:465:LEU:HB2	2.30	0.46
1:B:379:LEU:HD11	1:B:435:LEU:HD13	1.96	0.46
1:B:348:MET:SD	1:B:353:LEU:HD21	2.56	0.46
1:A:456:ILE:HG21	1:A:463:MET:HE1	1.97	0.46
1:A:453:LEU:O	1:A:457:LYS:HG2	2.16	0.45
1:A:204:LEU:H	1:A:204:LEU:HD23	1.80	0.45
1:A:392:ILE:HG22	1:A:393:LEU:HD22	1.99	0.45
1:B:379:LEU:HD21	1:B:431:LEU:HD11	1.99	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:205:ASN:HB2	1:A:206:PRO:HD2	1.99	0.44
1:A:440:THR:O	1:A:444:GLN:HG3	2.18	0.43
1:A:239:GLY:C	1:A:241:THR:N	2.68	0.43
1:B:245:SER:HA	1:B:246:PRO:HD3	1.85	0.43
1:B:380:ASP:OD1	1:B:382:SER:OG	2.31	0.43
1:B:469:LEU:HA	1:B:469:LEU:HD23	1.54	0.43
1:A:394:SER:O	1:A:397:ARG:HG2	2.19	0.43
1:A:443:ARG:NH1	1:B:444:GLN:OE1	2.50	0.43
1:B:362:ASP:N	1:B:362:ASP:OD1	2.51	0.42
1:A:415:GLN:HG2	1:B:430:GLN:HE21	1.84	0.42
1:A:222:TYR:CE1	1:A:381:ASP:HB3	2.54	0.42
1:B:320:TYR:HB3	1:B:397:ARG:HD2	2.01	0.42
1:A:453:LEU:HD12	1:A:453:LEU:HA	1.81	0.42
1:A:282:PHE:CD1	1:A:463:MET:HE3	2.55	0.41
1:A:240:LYS:NZ	3:A:619:HOH:O	2.54	0.40
1:A:287:PHE:CE1	1:A:466:HIS:HD2	2.40	0.40
1:A:290:VAL:HG21	1:A:466:HIS:CG	2.56	0.40
1:A:452:LEU:O	1:A:456:ILE:HG12	2.21	0.40
1:B:431:LEU:HD13	1:B:431:LEU:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	253/303 (84%)	247 (98%)	5 (2%)	1 (0%)	30	26
1	B	220/303 (73%)	218 (99%)	2 (1%)	0	100	100
All	All	473/606 (78%)	465 (98%)	7 (2%)	1 (0%)	43	42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	240	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	230/272 (85%)	224 (97%)	6 (3%)	40	42
1	B	202/272 (74%)	199 (98%)	3 (2%)	57	63
All	All	432/544 (79%)	423 (98%)	9 (2%)	47	51

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	221	SER
1	A	256	MET
1	A	277	VAL
1	A	311	LEU
1	A	322	VAL
1	A	372	VAL
1	B	210	ASP
1	B	255	LEU
1	B	445	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	294	GLN
1	A	312	ASN
1	A	314	GLN
1	A	410	GLN
1	B	253	ASN
1	B	286	GLN
1	B	415	GLN
1	B	430	GLN
1	B	437	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	470	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	LRA	A	501	-	22,23,23	2.78	5 (22%)	30,33,33	2.63	9 (30%)
2	LRA	B	501	-	22,23,23	2.50	3 (13%)	30,33,33	2.48	7 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LRA	A	501	-	-	4/8/24/24	0/3/3/3
2	LRA	B	501	-	-	2/8/24/24	0/3/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	LRA	CAF-CAH	-8.55	1.32	1.48
2	B	501	LRA	CAF-CAH	-7.18	1.35	1.48
2	B	501	LRA	CAO-CAL	-6.88	1.40	1.51
2	A	501	LRA	CAO-CAL	-6.46	1.41	1.51
2	A	501	LRA	CAI-CAJ	-5.89	1.31	1.45
2	B	501	LRA	CAI-CAJ	-5.35	1.32	1.45
2	A	501	LRA	CAI-CAH	-2.78	1.31	1.34
2	A	501	LRA	BRAG-CAC	-2.76	1.84	1.90

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	LRA	CAL-OAK-CAJ	-8.62	104.11	109.79
2	B	501	LRA	CAL-OAK-CAJ	-7.01	105.18	109.79
2	A	501	LRA	OAN-CAJ-CAI	-5.57	120.97	129.33
2	B	501	LRA	OAN-CAJ-CAI	-5.56	120.99	129.33
2	B	501	LRA	OAK-CAL-CAO	-5.14	100.91	110.16
2	B	501	LRA	OAK-CAJ-CAI	5.05	114.33	109.86
2	A	501	LRA	OAK-CAJ-CAI	5.02	114.30	109.86
2	B	501	LRA	CAJ-CAI-CAH	-4.88	105.78	109.40
2	A	501	LRA	CAO-CAL-CAH	4.10	122.64	115.24
2	A	501	LRA	CAJ-CAI-CAH	-3.78	106.59	109.40
2	A	501	LRA	CAT-CAO-CAL	3.03	125.29	120.13
2	A	501	LRA	CAP-CAO-CAL	-2.80	116.03	120.50
2	A	501	LRA	CAE-CAF-CAH	-2.71	117.39	120.91
2	A	501	LRA	OAK-CAJ-OAN	2.57	124.65	121.27
2	B	501	LRA	OAK-CAJ-OAN	2.56	124.64	121.27
2	B	501	LRA	BRAG-CAC-CAD	2.18	122.27	119.30

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	501	LRA	CAA-CAF-CAH-CAL
2	A	501	LRA	CAH-CAL-CAO-CAT
2	A	501	LRA	CAH-CAL-CAO-CAP
2	B	501	LRA	CAH-CAL-CAO-CAT
2	B	501	LRA	CAH-CAL-CAO-CAP
2	A	501	LRA	CAE-CAF-CAH-CAL


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	501	LRA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	257/303 (84%)	0.64	19 (7%) 20 24	41, 56, 83, 100	0
1	B	228/303 (75%)	0.86	29 (12%) 8 9	43, 59, 82, 95	0
All	All	485/606 (80%)	0.74	48 (9%) 13 15	41, 58, 82, 100	0

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	363	PHE	5.7
1	B	473	TYR	4.4
1	A	256	MET	4.3
1	B	363	PHE	4.1
1	A	242	THR	3.9
1	B	238	THR	3.5
1	B	468	LEU	3.5
1	B	469	LEU	3.4
1	A	204	LEU	3.2
1	A	287	PHE	3.2
1	A	207	GLU	3.0
1	A	464	SER	3.0
1	B	311	LEU	2.8
1	B	219	TYR	2.8
1	A	241	THR	2.8
1	A	473	TYR	2.7
1	B	252	MET	2.7
1	B	360	PHE	2.6
1	B	250	TYR	2.6
1	A	461	THR	2.6
1	A	405	PRO	2.6
1	B	251	ASP	2.5
1	B	228	LEU	2.4
1	B	287	PHE	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	231	ALA	2.4
1	A	472	ILE	2.3
1	B	249	ILE	2.3
1	B	472	ILE	2.3
1	B	307	VAL	2.3
1	B	327	TYR	2.3
1	B	248	VAL	2.3
1	B	426	PRO	2.3
1	A	253	ASN	2.3
1	A	465	LEU	2.3
1	B	254	SER	2.2
1	A	362	ASP	2.2
1	A	467	PRO	2.1
1	B	343	GLU	2.1
1	B	258	GLY	2.1
1	B	452	LEU	2.1
1	B	422	LYS	2.1
1	A	203	GLN	2.1
1	B	247	PHE	2.1
1	B	342	SER	2.1
1	A	476	LEU	2.0
1	B	236	ILE	2.0
1	B	257	MET	2.0
1	A	402	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	LRA	A	501	21/21	0.83	0.19	85,87,93,134	0
2	LRA	B	501	21/21	0.92	0.11	61,69,75,88	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.